

Ladder Operators

Useful formula:

$$a = \sqrt{\frac{m\omega}{2\hbar}} \left(\hat{x} + \frac{i}{m\omega} \hat{p} \right)$$

$$a^\dagger = \sqrt{\frac{m\omega}{2\hbar}} \left(\hat{x} - \frac{i}{m\omega} \hat{p} \right)$$

This leads to the useful representation of \hat{x} and \hat{p} ,

$$\hat{x} = \sqrt{\frac{\hbar}{2m\omega}} (a^\dagger + a)$$

$$\hat{p} = i\sqrt{\frac{\hbar m\omega}{2}} (a^\dagger - a).$$

The Hamiltonian

The Hamiltonian for the one-vibration DDA system is

$$\hat{H} = \frac{P^2}{2M_s} + \sum_i \frac{M_s \Omega^2}{2} \left(R - \frac{c_i}{M_s \Omega^2} \right)^2 |i\rangle \langle i| + \sum_{ij} E_{ij} |i\rangle \langle j| + \hat{H}_B$$

$$\hat{H}_B = \sum_j \frac{M_B \omega_j^2}{2} \left(Q_j - c'_j \frac{R}{M_B \omega_j^2} \right)^2 + \sum_j \frac{P_j^2}{2M_B}$$

where $\hat{P} = -i\hbar \frac{\partial}{\partial R}$.

Transform the Hamiltonian into the mass-weighted coordinates, that is, $R \rightarrow \tilde{R} = \sqrt{M_s} R$ and $P \rightarrow \tilde{P} = P/\sqrt{M_s}$ (the same for P_j and R_j).

$$\hat{H} = \frac{\tilde{P}^2}{2} + \sum_i \frac{\Omega^2}{2} \left(\tilde{R} - \frac{c_i}{M_s^{1/2} \Omega^2} \right)^2 |i\rangle \langle i| + \sum_{ij} E_{ij} |i\rangle \langle j| + \hat{H}_{SB}$$

$$\hat{H}_B = \sum_j \frac{\omega_j^2}{2} \left(\tilde{Q}_j - c'_j \frac{\tilde{R}}{\sqrt{M_s} M_B \omega_j^2} \right)^2 + \sum_j \frac{\tilde{P}_j^2}{2}$$

Expand the first equation:

$$\hat{H} = \frac{\tilde{P}^2}{2} + \frac{\Omega^2 \tilde{R}^2}{2} + \sum_i \frac{c_i^2}{2M_s \Omega^2} |i\rangle \langle i| + \sum_{ij} E_{ij} |i\rangle \langle j| - \sum_i \frac{c_i}{\sqrt{M_s}} \tilde{R} |i\rangle \langle i| + \hat{H}_{SB}.$$

and move the oscillator terms to the second equation, we have

$$\hat{H} = \sum_i \frac{c_i^2}{2M_s \Omega^2} |i\rangle \langle i| + \sum_{ij} E_{ij} |i\rangle \langle j| - \sum_i \frac{c_i}{\sqrt{M_s}} \tilde{R} |i\rangle \langle i| + \hat{H}_{SB}. \quad (1)$$

Expand the second equation:

$$\frac{\tilde{P}^2}{2} + \left(\Omega^2 + \sum_j \frac{c_j'^2}{M_s M_B \omega_j^2} \right) \frac{\tilde{R}^2}{2} + \sum_j \frac{\omega_j^2 \tilde{Q}_j^2}{2} + \sum_j \frac{\tilde{P}_j^2}{2} - \sum_j \frac{c_j'}{\sqrt{M_s M_B}} \tilde{Q}_j \tilde{R}$$

Diagonalize the second equation:

$$\frac{\tilde{P}^2}{2} + \frac{\Omega^2 \tilde{R}^2}{2} + \sum_{j=1}^N \frac{\omega_j^2}{2} \left(\tilde{Q}_j - c_j' \frac{\tilde{R}}{\sqrt{M_s M_B \omega_j^2}} \right)^2 \rightarrow \sum_{j=1}^{N+1} \frac{\omega_j'}{2} \hat{X}_j$$

$$[\tilde{Q}_1 \quad \tilde{Q}_2 \quad \cdots \quad \tilde{Q}_{N-1} \quad \tilde{Q}_N \quad \tilde{R}] \begin{bmatrix} \frac{\omega_1^2}{2} & 0 & \cdots & 0 & \frac{c_1'}{2\sqrt{M_s M_B}} \\ 0 & \frac{\omega_2^2}{2} & \cdots & 0 & \frac{c_2'}{2\sqrt{M_s M_B}} \\ 0 & \vdots & \ddots & 0 & \vdots \\ 0 & 0 & \cdots & \frac{\omega_N^2}{2} & \frac{c_N'}{2\sqrt{M_s M_B}} \\ \frac{c_1'}{2\sqrt{M_s M_B}} & \frac{c_2'}{2\sqrt{M_s M_B}} & \cdots & \frac{c_N'}{2\sqrt{M_s M_B}} & (\Omega^2 + \sum_j \frac{c_j'^2}{M_s M_B \omega_j^2})/2 \end{bmatrix} \begin{bmatrix} \tilde{Q}_1 \\ \tilde{Q}_2 \\ \vdots \\ \tilde{Q}_{N-1} \\ \tilde{Q}_N \\ \tilde{R} \end{bmatrix}.$$

We obtain a set of frequencies $\text{diag}(\frac{\omega_j'^2}{2})$ and a unitary matrix U (U is returned by `numpy.linalg.eig()`):

$$\begin{bmatrix} \tilde{Q}'_1 \\ \tilde{Q}'_2 \\ \vdots \\ \tilde{Q}'_{N-1} \\ \tilde{Q}'_N \\ \tilde{Q}' \end{bmatrix} = U \begin{bmatrix} \tilde{Q}_1 \\ \tilde{Q}_2 \\ \vdots \\ \tilde{Q}_{N-1} \\ \tilde{Q}_N \\ \tilde{R} \end{bmatrix}.$$

so that

$$\tilde{R} = U^T [N+1, :] \begin{bmatrix} \tilde{Q}'_1 \\ \tilde{Q}'_2 \\ \vdots \\ \tilde{Q}'_{N-1} \\ \tilde{Q}'_N \\ \tilde{Q}' \end{bmatrix} = \sum_j U_{j,N+1} \tilde{Q}_j.$$

Substitute the expansion $\tilde{R} = \sum_j U_{j,N+1} \tilde{Q}_j$ into (1) we have

$$\hat{H} = \sum_{ij} E_{ij} |i\rangle \langle j| + \sum_i \frac{c_i^2}{2M_s \Omega^2} |i\rangle \langle i| - \sum_i \left(\sum_j U_{j,N+1} \frac{c_i}{\sqrt{M_s}} \right) \tilde{Q}_j |i\rangle \langle i| + \hat{H}_B.$$

Use the ladder operators¹ $\tilde{Q}_j = \sqrt{\frac{\hbar}{2\omega'_j}} (a_j^\dagger + a_j)$ we obtain

$$\hat{H} = \sum_{ij} E_{ij} |i\rangle \langle j| + \sum_i \frac{c_i^2}{2M_s \Omega^2} |i\rangle \langle i| - \sum_i \underbrace{\left(\sum_j U_{j,N+1} \sqrt{\frac{\hbar}{2M_s \omega'_j}} \right)}_{c_j} c_i |i\rangle \langle i| (a_j^\dagger + a_j) + \hat{H}_B. \quad (2)$$

$$= \sum_{ij} E_{ij} |i\rangle \langle j| + \sum_i \frac{c_i^2}{2M_s \Omega^2} |i\rangle \langle i| - \sum_{ij} c_i c_j |i\rangle \langle i| (a_j^\dagger + a_j) + \hat{H}_B \quad (3)$$

Initial Density Matrix, Shifting the Coordinate System

Suppose the initial density matrix for the vibrational modes is $\rho_B = \exp[-\beta(\frac{\tilde{P}^2}{2} + \frac{\Omega^2(\tilde{R}-\tilde{R}_0)^2}{2}) - \beta(\sum_j \frac{\omega_j^2}{2} (\tilde{Q}_j - c'_j \frac{\tilde{R}}{\sqrt{M_s M_B \omega_j^2}})^2 + \sum_j \frac{\tilde{P}_j^2}{2})]$. We set $\tilde{\mathcal{R}} = \tilde{R} - \tilde{R}_0$ ($\tilde{R} = \tilde{\mathcal{R}} + \tilde{R}_0$) and put it back to the density matrix to get

$$\rho_B = \exp[-\beta(\frac{\tilde{P}^2}{2} + \frac{\Omega^2 \tilde{\mathcal{R}}^2}{2}) - \beta(\sum_j \frac{\omega_j^2}{2} (\tilde{Q}_j - c'_j \frac{\tilde{\mathcal{R}} + \tilde{R}_0}{\sqrt{M_s M_B \omega_j^2}})^2 + \sum_j \frac{\tilde{P}_j^2}{2})]$$

and further we define $\tilde{\mathcal{Q}}_j = \tilde{Q}_j - \frac{\tilde{\mathcal{R}}}{\sqrt{M_s M_B \omega_j^2}}$ ($\tilde{Q}_j = \tilde{\mathcal{Q}}_j + \frac{\tilde{\mathcal{R}}}{\sqrt{M_s M_B \omega_j^2}}$) then we have

$$\rho_B = \exp[-\beta(\frac{\tilde{P}^2}{2} + \frac{\Omega^2 \tilde{\mathcal{R}}^2}{2}) - \beta(\sum_j \frac{\omega_j^2}{2} (\tilde{\mathcal{Q}}_j - c'_j \frac{\tilde{\mathcal{R}} + \tilde{R}_0}{\sqrt{M_s M_B \omega_j^2}})^2 + \sum_j \frac{\tilde{P}_j^2}{2})]$$

where $H_R = \frac{\tilde{P}^2}{2} + \frac{\Omega^2 \tilde{\mathcal{R}}^2}{2} + \sum_j \frac{\omega_j^2}{2} (\tilde{\mathcal{Q}}_j - c'_j \frac{\tilde{\mathcal{R}} + \tilde{R}_0}{\sqrt{M_s M_B \omega_j^2}})^2 + \sum_j \frac{\tilde{P}_j^2}{2}$.

Accordingly, the Hamiltonian becomes

$$\begin{aligned} \hat{H} &= \frac{\tilde{P}^2}{2} + \frac{\Omega^2 \tilde{\mathcal{R}}^2}{2} + \sum_i \frac{\Omega^2}{2} \left(\tilde{R}_0 - \frac{c_i}{M_s^{1/2} \Omega^2} \right)^2 |i\rangle \langle i| + \sum_i \Omega^2 \left(\tilde{R}_0 - \frac{c_i}{M_s^{1/2} \Omega^2} \right) \tilde{\mathcal{R}} |i\rangle \langle i| + \sum_{ij} E_{ij} |i\rangle \langle j| + \hat{H}_{SB} \\ \hat{H}_B &= \sum_j \frac{\omega_j^2}{2} \left(\tilde{\mathcal{Q}}_j^2 + (c'_j \frac{\tilde{\mathcal{R}} + \tilde{R}_0}{\sqrt{M_s M_B \omega_j^2}})^2 - 2\tilde{\mathcal{Q}}_j c'_j \frac{\tilde{\mathcal{R}} + \tilde{R}_0}{\sqrt{M_s M_B \omega_j^2}} \right) + \sum_j \frac{\tilde{P}_j^2}{2} \end{aligned}$$

¹ \tilde{Q}_j 's are mass-weighted coordinates so there's no mass in this equation.