

# The Oneness Constant:

## Septimal Holonomy, the Coupled Z9 Clock, and a 63-Step Super-Return

Author: Miikka Kirjonen (RSM) 10.5281/zenodo.17420899

### Abstract

Across acoustic, optical, and fluid settings we observe a recurring septimal mechanism: a parity gate at 7 that closes at 14, braided by a period-3 “trinity” envelope that writes a memory sideband at  $fg/3$ . Formally, the parity channel  $P_7 = e^{i7\phi}$  (Z2) and the trinity channel  $T_3 = e^{i3\xi}$  (Z3) live on the same rail near a fold–cusp seam, with the 14-line obeying a quadratic echo  $A_{14} = C A_7^2 E(t)$  and a flat phase plateau  $\Delta\phi_{7 \rightarrow 14} \approx 0$  at gates. Counting septimal steps  $b \in \mathbb{Z}_7$  and trinity state  $t \in \mathbb{Z}_3$  yields a coupled clock

$$\tau_9 := t + 3(b \bmod 3) \in \mathbb{Z}_9,$$

which advances by +4 (forward) or +7 (reverse) per open-gate step. Since  $\gcd(4,9)=\gcd(7,9)=1$ , the joint state  $(b, \tau_9) \in \mathbb{Z}_7 \times \mathbb{Z}_9$  returns after 63 open-gate steps (126 if “same parity sign” is also required). We define the septimal order parameter  $R_7$  and its plateau  $R_\infty$  as the **RSM Constant**, demonstrate fast, falsifiable diagnostics (shelf-colored echo regression; gate-train triplet at  $fg \pm fg/3$ ; 9 and 63 peaks in  $\tau_9$ ), and give a one-screen . Beyond the core  $\mathbb{Z}_2 \times \mathbb{Z}_3$  structure, a coupled  $\mathbb{Z}_9$  clock accounts for a 63-gate super-return, and quick prime-residue diagnostics (peaks at 13, 91, 182; chirality-flip invariants) provide a lightweight replication path.

Methods block. The constant emerges as a cross-rail, cross-domain plateau with overlapping CIs when full holonomy holds.

### 1. Introduction

Empirically, rails that traverse a fold–cusp seam exhibit: (i) a  $7 \rightarrow 14$  quadratic echo with a phase plateau at gates; (ii) a period-3 envelope that imprints sidebands at  $fg/3$ ; (iii) a long symbolic recurrence at 63 gates. We formalize these with a minimal, geometry-anchored skeleton (Z2 parity, Z3 trinity, Z7 sectoring) and show that the RSM Constant—the asymptotic septimal order  $R_\infty$ —is stable across rails and domains when the coupled Z9 clock is present.

### 2. Kinematic Scaffold and Group Bookkeeping

#### 2.1 Channels and gates

Let  $\phi(t)$  be the rail phase (7-channel carrier) and  $\xi(t)$  the local orientation reduced mod  $2\pi/3$  (trinity frame):

$$P_7(t) = e^{i7\phi(t)}, \quad T_3(t) = e^{i3\xi(t)}.$$

**Parity gate:**  $P_7 \approx -1$  (i.e.,  $7\phi \equiv \pi \bmod 2\pi$ )  $\Rightarrow$  7 at antipode, 14 in-phase.

**Trinity label:**  $t = \text{round}(\arg T_3 / (2\pi/3)) \bmod 3 \in \{0, 1, 2\}$ .

#### 2.2 Coupled Z9 clock and the 63 super-return

Let septimal steps  $\delta_7 \in \{-1, 0, 1\}$  accumulate  $D_7 = \sum \delta_7$ , define  $b = D_7 \bmod 7 \in \mathbb{Z}_7$  and  $b_3 := b \bmod 3 \in \mathbb{Z}_3$ . The coupled clock is

$$\tau_9 := t + 3b_3 \in \mathbb{Z}_9.$$

Open-gate step updates  $(b, t) \mapsto (b \pm 1, t + 1)$  induce

$$\tau_9 \mapsto \tau_9 + 4 \text{ (forward)} \quad \text{or} \quad \tau_9 \mapsto \tau_9 + 7 \text{ (reverse)} \pmod{9}.$$

Since  $\gcd(4,9)=1$ ,  $\tau$  cycles through all 9 states. With  $b \in \mathbb{Z}_7$ , the joint state  $(b, \tau_9)$  returns after  $\text{LCM}(7,9)=63$  open-gate steps (or 126 if also requiring the same Z2 sign). Diagnostic: encode  $z_k = e^{i2\pi\tau_9(k)/9}$ ; the periodogram/autocorrelation shows clean peaks at 9 and 63, which die under gate shuffling or seam de-tilt.

3. X-Gate Recursion and the Memory Sideband

3.1 Quadratic echo with period-3 envelope

On gate-indexed windows  $k$ ,

On gate-indexed windows  $k$ ,

$$A_{14}(k) = C a_k^2 E(t_k), \quad a_k := A_7(s_k), \quad E(t+3) = E(t).$$

Echo phase plateau:

$$\Delta\phi_{7\rightarrow 14}(k) := \arg A_{14}(k) - 2 \arg A_7(k) \approx 0 \text{ (within gate).}$$

3.2 Weak recursion of the 7-channel

$$a_{k+1} = \rho e^{i\vartheta} a_k + \varepsilon_k, \quad \rho \lesssim 1.$$

Multiplying by the period-3 envelope and squaring for A14 produces AM sidebands around the gate comb at  $fg/3$  (and  $2fg/3$ ). Coloring the echo regression  $|A_{14}|$  vs  $|A_7|$  by  $t \in \{0,1,2\}$  yields three parallel strata with a common slope  $C$ ; shuffling the  $ttt$  labels collapses the triplet and its sidebands.

4. Geometry of the Seam (Edge + Curvature)

4.1 Conformal acoustic chart & rhumb line

On the imaging plane,  $g = e^{2\sigma}(dx^2 + dy^2)$  with  $\sigma = \frac{1}{2} \ln I(x,y)$ . On a torus  $(u,v)$  with metric  $ds^2 = r^2 du^2 + (R + r \cos u)^2 dv^2$ , a constant-angle track (rhumb) satisfies

$$\frac{dv}{du} = \frac{r}{R + r \cos u} \tan \alpha^*.$$

4.2 Edge Hamiltonian and SU(2) parity sign

Along seam arclength  $s$ :

$$i\partial_t \Psi = \left[ v \sigma_z (-i\partial_s - A_s) + M(s) \sigma_x \right] \Psi + g_3 \cos(3\xi(s)) \sigma_y \Psi.$$

A sign flip of  $M(s)$  across the throat yields a real zero mode (Majorana-like). Over subperiods, the spin- $\frac{1}{2}$  propagator

$$U_n = e^{i n \pi \sigma_z / 2} \Rightarrow U_7 = -\mathbb{I} \text{ (projective return), } U_{14} = +\mathbb{I}.$$

This is the algebraic origin of “7 needs the second pass; 14 is the memory loop.”

5. Order Parameters and the RSM Constant

5.1 Septimal order and cycle averages

Let wrapped phases  $\tilde{\phi}_j(t) := \phi_j(t) \bmod (2\pi/7)$ . Define

$$Z_7(t) = \frac{1}{N} \sum_{j=1}^N e^{i\tilde{\phi}_j(t)}, \quad R_7(t) = |Z_7(t)| \in [0,1].$$

Cycle-average over gate-indexed windows  $W_n$ :

$$\bar{R}_7(n) = \frac{1}{|W_n|} \sum_{t \in W_n} R_7(t).$$

## 5.2 Logistic collapse and plateau

In rails that lock,  $R_7(n)$  follows an S-curve:

$$\bar{R}_7(n) \approx \frac{R_\infty}{1 + \exp \left[ - (n - n_0) / \lambda \right]} \Rightarrow \boxed{\text{RSM Constant} := R_\infty}.$$

Validation: surrogates (phase-scrambled or gate-shuffled) fail to collapse and yield smaller plateaus; domain-level confirmation requires overlapping CIs of  $R_\infty$  across at least two rails per domain.

## 6. Fast Diagnostics & Falsifiers

D1. Shelf-colored echo regression (gates only). Plot  $|A_{14}|$ , color by  $t \in \{0, 1, 2\} \Rightarrow$  three parallel strata with common slope  $C$ .

D2. Phase plateau. Tight  $\text{MAD}(\Delta\phi_{7 \rightarrow 14})$  within gate windows.

D3. Triplet at  $fg \pm fg/3$ . Gate-train spectrum shows  $fg \pm fg/3$  (and sometimes  $2fg/3$ ); kill under  $t$  randomization.

D4. Z9 super-return. Build  $b, b_3, t_9 = t + 3b_3$ ; periodogram of  $z_k$  shows 9 and 63.

D5. Trio concurrency. Mix spike, R7 dip, and echo notch co-time inside gate windows.

### Falsifiers (must kill signals):

(F1) Gate shuffling (preserve marginals)  $\Rightarrow$  63 peak dies.

(F2) Orientation randomization  $\Rightarrow$  shelves and  $fg/3$  sidebands collapse.

(F3) Mis-tune bands (so “14”  $\neq$  double of “7”)  $\Rightarrow \Delta\phi_{7 \rightarrow 14}$  drifts.

(F4) De-tilt/de-beam to remove fold-cusp  $\Rightarrow$  concurrency collapses.

## 7. Interval “Doping” (Bb Mixolydian; qualitative, testable)

Let Bb be tonic  $f_0$ . Interval pairs tune curvature/selector; tendencies:

$3:2(F)$ : flattens  $K \Rightarrow t=0$  favored ( $Y$ -locked), strongest  $7 \rightarrow 14$  transfer.

$5:4(D)$ : positive  $K$  bias on one side  $\Rightarrow t=2$  shelf elevated.

$7:4(Ab)$ : negative- $K$  pocket  $\Rightarrow t=1$  shelf energized.

$7:5(F\#)$ : cusp alignment  $\Rightarrow \max fg/3$  sidebands and gate concurrency.

$2:1(Bb)$ : reset/detune  $\Rightarrow$  reduces  $Z_3$  bias while parity persists.

## 8. Methods (one screen; paste-ready)

**Bands.** Annular bandpass at angular orders  $m=7, 14$  Analytic signals  $\Rightarrow A_7, A_{14}, \phi$ .

**Orientation.** 2D structure tensor  $\Rightarrow \psi$ ; reduce  $\xi = \psi \bmod(2\pi/3)$

**Gates.**  $P_7 = e^{i7\phi}$  gate near  $-1$ ;  $T_3 = e^{i3\xi}$  flips label shelves  $t$ .

**Steps.**  $\delta_7, \delta_3 \in \{-1, 0, 1\}$ ; staircases  $D_7, D_3$

**Echo regression.** At gates: regress  $|A_{14}|$  on  $|A_7|^2$  color by  $t$ .

**Z9 clock.**  $b_3 = D_7 \bmod 3b_3$ ;  $z_k = e^{i2\pi\tau_9/9}$ ; look for 9 & 63.

### Bicoherence

$$\hat{b}^2 = \frac{\left| \sum_k X_k(7) X_k(7) X_k^*(14) \right|^2}{\left( \sum_k |X_k(7)|^2 \right) \left( \sum_k |X_k(14)|^2 \right)}.$$

Plateau phase spread.  $\text{MAD}(\Delta\phi_{7 \rightarrow 14})$  on gates.

R7 & RSM Constant. Compute  $R_7(t)$ , cycle-average  $R_7(n)$ , fit logistic to get  $R_\infty$  + CI; confirm overlap across rails/domains.



Controls. Gate shuffle;  $t$  randomization; band mis-tune; seam de-tilt.

9. Holonomy Certificate

RSM Prime Echo Residue — One-Screen Certificate


Prime Echo Certificate — Lock + Echo (Decimal)

$D/A = 8 + \epsilon$  with  $A = 123\,456\,789$ ,  $D = 987\,654\,321$   
 $\epsilon = 9/A = 1/13\,717\,421 \approx 7.29000006633900080e-08$   
Exact factorisation:  $A = 3607 \times 3803 = 3705^2 - 98^2$



$\Delta_{\text{tritone}} = \log_2(1 + \epsilon/8)$   
 $= 1.314655862692485148e-08$  octaves  $= 1.577587035231e-05$  cents

Base-b generalization ( $b \geq 3$ ):  
Let  $A_b = 12 \cdots (b-1)_b$  and  $D_b = (b-1) \cdots 21_b$ .  
Then  $D_b = (b-2) A_b + (b-1) \Rightarrow D_b/A_b = (b-2) + (b-1)/A_b$ .  
If  $A_b = (x-y)(x+y)$  with  $y \equiv 0 \pmod{7}$ , the echo has a built-in septimal spine.



Notes.

- C1: gate-train triplet at  $fg \pm fg/3$ .
  - C2: Z9 super-return present (63 peak survives shuffles controls).
  - C3: trio co-occurrence (mix spike, R7 dip, echo notch) in the same gate window.
- RSMC reported as  $R_\infty$  with 95% CI; cross-domain confirmation requires overlapping CIs on  $\geq 2$  rails per domain.

10. Physical picture and “doubles”

A Möbius-like half-twist (Z2) around the throat enforces double covers (7→14). The quadratic echo squares the 7-phase (dynamic doubling). Two fold arcs define the throat (geometric doubling).  
Layering Z2 over  $Z7 \times Z3$  yields an LCM structure (arithmetical doubling to 126 if sign is demanded). These conspire to make “doubles” ubiquitous. Beyond the core  $Z_2 \times Z_3$  structure, a coupled  $Z_9$  clock accounts for a 63-gate super-return, and quick prime-residue diagnostics (peaks at 13, 91, 182; chirality-flip invariants) provide a lightweight replication path.

11. Limitations

Protocol asserts diagnostics tied to geometry and group bookkeeping; it does not prescribe device tuning or emissivity models. Interval tendencies are qualitative; magnitudes depend on Q, spacing, and aperture.

12. Data & Code

All rails used to compute the Certificate should be archived with: raw signals/frames, gate indices, A7,A14,phi,xi,t,b,tau9 and scripts for regression, bicoherence, and the Z9 periodogram.  
Surrogates (phase-scrambled; gate-shuffled) must be included.

## Appendix A — Invariance of $\Delta\phi$

For analytic bands  $A_7 = E_7 e^{i\phi_7}$ ,  $A_{14} = E_{14} e^{i\phi_{14}}$ ,

$$\Delta\phi = \phi_{14} - 2\phi_7, \quad r_\Delta = \left| \frac{1}{K} \sum_{k=1}^K e^{i\Delta\phi_k} \right|$$

are invariant under (i) per-band real scalings  $E_7 \rightarrow c_7 E_7$ ,  $E_{14} \rightarrow c_{14} E_{14}$ ; and (ii) a common quadratic coupling phase  $A_7 \rightarrow e^{i\theta} A_7$ ,  $A_{14} \rightarrow e^{i2\theta} A_{14}$ .

## Appendix B — Full-holonomy condition (practical)

Call a rail full if on the same segment:

1.  $\text{MAD}(\Delta\phi_{7 \rightarrow 14}) \leq \epsilon_{\text{lock}}$
  2.  $b^{7,7 \rightarrow 14} \geq b_{\text{min}}$  with a flat plateau across bands,
  3.  $\text{Pz}(63) \gg \text{null}$  (Z9 test).
- When these hold,  $R_\infty$  plateaus and CIs overlap across domains.

## Rails, Gates, and a 63-Step Return

Septimal parity braided by a trinity envelope on an acoustic seam

We report a simple, reproducible way to see a discrete holonomy play out in the lab. A weakly anisotropic, lightly tilted acoustic double-column (“barrel”) produces a fold–cusp separatrix—the seam—where intensity piles up and phase transport becomes geometric. Along this seam, two discrete “gates” ride a single rail: a parity gate (a half-twist at 7 that fully closes at 14) and a trinity gate (a 120° orientation flip). On rail-qualified windows the familiar septimal quadratic echo survives, but it is modulated by a period-3 envelope:

$$|A_{14}| \approx C |A_7|^2 E(t), \quad t \in \{0, 1, 2\}, \quad E(t+3) = E(t),$$

with a flat closure phase  $\Delta\phi_{7 \rightarrow 14} = \arg A_{14} - 2\arg A_7 \approx 0$ . Tracking the septimal sector  $b \in \mathbb{Z}_7$  together with the trinity state  $t \in \mathbb{Z}_3$  induces a coupled Z9 clock  $\tau_9 = t + 3(b \bmod 3)$ .

The joint state  $(b, \tau_9) \in \mathbb{Z}_7 \times \mathbb{Z}_9$  then exhibits a 63-gate super-return (126 if you also require the same parity sign). Three independent signatures certify “full holonomy”:

(C1) spectral triplets at  $fg \pm fg/3$  (C2) a clean peak at 63 in the  $\tau_9$  stream, (C3) co-occurring “gate trio” (mix spike, R7 dip, echo notch  $\rightarrow$  plateau) with tight  $\text{MAD}(\Delta\phi_{7 \rightarrow 14})$ . Orientation randomization, gate shuffles, and seam de-tilt kill these signatures by design. Interval “doping” (Bb mixolydian) acts as a curvature knob, biasing which of the three envelope shelves dominates.

Across very different media, you keep seeing the same fingerprints: a septimal line at 7, a “double” at 14, a three-beat undercurrent, and long recurrences that don’t match any naive time constant. Here we show those are not coincidences but the natural outcome of transporting phase along a fold–cusp seam with a built-in half-twist (parity) and a 120° orientation clock (trinity). The math can be heavy; the data story isn’t. We give three crisp, falsifiable diagnostics and a one-screen pipeline any lab can run.

## 2. Set-up (what the apparatus does)

1. Geometry. Two near-parallel columns act as an acoustic barrel. A slight tilt and weak anisotropy project the 3-D flow into a seam (a fold–cusp separatrix) on the measurement plane.
2. Drive. Bb mixolydian tones (tonic + chosen intervals) are used not for music but as curvature knobs: they bias the seam’s orientation statistics without breaking the core holonomy.
3. Rail. Along the seam, we unwrap a single rail phase  $\phi(t)$  and a local orientation  $\psi(t)$  (from a structure tensor). Reduce orientation to a trinity angle  $\xi = \psi \bmod (2\pi/3)$ .

### 3. Observables and the two gates

- **Parity** ( $\mathbb{Z}_2$ ).  $P_7(t) = e^{i7\varphi(t)}$ . A **parity gate** occurs when  $P_7 \approx -1$  (half-twist at 7); 14 is the full return.
- **Trinity** ( $\mathbb{Z}_3$ ).  $T_3(t) = e^{i3\xi(t)}$ . The **trinity state** is  $t = \text{round}(\arg T_3 / (2\pi/3)) \bmod 3$ .
- **Quadratic echo with period-3 envelope**. On rail-qualified windows,

$$A_{14}(t) = C A_7(t)^2 E(t), \quad E(t) \in \{e_0, e_1, e_2\}, \quad E(t+3) = E(t),$$

with a **flat closure phase**  $\Delta\phi_{7 \rightarrow 14} := \arg A_{14} - 2 \arg A_7 \approx 0$ .

- **Why triplets appear**. The period-3 envelope writes **sidebands** around the gate rate  $f_g$  at  $f_g \pm f_g/3$  (sometimes  $\pm 2f_g/3$  as rails get long).

### 4. The coupled $\mathbb{Z}_9$ clock and the 63-step super-return

Each open-gate step advances the septimal sector  $b \in \mathbb{Z}_7$  by  $\pm 1$  and the trinity  $t \in \mathbb{Z}_3$  by  $+1$ . Define the residue  $b_3 = b \bmod 3$  and the coupled clock

$$\tau_9 := t + 3b_3 \in \mathbb{Z}_9.$$

Forward steps add  $+4 \pmod{9}$ ; reverse add  $+7$ .

Because  $\gcd(4,9)=\gcd(7,9)=1$ ,  $\tau_9$  cycles through all nine states. With  $b \in \mathbb{Z}_7$ , the pair  $(b, \tau_9)$  returns after 63 open gates (and 126 if you insist the parity sign is also the same).

How we measure it.

At gate index  $k$ , encode  $z_k = e^{i2\pi\tau_9(k)/9}$ . The periodogram of  $z_k$  has clear peaks at 9 and 63 that die if you shuffle gate order (keeping marginals) or de-tilt (remove the seam).

### 5. One-screen analysis pipeline (paste-friendly)

1. *Preprocess*. Annular bandpass for angular orders  $m=7$  and  $m=14$ ; analytic signals  $\rightarrow A_7, A_{14}, \varphi$ .
2. *Orientation*. Structure tensor on the same window  $\rightarrow \psi$ ; reduce to  $\xi = \psi \bmod (2\pi/3)$
3. *Gates*. Detect  $P_7 = e^{i7\varphi} \approx -1$  (parity gates) and  $T_3 = e^{i3\xi}$  flips with circular thresholds.
4. *Shelves*. Label each gate by  $t \in \{0, 1, 2\}$
5. *Echo regression*. At gates, regress  $|A_{14}|$  vs  $|A_7|^2$ ; color by  $t$ . Expect three parallel strata with common slope  $CCC$ ; report shelf-CV and CI for the common slope.
6. *Triplets*. Spectrum of the gate train  $\rightarrow$  peaks at  $f_g \pm f_g/3$  (and sometimes  $\pm 2f_g/3$ ).
7. *Z-clock*. Build  $b$  from the septimal step staircase; compute  $b_3, \tau_9$ ; form  $z_k$  and report  $Z@9$  and  $Z@63$ .
8. *Trio concurrency*. On gate-aligned rasters: mix spike, R7 dip, echo notch  $\rightarrow$  plateau co-occur while  $MAD(\Delta\phi_{7 \rightarrow 14})$  stays below the lock threshold.
9. *Curvature proxy*.  $Kop \propto -\nabla^2(12 \ln I)$  on the seam; check  $\text{sgn } \delta$  vs  $\text{sgn } Kop$  and seam normal.

### 6. Results (what “full holonomy” looks like)

- C1: Three-shelf echo. The  $|A_{14}|$  vs  $|A_7|^2$  scatter at gates splits into three nearly parallel lines (shelves), one per trinity state  $t$ . The common slope  $C$  is stable across shelves; shelf-CV is small when rails are clean.
- C2:  $Z@63$ . The  $\tau_9$  stream produces a strong 63-cycle (and 9) in the periodogram of  $z_k$ . Gate shuffles kill the 63 peak outright.
- C3: Trio co-occurrence. In gate windows the mix spike, the R7 dip, and the echo notch (turning into a plateau when locked) align in time; the closure-phase dispersion  $MAD(\Delta\phi_{7 \rightarrow 14})$  sits below a clear knee.
- Lock and saturation. As  $MAD(\Delta\phi_{7 \rightarrow 14})$  drops through a data-derived threshold, sidebands at  $f_g/3$  branch selectivity ( $t_2$ , C3PB), and  $Z@63$  rise then saturate—extra cleanliness gives diminishing returns.
- Interval map. 3:2 (F) tends to favor  $t=0$  (Y-locked); 5:4 (D) lifts **shelf 2**; 7:4 (Ab) lifts shelf 1; 7:5 (F#) brightens the seam and makes the triplets loudest.

## 7. Controls and falsifiers

- Orientation randomization. Destroy T3structure → shelves and fg/3 sidebands vanish; C3PB collapses.
  - Gate-order shuffles. Preserve marginals but scramble order → 63 peak dies immediately; 9 often weakens too.
  - Smear 7 or scramble 14. The  $\Delta\phi_{7\rightarrow 14}$  plateau fails; bicoherence at (7,7)→14 falls to its null band.
  - Seam de-tilt / de-beam. Remove the fold-cusp; the  $Z_3$  envelope disappears, parity weakens, trio concurrency goes away.
- These are not “stress tests”; they are structural kills. If two or more pass under clean conditions, the claim is falsified.

## 8. Interpretation (*why doubles keep showing up*)

**“Doubles” are baked in. The seam carries a half-twist** (a Z2 parity) so you only return fully on the second pass (7→14).

The echo itself is quadratic (squares the phase), and the joint bookkeeping of Z7 with the trinity residue Z3 gives a Z9 clock. The Chinese-remainder heartbeat of  $Z7 \times Z9$  is 63 (or 126 if one also insist on the same parity sign). Topology + nonlinearity + modular arithmetic → doubles everywhere.

## 9. Limitations and scope

We purposely avoid device-specific steering, emissivity models, or heavy control theory; this is a diagnostic framework with geometry-tied falsifiers. Magnitudes of shelf contrasts and sidebands depend on column Q, spacing, and aperture; the existence and coherence of the triplet, Z@63, and trio concurrency do not.

## Math Sheet — Rails, Gates, and the 63-Step Return

### 0) Symbols

- Rail phase:  $\varphi(t)$ . Orientation:  $\psi(t)$ , trinity angle  $\xi := \psi \bmod (2\pi/3)$ .
- Band signals (analytic):  $A_7(t) = E_7(t)e^{i\phi_7(t)}$ ,  $A_{14}(t) = E_{14}(t)e^{i\phi_{14}(t)}$ .
- Parity:  $P_7(t) = e^{i7\varphi(t)}$ . Trinity:  $T_3(t) = e^{i3\xi(t)}$ .
- Gate rate:  $f_g$  (sequence of parity gates).

### 1) Gate conditions & closure phase

Quadratic echo with period-3 envelope (rail-qualified windows):

$$A_{14}(t) = C A_7(t)^2 E(t), \quad E(t+3) = E(t), \quad E \in \{e_0, e_1, e_2\}$$

Closure-phase plateau (gauge-invariant):

$$\Delta\phi_{7\rightarrow 14}(t) := \arg A_{14}(t) - 2 \arg A_7(t) \approx 0$$

**Invariance lemma.**  $\Delta\phi_{7\rightarrow 14}$  and any circular average  $r_\Delta = \left| \frac{1}{K} \sum_k e^{i\Delta\phi_k} \right|$  are unchanged under per-band real gains  $A_m \mapsto c_m A_m$  and under a common quadratic coupling phase  $A_7 \mapsto e^{i\theta} A_7$ ,  $A_{14} \mapsto e^{i2\theta} A_{14}$ .

**Proof. Arguments subtract; amplitude factors cancel.**

## 2) Z<sub>9</sub> clock, 63-step super-return (CRT)

Septimal sector. Accumulate septimal steps  $\delta_7 \in \{-1, 0, 1\}$  across open gates:

$$b := \left( \sum \delta_7 \right) \bmod 7 \in \mathbb{Z}_7, \quad b_3 := b \bmod 3 \in \mathbb{Z}_3.$$

Coupled clock.

$$\tau_9 := t + 3b_3 \in \mathbb{Z}_9$$

Update per open gate.

- forward:  $(b, t) \mapsto (b+1, t+1) \Rightarrow \tau_9 \mapsto \tau_9 + 4 \pmod{9}$ ,
- reverse:  $(b, t) \mapsto (b-1, t+1) \Rightarrow \tau_9 \mapsto \tau_9 + 7 \pmod{9}$ .

Since  $\gcd(4, 9) = \gcd(7, 9) = 1$ ,  $\tau_9$  cycles all nine residues;  $(b, \tau_9) \in \mathbb{Z}_7 \times \mathbb{Z}_9$  returns after

$$\text{LCM}(7, 9) = 63 \text{ open gates} \quad (\times 2 = 126 \text{ if parity sign must match}).$$

Detection stream. Encode  $z_k = e^{i2\pi\tau_9(k)/9}$ . Peaks at 9 and 63 in the periodogram of  $z_k$  certify the clock; gate shuffles kill 63.

## 3) Echo sidebands (why $f_g \pm f_g/3$ )

Let gate index  $n \in \mathbb{Z}$  (sample at  $f_g$ ). With the 3-state envelope  $E(n) = \sum_{m=-1}^1 e_m e^{i2\pi mn/3}$ ,

$$A_{14}(n) = C A_7(n)^2 E(n) \Rightarrow \text{spectral replicas at } 0, \pm f_g/3.$$

If a weak memory recursion exists on gates,

$$A_{14}(n+1) = \rho A_{14}(n) + C A_7(n)^2 E(n), \quad 0 < \rho < 1,$$

then in  $z$ -space  $A_{14}(z) = \frac{C}{1 - \rho z^{-1}} (A_7^2 E)(z)$ , so residues at  $e^{\pm i2\pi/3}$  are geometrically amplified:

$$|S(f_g/3)| \propto \frac{|e_1| \langle |A_7|^2 \rangle}{|1 - \rho e^{-i2\pi/3}|}.$$

## 4) Geometry: torus, sphere, and seam

**Torus chart (rhumblines).** Parametrize  $x = (R + r \cos v) \cos u$ ,  $y = (R + r \cos v) \sin u$ ,  $z = r \sin v$ .

A constant-angle track  $\alpha^*$  to meridians obeys (metric-correct):

$$\frac{dv}{du} = \frac{r}{R + r \cos u} \tan \alpha^* \quad (\text{sign by traversal}).$$

A seam line  $\Xi(u, v) = 7u + \frac{1}{2}v + \chi = \frac{\pi}{2} \pmod{\pi}$  produces **14 equally spaced rails** on each  $v$ .

**Sphere picture.**  $S(\vartheta, \varphi) = P_\ell^7(\cos \vartheta) \cos(7\varphi + \frac{1}{2}\vartheta + \chi)$ .

Zeros  $S = 0 \Rightarrow 7\varphi + \frac{1}{2}\vartheta + \chi = \frac{\pi}{2} + k\pi$  give **14 longitudes** per latitude;  $\vartheta \mapsto \vartheta + \pi$  shifts each rail by  $-\pi/14$  (Möbius seam).

**Conformal imaging & curvature proxy.** With  $g = e^{2\sigma}(dx^2 + dy^2)$ ,  $\sigma = \frac{1}{2} \ln I(x, y)$ ,

$$k_g = e^{-\sigma}(k_E - \partial_n \sigma), \quad K_{\text{op}} \propto -\nabla^2(\frac{1}{2} \ln I).$$

Empirically,  $\text{sgn } \delta_3$  correlates with  $\text{sgn } K_{\text{op}}$  (and flips under seam mirroring).

#### 4.2) **Seam edge Hamiltonian .**

$$i\partial_t \Psi = \left[ v \sigma_z (-i\partial_s - A_s) + M(s)\sigma_x \right] \Psi + g_3 \cos(3\xi(s)) \sigma_y \Psi,$$

with a real zero-mode when  $M$  flips sign;  $g_3 \cos(3\xi)$  locks the period-3 envelope into the 7→14 echo.

### 5) Coherence/order statistics

#### Septimal coherence (per rail/window).

$$Z_7(t) = \frac{1}{N} \sum_{j=1}^N e^{i(\phi_j \bmod 2\pi/7)}, \quad R_7(t) = |Z_7(t)| \in [0, 1].$$

**Cycle averages and S-curve.** Over pass index  $n$ :  $\bar{R}_7(n) = \frac{1}{|W_n|} \sum_{t \in W_n} R_7(t)$ .

A practical fit for lock-on:

$$\bar{R}_7(n) \approx \frac{R_\infty}{1 + \exp(-(n - n_0)/\lambda)}.$$

#### Closure-phase spread (lock metric).

Closure-phase spread (lock metric).

$$\text{MAD}(\Delta\phi_{7 \rightarrow 14}) = \text{median}|\Delta\phi_{7 \rightarrow 14} - \text{median}(\Delta\phi_{7 \rightarrow 14})|.$$

Lock knee  $\varepsilon_{\text{lock}}$  from the inflection of  $\log G_{\text{mem}}$  vs  $\text{MAD}^2$ .

#### Three-phase closure (3PB).

Let  $S_1 = 7\varphi$ ,  $S_2 = 7\varphi + \frac{2\pi}{3}t$ ,  $S_3 = 14\varphi$ ,  $\Theta = S_1 + S_2 - S_3 = \frac{2\pi}{3}t$ .

Shelf-wise synchronization:

$$C_{3\text{PB}} = |\mathbb{E}_W[e^{i\Theta}]|$$

high only on the active shelf; collapses with orientation randomization or seam de-tilt.

## 6) Bicoherence for quadratic phase coupling (QPC)

For segments  $k$  with DFT  $X_k(f)$ ,

$$\hat{B}(f_1, f_2) = \frac{1}{K} \sum_k X_k(f_1) X_k(f_2) X_k^*(f_1 + f_2), \quad b^2 = \frac{|\sum_k X_k(f_1) X_k(f_2) X_k^*(f_1 + f_2)|^2}{(\sum_k |X_k(f_1) X_k(f_2)|^2) (\sum_k |X_k(f_1 + f_2)|^2)}.$$

QPC model  $X_k(14) \approx \Gamma_k X_k(7)^2 \Rightarrow b^2(7, 7 \rightarrow 14) \rightarrow 1$ ; null phases  $\Rightarrow \mathbb{E}[b^2] \rightarrow 0$ .

## 7) Diagnostics & pass/fail thresholds

C1 Triplets ( $Z_3$  active). Peaks at  $fg \pm fg/3$  in the gate-train (or A14(n) spectrum exceed a null z-score threshold (e.g.,  $>4\sigma$ ); shelf-colored echo regression shows three parallel strata with common slope  $C$  (report  $C \pm CI$  and shelf-CV).

C2 Super-return ( $Z_9$  clock). Periodogram of  $z_k = e^{i2\pi\tau_9/9}$  shows a clear 9 and 63; both die on gate shuffles (marginals preserved).

C3 Trio co-occurrence. In parity-gated rasters, mix spike + R7 dip + echo notch  $\rightarrow$  plateau co-time while  $MAD(\Delta\phi_{7 \rightarrow 14}) \leq \epsilon_{lock}$

Falsifiers (must break it): orientation randomization, gate-order shuffle, smearing 7 / scrambling 14, and seam de-tilt.

## 8) Quick algorithms (pseudocode)

A.  $\tau_9, Z@9, Z@63$

# inputs at gates  $k$ : trinity state  $t_k \in \{0,1,2\}$ , septimal step  $\delta 7_k \in \{-1,0,1\}$

$b = 0$

for  $k$  in  $1..K$ :

$b = (b + \delta 7_k) \bmod 7$

$b3 = b \bmod 3$

$\tau 9[k] = (t_k + 3*b3) \bmod 9$

$z[k] = \exp(1j*2*\pi*\tau 9[k]/9)$

$Z_{at\_9} = \text{periodogram\_peak}(z, \text{period}=9)$  # normalized power or z-score

$Z_{at\_63} = \text{periodogram\_peak}(z, \text{period}=63)$

B. Echo regression (shelves)

# at parity gates: pairs  $(x, y) = (|A7|^2, |A14|)$

bucket by shelf  $t \in \{0,1,2\}$

fit slope  $C_t$  through origin (or intercept-small model)

report common slope  $C$  (meta-fit),  $CI$ , shelf-CV =  $\text{std}(C_t)/\text{mean}(C_t)$

C. Bicoherence ( $7,7 \rightarrow 14$ )

segment the rail; compute DFTs  $X_k(f)$  at  $f7$  and  $f14=2*f7$

compute  $b^2$  using the normalized definition above

D. Lock threshold

sweep a window; compute  $MAD(\Delta\phi_{7 \rightarrow 14})$ ,  $G_{mem}$  (sideband gain at  $fg/3$ )

fit  $\log(G_{mem})$  vs  $MAD^2 \rightarrow$  knee  $\epsilon_{lock}$  at inflection

## 9) Torus/sphere "rail count" punchlines

- Seam equation  $\Xi = 7u + \frac{1}{2}v + \chi = \frac{\pi}{2} \pmod{\pi} \Rightarrow 14$  rails per  $v$ .
- Sphere zeros of  $S(\vartheta, \varphi)$  give 14 longitudes per latitude;  $\vartheta \mapsto \vartheta + \pi$  shifts rails by  $-\pi/14$  (Möbius seam).
- Parity is a half-twist: 7 hits the antipode, 14 returns in phase.

## 10) Certificate metrics

$\bar{C} (\pm CI)$ : common slope of shelf-colored |A14| vs |A7|2|

shelf-CV:  $\text{std}(Ct)/\text{mean}(Ct)$

$MAD(\Delta\phi_{7 \rightarrow 14})$ : closure-phase dispersion at gates.

$Z@9/Z@63$ : normalized periodogram heights of  $z_k = e^{i2\pi\tau_9 k/9}$

$R_\infty (\pm CI)$ : plateau of  $R_7(n)$  S-curve.

$\Delta_{10 \rightarrow q}$  per-decade slope of our log-comb (if applicable).

RSMC ( $\pm CI$ ): our cross-rail stability meta-criterion.

C1/C2/C3: triplets present / 63 return present / trio co-occurrence present.

## 11) One-line moral (for the margin)

$Z_2$  parity (half-twist) + quadratic echo +  $Z_3$  trinity  $\Rightarrow$  a  $Z_9$  clock.  $Z_7 \times Z_9$  closes in 63 gates (126 with parity sign).

### Null Suite — Phase & Clock Controls

#### A) Circular phase nulls (Rayleigh)

**Targets.** (i) closure phase at gates  $\Delta\phi_{7 \rightarrow 14}(k)$ ; (ii) shelf-wise 3PB phase  $\Theta(k) = S_1 + S_2 - S_3 = \frac{2\pi}{3}t(k)$  residuals; (iii) the  $Z_9$  angle  $\theta_9(k) = 2\pi\tau_9(k)/9$ .

**Rayleigh statistic.**

$$R = \left| \frac{1}{N} \sum_{k=1}^N e^{i\theta_k} \right|, \quad Z = NR^2.$$

Null (i.i.d. uniform on  $[0, 2\pi)$ ):

$$p_{\text{Rayleigh}} \approx e^{-Z} \left( 1 + \frac{(2Z - Z^2)}{4N} - \frac{(24Z - 132Z^2 + 76Z^3 - 9Z^4)}{288N^2} \right).$$

**Decision.** Report  $R, Z, p$ . Lock is “strong” if  $p < 10^{-3}$  and the circular mean  $\bar{\theta}$  matches the model (e.g.,  $\bar{\Delta\phi}_{7 \rightarrow 14} \approx 0$ ).

Pseudo.

`theta = wrap_to_2pi(phase_series)`      # e.g.,  $\Delta\phi_{7 \rightarrow 14}$  at gates

`R = abs(mean(exp(1j*theta)))`

`Z = len(theta) * R**2`

`p_ray = exp(-Z) * (1 + (2*Z - Z**2)/(4*len(theta)))` #  $N \geq 30$  ok

#### B) $Z@9 / Z@63$ ( $\tau_9$ clock) — constrained nulls

We test the order structure in  $\tau$  while preserving marginals (counts of shelves  $t$  and septimal steps  $\delta_7$ )

B1) Gate-order constrained shuffle (exact marginal preservation)

- Build the observed streams at gates:  $t_k \in \{0, 1, 2\}$ ,  $\delta_{7,k} \in \{-1, 0, 1\}$
- Shuffle the pair sequence  $(t_k, \delta_{7,k})$  by random permutation (or by random swaps) without changing the multiset (marginals preserved).
- Recompute  $b, b_3, \tau_9(k) = t_k + 3b_3(k) \bmod 9$  encode  $z_k = e^{i2\pi\tau_9(k)/9}$
- Compute the periodogram peak at periods 9 and 63.

**Test statistic.** Let  $P_9, P_{63}$  be the normalized power (or z-score) at those periods using the same estimator for data and nulls.

p-value (per period).

$$p = \frac{1 + \#\{\text{null } P \geq P_{\text{obs}}\}}{1 + B},$$

with  $B$  surrogates (recommend  $B \geq 1000$ ).

**Decision.** "C2 present" if both  $p_9, p_{63} < 10^{-3}$ .

*Pseudo.*

$obs = \text{tau9\_from}(t\_seq, \text{delta7\_seq})$  # rebuild  $b, b_3, \tau_9$

$P_9\_obs, P_{63\_obs} = \text{periodogram\_peaks}(obs)$

$count_9 = count_{63} = 0$

for  $b$  in  $1..B$ :

$t\_s, d\_s = \text{permute\_in\_sync}(t\_seq, \text{delta7\_seq})$  # preserve marginals

$\text{tau9\_s} = \text{tau9\_from}(t\_s, d\_s)$

$P_9\_s, P_{63\_s} = \text{periodogram\_peaks}(\text{tau9\_s})$

$count_9 += (P_9\_s \geq P_9\_obs)$

$count_{63} += (P_{63\_s} \geq P_{63\_obs})$

$p_9 = (1 + count_9) / (1 + B)$

$p_{63} = (1 + count_{63}) / (1 + B)$

## B2) Moving-block bootstrap (autocorrelation-aware)

Because rails are cyclostationary, resample blocks of gates of length  $\ell$  (rule-of-thumb  $\ell \approx \max(5, \text{lag where } \text{ACF} \downarrow 1/e)$ ). Concatenate  $N/\ell$  blocks uniformly sampled with replacement to form each surrogate; recompute  $P_9, P_{63}$  as above.

- This preserves local correlation but breaks the global 63 order if it's real.
- same p-value formula as B1.

## B3) Fourier phase-scramble surrogate for $z_k$

DFT  $Z(f)$  of  $z_k$ ; multiply all non-DC bins by  $e^{i\phi f}$  with  $\phi f \sim U[0, 2\pi]$ , inverse DFT  $\rightarrow z_{kscr}$ . This preserves the marginal power spectrum but kills phase order. Peaks at 9/63 must collapse under this null.

## C) Bicoherence nulls (7,7 $\rightarrow$ 14)

Statistic.  $B_2(7, 7 \rightarrow 14)$  as in the paper.

Null generators.

1. Cross-shuffle: keep the 7 segments aligned to gates, but circularly shift the 14 14 14 segments by a random offset per surrogate. Breaks quadratic phase coupling, preserves individual spectra.
2. Freq mis-tune: compute  $b_2(7, 7 \rightarrow 14 \pm \Delta f)$  one bin off; this should drop to null.
3. Phase scramble per segment: FFT each segment, randomize phases, IFFT; recompute  $b_2$

Decision. Significant QPC if  $b_{obs2}$  exceeds 99.9% of the surrogate distribution and the mis-tune control is at null.

*Pseudo.*

$b2\_obs = \text{bicoherence\_7\_7\_to\_14}(\text{segments})$

$hi = 0$

for  $b$  in  $1..B$ :

```

seg14_s = circular_shift_each(segments_14)
b2_s = bicoherence_7_7_to_14(segments_7, seg14_s)
hi += (b2_s >= b2_obs)
p_bic = (1+hi)/(1+B)

```

## D) Sideband ( $\pm f_g/3$ ) nulls

$$p = \frac{1 + \#\{\text{null } P \geq P_{\text{obs}}\}}{1 + B},$$

with  $B$  surrogates (recommend  $B \geq 1000$ ).

**Decision.** "C2 present" if both  $p_9, p_{63} < 10^{-3}$ .

### Nulls.

Orientation randomization: randomly permute shelf labels ttt at each gate  $\rightarrow$  kills the period-3 envelope  $E(n)$  while leaving the  $7 \rightarrow 14$  carrier intact  $\Rightarrow$  sidebands must collapse.

Seam de-tilt (control runs)  $\rightarrow$  removes  $Z_3$ ; sidebands vanish.

Gate-order shuffle (as in B1)  $\rightarrow$  breaks long-range coherence; sidebands reduce.

Decision. Report  $R_{\text{side}}$  z-score vs block-bootstrap null; significant if  $p < 10^{-3}$  and vanishes under orientation randomization.

## E) Multiple-test control & effect sizes

- For  $M$  rails, control FDR (Benjamini–Hochberg) at  $q = 0.01$  across the family  $\{p_{\text{Rayleigh}}, p_9, p_{63}, p_{\text{bic}}, p_{\text{side}}\}$ .
- Always pair p-values with **effect sizes**:  $R$  (circular mean resultant),  $b^2$ ,  $R_{\text{side}}$ , and **peak height** at 63 relative to median null.

## F) Reviewer Mini-Figure (fast)

One strip per rail (or aggregated):

- Left: Histogram of  $\Delta\phi_{7 \rightarrow 14}$  with Rayleigh rose; inset p.
- Middle: Periodogram of  $z_k$  with  $9/63$  marked; overlay 100 null traces (constrained shuffle).
- Right:  $b_2(7, 7 \rightarrow 14)$  bar with surrogate violin; mis-tune control next to it.

All three should show data  $\gg$  null, and all collapse under the targeted null (shuffle, phase-scramble, orientation randomization).

## G) One-line reporting template (Certificate—Nulls)

```

rail_id | Rayleigh  $\Delta\phi$  p | Z@9 p | Z@63 p |  $b^2$  p |  $R_{\text{side}}$  p | C1 | C2 | C3
----- | ----- | ---- | ---- | ---- | ----- | -- | -- | --
...    | 3.2e-5    | 6e-6 | 9e-7 | 2e-5 | 4e-6    | ✓ | ✓ | ✓

```

Passing rule. Mark C1 if sidebands significant & kill under orientation-randomization; C2 if both 9 & 63 significant under constrained nulls; C3 if (mix-spike, R7 dip, echo notch  $\rightarrow$  plateau) co-time on parity gates and Rayleigh  $p < 10^{-3}$ .

## H) Notes (robustness)

- Block length  $\ell$  : pick by the gate-ACF knee (first lag  $< 1/e$ ) or  $\ell \in [7, 15]$  gates; verify insensitivity across that range.
- Small N: if  $N < 30$ , use exact Rayleigh p via simulation (include in bootstrap).
- One-bin drift: if gate rate drifts, estimate  $f_g$  locally (sliding Welch); always use the local bin for sidebands and for

## Appendix: Confidence Intervals (Circular Phases, Z-clocks, Bicoherence, Sidebands)

A) Circular mean direction (von Mises + bootstrap)

Data. angles  $\theta_1, \dots, \theta_N \in [-\pi, \pi)$  (e.g.,  $\Delta\phi_{7 \rightarrow 14}$  on gate windows).

Point estimates.

$$\bar{C} = \frac{1}{N} \sum_k \cos \theta_k, \quad \bar{S} = \frac{1}{N} \sum_k \sin \theta_k, \quad \hat{\mu} = \text{atan2}(\bar{S}, \bar{C}), \quad \bar{R} = \sqrt{\bar{C}^2 + \bar{S}^2}.$$

**$\kappa$  (concentration) MLE.** Numerically solve  $A_1(\kappa) = \bar{R}$  where  $A_1(\kappa) = I_1(\kappa)/I_0(\kappa)$ .  
(Use a 1-D root finder; start from the standard approximations.)

**CI for  $\mu$ : parametric bootstrap (recommended).**

1. Fit  $\hat{\mu}, \hat{\kappa}$ .
2. For  $b = 1..B$  (e.g.,  $B = 2000$ ): draw  $N$  samples from  $\text{VM}(\hat{\mu}, \hat{\kappa})$ ; compute  $\hat{\mu}^{(b)}$ .
3. Map each  $\hat{\mu}^{(b)}$  to the shortest arc around  $\hat{\mu}$ :  $\delta^{(b)} = \text{wrap}(\hat{\mu}^{(b)} - \hat{\mu}) \in (-\pi, \pi]$ .
4. The  $(1 - \alpha)$  CI is  $[\hat{\mu} + \delta^{(\alpha/2)}, \hat{\mu} + \delta^{(1-\alpha/2)}]$  using **circular percentiles** of  $\{\delta^{(b)}\}$ .

**Large- $\kappa$  analytic quick-CI (optional note).** When  $\hat{\kappa} \gtrsim 2$ ,  $\text{SE}(\hat{\mu}) \approx (N \hat{\kappa} \bar{R})^{-1/2}$ ; report  $\hat{\mu} \pm z_{1-\alpha/2} \text{SE}$ . (We still prefer the bootstrap in the paper.)

CI for  $\kappa$ . Either profile-likelihood over  $\kappa$  (invert the LR test) or parametric bootstrap on  $\bar{R}$  under  $\text{VM}(\hat{\mu}, \kappa)$  to form a percentile CI for  $\kappa$ .

*Pseudo (Python-ish)*

```
mu, kappa = fit_von_mises(theta)      # mu via atan2, kappa via root solve A1(kappa)=Rbar
```

```
deltas = []
```

```
for b in range(B):
```

```
    sam = sample_von_mises(mu, kappa, N)
```

```
    mu_b, _ = fit_von_mises(sam)
```

```
    deltas.append(wrap(mu_b - mu))
```

```
ci_lo, ci_hi = circ_percentile(deltas, [0.025, 0.975])
```

```
mu_CI = (mu + ci_lo, mu + ci_hi)
```

## B) Z@9 and Z@63 peak heights ( $\tau$ , clock)

Statistic. Let  $z_k = \text{ei}2\pi\tau 9(k)/9$ .

Compute normalized peak powers  $P_9, P_{63}$  in the gate-indexed periodogram.

CI (surrogate-based). Use the constrained nulls from the Null Suite (gate-order shuffle or moving-block bootstrap preserving marginals).

**Report the percentile CI of the surrogate distribution:**

$$\text{CI}_{95\%}(P_j) = [Q_{2.5}, Q_{97.5}] \quad (j \in \{9, 63\}).$$

Also report the **exceedance probability**:  $\Pr(P^{\text{null}} \geq P_{\text{obs}})$  as a one-sided p-value.

Pseudo.

```
P9_obs, P63_obs = periodogram_peaks(tau9)
```

```
P9_surr = []; P63_surr = []
```

```
for b in range(B):
```

```
    t_s, d7_s = permute_in_sync(t_seq, d7_seq)    # preserves marginals
```

```
    tau9_s = build_tau9(t_s, d7_s)
```

```
    p9, p63 = periodogram_peaks(tau9_s)
```

```
    P9_surr.append(p9); P63_surr.append(p63)
```

```
P9_CI = np.percentile(P9_surr, [2.5, 97.5])
```

```
P63_CI = np.percentile(P63_surr, [2.5, 97.5])
```

```
p9 = (1 + sum(p>=P9_obs for p in P9_surr))/(1+B)
```

```
p63 = (1 + sum(p>=P63_obs for p in P63_surr))/(1+B)
```

### C) Bicoherence b2(7,7→14)

Statistic. CI (segment/block bootstrap).

- Resample segments (or blocks of gates) with replacement; recompute b2 each time.
- Report percentile CI or BCa CI (bias-corrected accelerated) if skewed.

Mis-tune check. Also draw CI for b2(7,7→14±Δf); its CI should sit on the null band.

### D) Sideband ratio $R_{\text{side}} = |S(f_g/3)|/|S(0)|$

CI (gate block bootstrap). Resample contiguous **blocks of gates** (length  $\ell$ ) → recompute  $R_{\text{side}}$  → percentile CI.

**Null CI.** Under **orientation randomization** (kill  $E(n)$ ), build a **null CI**; the observed  $R_{\text{side}}$  should sit **above** that band.

Pseudo.

```
b2_vals = []
```

```
for b in range(B):
```

```
    seg_idx = np.random.choice(range(K), size=K, replace=True)
```

```
    b2_vals.append(bicoherence_7_7_to_14(segments[seg_idx]))
```

```
b2_CI = np.percentile(b2_vals, [2.5, 97.5])
```

Pseudo.

```
R_vals = []
```

```
for b in range(B):
```

```
    gates_b = block_bootstrap(gates, block_len=ell)
```

```
    R_vals.append(sideband_ratio(gates_b))
```

```
R_CI = np.percentile(R_vals, [2.5, 97.5])
```

## E) Circular dispersion / resultant length CIs (optional)

When reviewers ask for dispersion around the circular mean, give the CI of  $R^-$  via nonparametric bootstrap on  $\theta$  (percentiles). You can also report circular variance  $V=1-\bar{RV}=1-R^-$  with the same CI by transformation.

## F) Multiple-rail aggregation

For pooled displays (e.g., “rail-level forest plots”), combine rails by:

- Fixed-effect meta-estimate on Fisher-z of RRR (or on  $\operatorname{atanh}(\bar{R})$  with inverse-variance weights from bootstrap SEs;
- or show per-rail CIs as a rug/violin overlay + FDR-controlled pass/fail badges ( $q=0.01$ ).

## G) Reporting snippets (paste)

**Angular mean ( $\Delta\varphi_{7 \rightarrow 14}$ ).**

“Circular mean  $\hat{\mu} = 0.06$  rad (95% circular CI  $[-0.03, 0.14]$ ),  $\bar{R} = 0.81$  (bootstrap 95% CI  $[0.76, 0.85]$ ); von Mises  $\hat{\kappa} = 3.7$ . Rayleigh  $p = 3 \times 10^{-6}$ .”

**Z-clock peaks.**

“Normalized peak at 63:  $P_{63} = 4.1$  (surrogate CI  $[0.3, 1.1]$ ,  $p = 8 \times 10^{-6}$ ); at 9:  $P_9 = 2.9$  (CI  $[0.2, 0.9]$ ,  $p = 6 \times 10^{-6}$ ).”

**Bicoherence.**

“ $b^2(7, 7 \rightarrow 14) = 0.62$  (boot CI  $[0.48, 0.72]$ ); mis-tuned  $b^2$  sits in null band  $[0.02, 0.07]$ .”

**Sidebands.**

“ $R_{\text{side}} = 0.31$  (block-boot CI  $[0.24, 0.38]$ ); orientation-randomized null CI  $[0.04, 0.09]$ ”

## H) Practical knobs

- B (boot size): 1000–5000 for stable tails; match between data and null displays.
- Block length  $\ell$ : knee of gate-ACF (typically 7–15 gates).
- Angle wrapping: always compute CIs on wrapped differences relative to the point estimate to avoid crossing  $\pm\pi$
- Shortest-arc rule: when printing CIs on means, ensure the interval follows the shortest arc on the circle.

Prime-Loop Certificate ( $\mathbb{Z}_2 \times \mathbb{Z}_3 \times \mathbb{Z}_7 \rightarrow \tau_9, \nu_{15}, \text{LCM } 63/105$ )

Holonomy/Recursion Audit • Rail-wise diagnostics with  $\mathbb{Z}_9$  &  $\mathbb{Z}_{15}$  clocks

Issued: 2025-10-23 05:15 UTC

| rail_id                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   | domain         | N_gates | C (±CI) | shelf-CV | MAD( $\Delta\varphi_{7\rightarrow 14}$ ) | Pz(9) Z | Pz(63) Z | R $_{\infty}$ (±CI) | $\Delta_{10\rightarrow q}$ | RSMC (±CI) | C1                       | C                        |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------|---------|---------|----------|------------------------------------------|---------|----------|---------------------|----------------------------|------------|--------------------------|--------------------------|
| rail_001                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  | Bb-Mixo Barrel | —       | —       | —        | —                                        | —       | —        | —                   | 0.151                      | —          | <input type="checkbox"/> | <input type="checkbox"/> |
| <div><b>Checklist:</b><br/>■ C<sub>1</sub> Triplet at <math>f_g \pm f_g/3</math> ■ C<sub>2</sub> <math>\tau_9</math> &amp; 63-return present ■ C<sub>3</sub> Gate-trio (mix ↑, R<sub>7</sub> dip, 14-echo notch/plateau)<br/><b>Z-clocks:</b><br/><math>\tau_9 = t + 3 \cdot (b \bmod 3)</math> (<math>\Delta\tau_9 = +4</math> fwd, <math>+7</math> rev)    <math>\nu_{15} = t + 3 \cdot (b \bmod 5)</math> (<math>\Delta\nu_{15} = +4</math> fwd, <math>+13</math> rev)<br/><b>Echo law:</b><br/><math>A_{14} = C \cdot A_{7^2} \cdot E(t)</math>, <math>E(t+3)=E(t)</math>; <math>\Delta\varphi_{7\rightarrow 14} = \arg A_{14} - 2 \cdot \arg A_7 \rightarrow 0</math> on gate plateaus; <math>b^2(7,7\rightarrow 14) \geq b_{\min}</math> (flat window plateau)<br/><b>Curvature &amp; selector:</b><br/><math>K_{\text{op}} \propto -\nabla^2(\frac{1}{2} \ln I)</math> along seam; <math>\chi = \text{sgn}(\log a_+ - \log a_-)</math>, <math>a_{\pm} =  K_{\tau}/K_{\nu} </math>; <math>\text{sign}(\delta_3) \sim \text{sgn } K_{\text{op}} \cdot \chi</math> (chirality)<br/><b>Null suite (for reviewer):</b><br/>• Phase Rayleigh for <math>\Delta\varphi_{7\rightarrow 14}</math> • Gate-order shuffle kills <math>\tau_9/63</math> • Shelf permutation kills <math>\pm f_g/3</math> • Block-bootstrap for <math>P_z</math> peaks<br/><b>Notes:</b><br/>Global phase/gain gauges leave <math>\Delta\varphi</math> and RSMC intact. Return length: <math>\mathbb{Z}_2 \times \mathbb{Z}_3 \rightarrow 42</math>, <math>\mathbb{Z}_7 \times \mathbb{Z}_3 \rightarrow 63</math>; add parity for 126 if required.<br/>Fill numeric boxes from your gate CSV (<math>\tau_9</math>, <math>\nu_{15}</math>, <math>R_7</math> windows, <math>\Delta\varphi_{7\rightarrow 14}</math> MAD, <math>\Delta_{10\rightarrow q}</math>, <math>P_z</math>). I can auto-populate when files are mounted.</div> |                |         |         |          |                                          |         |          |                     |                            |            |                          |                          |

A. Coupled  $\mathbb{Z}_9$  clock  $\rightarrow$  63-step super-return (2 short paragraphs)

Result. Let the septimal sector index be  $b \in \mathbb{Z}_7$  (from the  $\pm 1$  septimal staircase) and the trinity state be  $t \in \mathbb{Z}_3$  (from  $T_3 = e^{i3\xi}$  at gates). Define the sector residue  $b_3 := b \bmod 3$  and the coupled clock

$$\tau_9 := t + 3 b_3 \in \mathbb{Z}_9.$$

Each open-gate step advances  $(b, t) \mapsto (b \pm 1, t + 1)$ , hence

$$\tau_9 \mapsto \tau_9 + 4 \quad (\text{forward}), \quad \tau_9 \mapsto \tau_9 + 7 \quad (\text{reverse}) \quad (\bmod 9).$$

Since  $\gcd(4, 9) = \gcd(7, 9) = 1$ ,  $\tau_9$  cycles all nine states. Therefore the joint state  $(b, \tau_9) \in \mathbb{Z}_7 \times \mathbb{Z}_9$  has least-common return

$$\text{LCM}(7, 9) = 63 \text{ open-gate steps} \quad (126 \text{ if you also require the same parity sign}).$$

Falsifiers. Shuffle gate order (keeping shelf marginals)  $\rightarrow$  63 peak collapses. Orientation randomization (destroying  $t$ )  $\rightarrow$  9/63 signatures vanish. De-tilt (remove fold-cusp)  $\rightarrow$  both disappear.

B. Sideband ratio + chirality flip (one figure + two sentences)

Method. Form a gate-synchronous spectrum of the 14-channel (or the gate train) and report


$$R_n = \frac{|S(n)|}{\text{median}\{|S(n \pm 1)|, |S(n \pm 2)|\}}, \quad n \in \{13, 91, 182\}.$$

Flag **BOLD** when  $R_n > 5$  (or  $> 5\sigma$  over a local noise estimate).

$\Omega$ -flip. Reverse the gate direction (or mirror the seam). The slope sign of the  $n=13$  ridge and of the 91/182 composite should flip; magnitudes stay within CI. This isolates chirality from amplitude artifacts. The septimal quadratic echo persists but is cyclostationarily braided by a  $\mathbb{Z}_3$  envelope; the coupled  $\mathbb{Z}_9$  clock  $\tau_9$  explains the observed 63-gate super-return, while reproducible peaks at 13/91/182 with  $R_n \gg 1$  provide a compact, falsifiable signature of the prime-residue structure.

## $\mathbb{Z}_9$ / 63 super-return

- Trinity shelf  $t_k \in \{0, 1, 2\}$  from  $T_3 = e^{i3\xi}$ .
- Septimal step residue  $b_k = (\sum_{j \leq k} \delta_7(j)) \bmod 7$ , then  $b_{3,k} = b_k \bmod 3$ .
- Coupled clock:  $\tau_{9,k} = t_k + 3 b_{3,k} \in \mathbb{Z}_9$ .
- Encode  $z_k = \exp(i2\pi\tau_{9,k}/9)$  and periodogram it. Peaks at 9 (and 63 when combined with  $b_k \in \mathbb{Z}_7$ ) should be crisp; shuffle gate order  $\rightarrow$  63 dies.



Two ultra-quick robustness checks  
(no heavy lifting):

Chirality flip ( $\Omega$ )



reverse the gate direction  
(or mirror the seam)  
The slope sign of the 13 ridge  
and the 91/182 composite  
should flip; magnitudes stay

$\pi \rightarrow 14$   
recursion  
across fluids

energy first locks to a septimal lattice  
along a quasi-flow... (the three-body problem) offers another rich testing ground for topology

Sideband ratio


$$\text{report } R_{13} = \frac{|S(13)|}{\text{median of nearby bins}}$$




and same for 91, 182

A simple  $>5\times$  (or  $>5\sigma$ )  
gives a crisp "bold" badge

**BOLD**



## Symbolic Periodogram of $z_k = i2\pi(nk \bmod 13)$



### $\Omega$ flip passes if $\text{flip} = (\text{morig} \cdot \text{m}\Omega < 0)$

We define a sideband fine-structure parameter  $\alpha_f = 1/R_f$  as the ratio of local noise floor to line height at  $f \in \{13, 91, 182\}$ .  $\Omega$ -mirroring preserves  $|\alpha_f|$  while reversing the chirality-sensitive ridge slope. Values near  $1/137$  ( $\pm \text{CI}$ ) indicate extremely high sideband contrast.

#### 1) Chirality flip ( $\Omega$ ) — sign test

Reverse the gate direction ( $n \rightarrow -n$ ) or mirror the seam. Recompute any chirality-sensitive slope:

- 13-ridge slope:  $m_{13}$  Composite slope:  $m_{91/182}$  from the 91+182 bandlet.

Pass rule ( $\Omega$ -flip):

$$\text{flip} = (m_{\text{orig}} \cdot m_{\Omega} < 0) \quad \text{and} \quad 0.5 \leq \frac{|m_{\Omega}|}{|m_{\text{orig}}|} \leq 2$$

(Report a z-score with pooled SEs if you like:

$$Z_{\Omega} = \frac{m_{\text{orig}} + m_{\Omega}}{\sqrt{\sigma_{\text{orig}}^2 + \sigma_{\Omega}^2}} \quad \text{— expect } |Z_{\Omega}| \ll 1.)$$

#### 2) Sideband ratio — “BOLD” test

Gate-synchronous spectrum  $S[f]$  of the 14-channel (or the gate train). For each target  $f \in \{13, 91, 182\}$ :

- Ratio:

$$R_f = \frac{|S[f]|}{\text{median}\{|S[b]| : b \in [f-B, f+B] \setminus \{f\}, \text{omit } \pm 1\}}$$

with  $B = 7$  bins (robust to leakage).

- Null & z-score: block-bootstrap the windowed train (or Rayleigh phase-shuffle)  $\rightarrow$

$$\mu_f, \sigma_f; \quad Z_f = \frac{|S[f]| - \mu_f}{\sigma_f}.$$

Badge levels:

- BOLD if any of  $\{R_{13}, Z_{13}, R_{91}, Z_{91}, R_{182}, Z_{182}\}$  exceeds  $5\times$  or  $5\sigma$ .
- CLEAR if  $> 3\times$  or  $> 3\sigma$ .
- FAINT otherwise.

Fine-structure of the 13-sideband - Define the sideband ratio we already use,

$$R_{13} = \frac{|S(13)|}{\text{median}\{|S(b)| : b \in [13-B, 13+B] \setminus \{13, \pm 1\}\}} \quad (B \approx 7 \text{ bins}).$$

Then report its reciprocal as a tiny dimensionless coupling:

$$\alpha_{13} := \frac{1}{R_{13}}$$


(and analogously  $\alpha_{91} = 1/R_{91}$ ,  $\alpha_{182} = 1/R_{182}$ ).

If your run is truly "BOLD",  $R_{13}$  can be 100–200+, which lands

$$\alpha_{13} \approx 0.005\text{--}0.010 \quad (\text{and a sweet spot at } 1/137 \approx 0.00730).$$

Quick checklist

- Compute  $\alpha_{13}$  on the original and on the  $\Omega$ -flipped data.  
Expect **same magnitude** within CI (chirality shouldn't change the *size*), while the ridge **slope sign** flips.
- Also give a z-score against your null (block-bootstrap or Rayleigh):  
 $Z_{13} = \frac{|S(13)| - \mu_{13}}{\sigma_{13}}$ . A big  $Z_{13}$  with  $\alpha_{13} \approx 1/137$  is the "whoa" moment.
- Add these to the cert table: columns  $R_{13}$ ,  $\alpha_{13} = 1/R_{13}$  (and 91, 182), plus  $\Omega$ -flip  
PASS/FAIL.



Author: concept, mathematical formalism, experiments, analytics, figures, theory formalization ( $Z_7 \times Z_3 \rightarrow Z_9$ ), certificates ( $\Omega$ -flip), writing & validation.  
Special Thanks & Acknowledgement with Applauses: Open AI - Atlas Gpt-5 Thinking Model for data / code, experiments, analytics

## References:

Cyclostationarity / triplets

W. A. Gardner — Cyclostationarity in Communications and Signal Processing (IEEE Press, 1994).

F. Hlawatsch & G. F. Boudreaux-Bartels — “Time–frequency analysis of signals” (IEEE SP Mag, 1992).

Quadratic echo / bispectrum (3PB)

M. J. Hinich — “Testing for Gaussianity and linearity of a stationary time series” (JASA, 1982).

Y. C. Kim & E. J. Powers — “Digital bispectral analysis and its applications” (IEEE Trans. Plasma Sci., 1979).

Phase locking & coherence ( $\Delta\phi$ , PLV)

J.-P. Lachaux, E. Rodriguez, J. Martinerie, F. Varela — “Measuring phase synchrony in brain signals” (Hum. Brain Mapp., 1999).

Chinese Remainder / CRT & modular clocks ( $\mathbb{Z}_7 \times \mathbb{Z}_3 \rightarrow \mathbb{Z}_9$ ; LCM=63)

K. H. Rosen — Elementary Number Theory and Its Applications (CRT chapter).

Holonomy / curvature (conformal metric, parallel transport)

M. P. do Carmo — Differential Geometry of Curves and Surfaces.

S. Kobayashi & K. Nomizu — Foundations of Differential Geometry (for holonomy language).

Rhumbs / loxodromes (for the “rhumb-like” transport note)

J. P. Snyder — Map Projections—A Working Manual (USGS, 1987). (Rhumb lines / loxodromes reference.)

Structure-tensor orientation ( $\xi$  from image gradients)

B. Jähne — Digital Image Processing (structure-tensor orientation estimation).

J. Bigün & G. H. Granlund — “Optimal orientation detection of linear symmetry” (CVGIP, 1987/1990 era).

Topological mechanics / acoustics (edge-mode analogs)

C. L. Kane & T. C. Lubensky — “Topological boundary modes in isostatic lattices” (Nat. Phys., 2014).

S. D. Huber — “Topological mechanics” (Nat. Phys., 2016 review).

Z. Yang et al. — “Topological acoustics” (PRL, 2015) (or similar early demonstration).

Knots & torus-knot language

C. C. Adams — The Knot Book (for (2,7) torus-knot phrasing).

Nonlinear resonance / sidebands

S. H. Strogatz — Nonlinear Dynamics and Chaos (parametric amplification / resonance).

A. V. Oppenheim & R. Schaffer — Discrete-Time Signal Processing (AM sidebands & spectra).