

# Algebraic Topology

Lectured by Prof Ivan Smith  
Typed by David Kurniadi Angdinata

Michaelmas 2020

**Syllabus**

# Contents

<b>0</b>	<b>Introduction</b>	<b>3</b>
0.1	Connectedness . . . . .	3
0.2	Homotopy . . . . .	4
<b>1</b>	<b>Singular homology and cohomology</b>	<b>5</b>
1.1	Chain and cochain complexes . . . . .	5
1.2	Singular homology and cohomology . . . . .	6
1.3	Basic examples . . . . .	8
1.4	Structural theorems . . . . .	9
1.5	The sphere . . . . .	10
1.6	Degrees . . . . .	12
1.7	The Klein bottle . . . . .	13
<b>2</b>	<b>Structural theorems</b>	<b>15</b>
2.1	Chain homotopy . . . . .	15
2.2	Homotopy invariance . . . . .	15
2.3	The long exact sequence . . . . .	17
2.4	Relative homology . . . . .	18
2.5	Reduced homology and good pairs . . . . .	20
2.6	Mayer-Vietoris and excision . . . . .	21
2.7	Small simplices theorem . . . . .	22
<b>3</b>	<b>Cellular homology and cohomology</b>	<b>24</b>
3.1	Cell complexes . . . . .	24
3.2	Point-set digression . . . . .	24
3.3	Cellular homology . . . . .	26
3.4	Degrees . . . . .	28
3.5	Cellular cohomology . . . . .	29
3.6	Euler characteristic . . . . .	31

## 0 Introduction

### 0.1 Connectedness

Lecture 1  
Friday  
09/10/20

Algebraic topology concerns the connectivity properties of topological spaces.

**Definition.** A space  $X$  is **path-connected** if for  $p, q \in X$ , there exists  $\gamma : [0, 1] \rightarrow X$  continuous with  $\gamma(0) = p$  and  $\gamma(1) = q$ .

**Example.**  $\mathbb{R}$  is path-connected, and  $\mathbb{R} \setminus \{0\}$  is not.

**Corollary 0.1** (Intermediate value theorem). *If  $f : \mathbb{R} \rightarrow \mathbb{R}$  is continuous and  $x < y$  satisfy  $f(x) > 0$  and  $f(y) < 0$  then  $f$  takes the value zero on  $[x, y]$ .*

*Proof.* Otherwise  $f^{-1}(-\infty, 0) \cup f^{-1}(0, \infty)$  disconnect  $[x, y]$ , a contradiction.  $\square$

**Definition.** Let  $X$  and  $Y$  be topological spaces. Maps  $f_0, f_1 : Y \rightarrow X$  are **homotopic** if there exists  $F : Y \times [0, 1] \rightarrow X$  continuous such that  $F|_{Y \times \{0\}} = f_0$  and  $F|_{Y \times \{1\}} = f_1$ . Write  $f_0 \simeq f_1$ , or  $f_0 \simeq_F f_1$ .

**Exercise.**  $\simeq$  is an equivalence relation on the set of maps from  $Y$  to  $X$ .

Note that  $X$  is **path-connected** if and only if every two maps  $\{\text{point}\} \rightarrow X$  are homotopic. Let

$$S^n = \{x \in \mathbb{R}^{n+1} \mid \|x\| = 1\},$$

so  $S^1 = \{z \in \mathbb{C} \mid |z| = 1\}$ .

**Definition.**  $X$  is **simply-connected** if every two maps  $S^1 \rightarrow X$  are homotopic.

**Example.**  $\mathbb{R}^2$  is simply-connected, and  $\mathbb{R}^2 \setminus \{0\}$  is not. From complex analysis you know  $\gamma : S^1 \rightarrow \mathbb{R}^2 \setminus \{0\}$  has a **winding number** or **degree**  $\deg \gamma \in \mathbb{Z}$ , for which

- if  $\gamma_n(t) = e^{2\pi i n t}$  then  $\deg \gamma_n = n$ , and
- $\deg \gamma_1 = \deg \gamma_2$  if  $\gamma_1 \simeq \gamma_2$ .

For differentiable  $\gamma$ ,  $\deg \gamma = \frac{1}{2\pi i} \int_\gamma \frac{1}{z} dz$ .

**Corollary 0.2** (Fundamental theorem of algebra). *Every non-constant complex polynomial has a root.*

*Proof.* Let  $f(z) = z^n + a_1 z^{n-1} + \dots + a_n$  be non-constant, and without loss of generality monic. Suppose  $f(z) \neq 0$  for all  $z \in \mathbb{C}$ . Let

$$\gamma_R(t) = f(Re^{2\pi i t}),$$

so  $\gamma_R : S^1 \rightarrow \mathbb{R}^2 \setminus \{0\}$ . Since  $\gamma_0$  is constant,  $\deg \gamma_0 = 0$ , so  $\deg \gamma_R = 0$  for all  $R$ . But take  $R \gg \sum_i |a_i|$ . Let

$$f_s(z) = z^n + s(a_1 z^{n-1} + \dots + a_n), \quad 0 \leq s \leq 1.$$

On the circle  $|z| = R$ ,  $f_s(z) \neq 0$  for all  $s$ . So if

$$\gamma_{R,s}(t) = f_s(Re^{2\pi i t}),$$

then  $\gamma_{R,1} = \gamma_R$ , which has degree zero from before, and  $\gamma_{R,0} : t \mapsto R^n e^{2\pi i n t}$ , which has degree  $n \neq 0$ , a contradiction.  $\square$

**Definition.**  $X$  is  **$k$ -connected** if every two maps  $S^i \rightarrow X$  are homotopic whenever  $i \leq k$ .

**Example.**  $\mathbb{R}^n$  is  $(n-1)$ -connected, and  $\mathbb{R}^n \setminus \{0\}$  is not. Maps  $S^{n-1} \rightarrow \mathbb{R}^n \setminus \{0\}$  have a homotopy invariant degree in  $\mathbb{Z}$ , where the degree of the inclusion is one and the degree of the constant map is zero. You may well not have seen this, and we will prove it later.

**Corollary 0.3** (Brouwer's theorem). *Any map  $f : \overline{B^n} = \{x \in \mathbb{R}^n \mid \|x\| \leq 1\} \rightarrow \overline{B^n}$  has a fixed point.*

*Proof.* Suppose  $f$  has no fixed point. Let

$$\gamma_R(v) = Rv - f(Rv), \quad 0 \leq R \leq 1, \quad v \in S^{n-1} = \partial \overline{B^n}.$$

Since  $f$  has no fixed point,  $\gamma_R$  takes values in  $\mathbb{R}^n \setminus \{0\}$ . Since  $\gamma_0$  is constant,  $\deg \gamma_0 = 0$ , so  $\deg \gamma_1 = 0$  by homotopy invariance. Let

$$\gamma_{1,s}(v) = v - sf(v), \quad 0 \leq s \leq 1.$$

Then  $\gamma_{1,1} = \gamma_1$ , and  $\text{im } \gamma_{1,s} \subseteq \mathbb{R}^n \setminus \{0\}$  as  $\|v\| = 1$  and  $\|sf(v)\| = |s|\|f(v)\| < 1$  if  $s < 1$ , so  $\deg \gamma_{1,0} = \deg \gamma_{1,1}$ . The inclusion has  $\deg \gamma_{1,0} = 1$  and  $\deg \gamma_{1,1} = 0$  from above, a contradiction.  $\square$

## 0.2 Homotopy

**Definition.**  $f : X \rightarrow Y$  is a **homotopy equivalence** if there exists  $g : Y \rightarrow X$  such that  $f \circ g \simeq \text{id}_Y$  and  $g \circ f \simeq \text{id}_X$ . Then  $g$  is a **homotopy inverse** for  $f$ , and  $\simeq$  is an equivalence relation on spaces.

**Example.** If  $X$  and  $Y$  are homeomorphic they are trivially homotopy equivalent, by taking  $g = f^{-1}$ .

**Example.**  $\mathbb{R}^n \setminus \{0\} \simeq S^{n-1}$ . Let

$$\begin{aligned} f : \mathbb{R}^n \setminus \{0\} &\longrightarrow S^{n-1} \\ v &\longmapsto \frac{v}{\|v\|}, \end{aligned}$$

and let  $g : S^{n-1} \hookrightarrow \mathbb{R}^n \setminus \{0\}$  be the inclusion. Then  $f \circ g = \text{id}_{S^{n-1}}$  and  $g \circ f \simeq_F \text{id}_{\mathbb{R}^n \setminus \{0\}}$  via the homotopy

$$F(t, v) = tv + (1-t) \frac{v}{\|v\|}.$$

**Example.**  $\{0\} \simeq \mathbb{R}^n$  is a homotopy equivalence.<sup>1</sup> If  $X \simeq \{\text{point}\}$  we say  $X$  is **contractible**.

Algebraic topology is the study of topological spaces up to homotopy equivalence. The idea is that homeomorphism is too delicate a relation. Homotopy equivalence keeps track of essential topological information. More precisely, we assign

$$\{\text{spaces}\} \rightarrow \{\text{groups}\}, \quad \{\text{maps of spaces}\} \rightarrow \{\text{homomorphism of groups}\},$$

so we get algebraic invariants. They are defined for all spaces, but have more structure and use or interest for nicer spaces. The classical first attempt is homotopy theory. One can concatenate loops  $\gamma$  and  $\tau$  by

$$(\gamma * \tau)(t) = \begin{cases} \gamma(2t) & t \leq \frac{1}{2} \\ \tau(1-2t) & t \geq \frac{1}{2} \end{cases}.$$

This is a well-defined operation on the **fundamental group**

$$\pi_1(X, x_0) = \{\text{maps } \gamma : S^1 \rightarrow X \mid \gamma(0) = x_0 \text{ fixed}\} / (\simeq \text{ preserving } x_0).$$

Similarly, the  **$n$ -th homotopy group** is

$$\pi_n(X, x_0) = \{\text{based maps } S^n \rightarrow X \text{ at } x_0\} / \simeq.$$

The issue is that they are very hard to compute, such as  $\pi_n(S^2, x_0)$  not known for all  $n$ . There is no simply-connected **manifold**, a Hausdorff second countable space  $X$  locally homeomorphic to  $\mathbb{R}^n$ , of dimension greater than zero, with  $\pi_n(X)$  known for all  $n$ . So we will do something else, homology and cohomology. It is algebraically harder to set up, but the computational gain is worth it. Note that computing cohomology of harder spaces, such as the space of diffeomorphisms of some manifold or the space of embeddings of one manifold into another, is still very hard.

**Remark.**

- Algebraic topology is all about being able to compute. It is important to do lots of examples.
- Our nice spaces are manifolds and indeed smooth manifolds. There is some overlap with differential geometry which will be useful, not essential but advised.

---

<sup>1</sup>Exercise: check

# 1 Singular homology and cohomology

Lecture 2  
Monday  
12/10/20

We will define invariants of spaces in two stages.

- Associate to  $X$  a chain or cochain complex.
- Take the homology or cohomology of that complex.

## 1.1 Chain and cochain complexes

**Definition.** A **chain complex**  $(C_\bullet, \partial)$  is a sequence of abelian groups and homomorphisms

$$\cdots \rightarrow C_{n+1} \xrightarrow{\partial_{n+1}} C_n \xrightarrow{\partial_n} C_{n-1} \rightarrow \cdots,$$

such that  $\partial_n \circ \partial_{n+1} = 0$  for all  $n$ . We write  $\partial^2 = 0$ , and  $\partial$  is the **differential** or **boundary map**. The **homology groups**  $H_\bullet(C_\bullet, \partial)$  are the graded groups

$$H_n(C_\bullet) = \ker \partial_n / \operatorname{im} \partial_{n+1}.$$

**Definition.** A **cochain complex**  $(C^\bullet, \partial)$  is a sequence of abelian groups and homomorphisms

$$\cdots \rightarrow C^{n-1} \xrightarrow{\partial^{n-1}} C^n \xrightarrow{\partial^n} C^{n+1} \rightarrow \cdots,$$

such that  $\partial^n \circ \partial^{n-1} = 0$  for all  $n$ . We write  $\partial^2 = 0$ , and  $\partial$  is still the **differential** or **boundary map**. The **cohomology groups**  $H^\bullet(C^\bullet, \partial)$  are

$$H^n(C^\bullet) = \ker \partial^n / \operatorname{im} \partial^{n-1}.$$

Elements of  $\ker(\partial : C_n \rightarrow C_{n-1})$  are **cycles**. Elements of  $\operatorname{im}(\partial : C_{n+1} \rightarrow C_n)$  are **boundaries**. Elements of  $\ker(\partial : C^n \rightarrow C^{n+1})$  are **cocycles**. Elements of  $\operatorname{im}(\partial : C^{n-1} \rightarrow C^n)$  are **coboundaries**. Write all  $\partial_i$  and  $\partial^i$  as  $\partial$ , or occasionally  $\partial_\bullet$  and  $\partial^\bullet$ . Elements of  $H_\bullet(C_\bullet)$  are **homology classes** and of  $H^\bullet(C^\bullet)$  are **cohomology classes**.

**Definition.** A **chain map** between chain complexes  $(C_\bullet, \partial)$  and  $(D_\bullet, \partial)$  is a sequence of homomorphisms  $f_n : C_n \rightarrow D_n$  such that for all  $n$  the diagram

$$\begin{array}{ccccccc} \cdots & \longrightarrow & C_n & \xrightarrow{\partial} & C_{n-1} & \longrightarrow & \cdots \\ & & f_n \downarrow & & \downarrow f_{n-1} & & \\ \cdots & \longrightarrow & D_n & \xrightarrow{\partial} & D_{n-1} & \longrightarrow & \cdots \end{array}$$

commutes. That is,  $f_{n-1} \circ \partial_n^{C_\bullet} = \partial_{n-1}^{D_\bullet} \circ f_n$ .

**Exercise.** Define a **cochain map** of cochain complexes.

**Lemma 1.1.** A chain map  $f : C_\bullet \rightarrow D_\bullet$  induces homomorphisms  $(f_*)_n : H_n(C_\bullet) \rightarrow H_n(D_\bullet)$  for each  $n$ .

*Proof.* Let  $[a] \in H_n(C_\bullet)$ , so  $a$  is represented by a cycle  $\alpha \in C_n$ , where  $\partial(\alpha) = 0$ . Then  $\partial(f_n(\alpha)) = f_{n-1}(\partial(\alpha)) = 0$ , so  $f_n(\alpha)$  is a cycle. Define  $(f_*)_n([a]) = [f_n(\alpha)] \in H_n(D_\bullet)$ . We made a choice of representing the cycle  $\alpha$ . But if  $[a]$  is represented by  $\alpha$  and  $\alpha'$ , then  $\alpha - \alpha' \in \operatorname{im}(\partial_{n+1} : C_{n+1} \rightarrow C_n)$ . Say  $\alpha - \alpha' = \partial(\tau)$ . Then  $f_n(\alpha) - f_n(\alpha') = f_n(\alpha - \alpha') = f_n(\partial(\tau)) = \partial(f_{n+1}(\tau))$ , so  $[f_n(\alpha)] = [f_n(\alpha') + \partial(f_{n+1}(\tau))] = [f_n(\alpha')]$  as  $[\operatorname{im} \partial] = 0$  in  $H_n(D_\bullet)$ . So  $(f_*)_n$  is well-defined, and it is easy to see it is a homomorphism.  $\square$

**Exercise.** If  $C_\bullet, D_\bullet, E_\bullet$  are chain complexes and  $f : C_\bullet \rightarrow D_\bullet$  and  $g : D_\bullet \rightarrow E_\bullet$  are chain maps then  $\{g_n \circ f_n : C_n \rightarrow E_n\}_n$  defines a chain map. Also

$$(g \circ f)_* = g_* \circ f_*, \quad (\operatorname{id}_{C_\bullet})_* = \operatorname{id}_{H_\bullet(C_\bullet)} \quad (1)$$

The goal is to associate to a space  $X$  chain complexes  $C_\bullet(X)$  and cochain complexes  $C^\bullet(X)$  such that a map  $f : X \rightarrow Y$  yields chain maps  $f : C_\bullet(X) \rightarrow C_\bullet(Y)$  and cochain maps  $f : C^\bullet(Y) \rightarrow C^\bullet(X)$ . Then (1) will say we have a functor

$$\begin{array}{ccc} \mathbf{Top} & \longrightarrow & \mathbf{Ab} \\ X & \longmapsto & H_\bullet(X) \end{array} ,$$

from the category of topological spaces and continuous maps to the category of abelian groups and homomorphisms. Our complexes  $C_\bullet$  and  $C^\bullet$  will have the benefit that they are intrinsic but will be huge and unwieldy. We will

- prove structure theorems to help compute, and
- find smaller complexes later for nice spaces, such as CW-complexes.

## 1.2 Singular homology and cohomology

**Definition.** The **standard simplex** is

$$\Delta^n = \left\{ (t_0, \dots, t_n) \in \mathbb{R}^{n+1} \mid \forall i, t_i \geq 0, \sum_i t_i = 1 \right\}.$$

The  $i$ -th **face** of  $\Delta^n$  is

$$\Delta_i^n = \{ \underline{t} \in \Delta^n \mid t_i = 0 \}.$$

Note that there exists a canonical homeomorphism

$$\begin{array}{ccc} \delta_i : & \Delta^{n-1} & \longrightarrow \Delta_i^n \subseteq \Delta^n \\ & (t_0, \dots, t_{n-1}) & \longmapsto (t_0, \dots, t_{i-1}, 0, t_i, \dots, t_{n-1}) \end{array} .$$

**Definition.** If  $X$  is a space, a **singular  $n$ -simplex** in  $X$  is a map  $\sigma : \Delta^n \rightarrow X$ . The **singular chain complex**  $(C_\bullet(X), \partial)$  has

$$C_n(X) = \left\{ \sum_{i=1}^N n_i \sigma_i \mid N < \infty, n_i \in \mathbb{Z}, \sigma_i : \Delta^n \rightarrow X \right\},$$

the free abelian group on the singular  $n$ -simplices in  $X$ , and

$$\begin{array}{ccc} \partial : C_n(X) & \longrightarrow & C_{n-1}(X) \\ \sigma & \longmapsto & \sum_{i=0}^n (-1)^i (\sigma \circ \delta_i) \end{array} ,$$

extended linearly.

**Example.**  $\Delta^0$  is a point,  $\Delta^1$  is a line,  $\Delta^2$  is a triangle, and  $\Delta^3$  is a tetrahedron.

Note that  $n+1$  ordered points  $\{v_i\}_{0 \leq i \leq n} \subseteq \mathbb{R}^{n+1}$  determine an  $n$ -simplex if  $\{v_i - v_0 \mid 1 \leq i \leq n\}$  are linearly independent, by taking their convex hull, and

$$\begin{array}{ccc} \sigma : \Delta^n & \longrightarrow & \mathbb{R}^{n+1} \\ \underline{t} & \longmapsto & \sum_{i=0}^n t_i v_i \end{array} .$$

We orient the edges  $v_i \rightarrow v_j$  if  $i < j$ . Write  $[v_0, \dots, v_n]$  for this  $n$ -simplex, then

$$\partial(\sigma) = \sum_{i=0}^n (-1)^i \sigma|_{[v_0, \dots, \widehat{v_i}, \dots, v_n]},$$

where the index  $\widehat{v_i}$  is omitted.

**Lemma 1.2.**  $\partial^2 = 0$ .

*Proof.*

$$\partial(\partial(\sigma)) = \sum_{j < i} (-1)^i (-1)^j \sigma|_{[v_0, \dots, \widehat{v_j}, \dots, \widehat{v_i}, \dots, v_n]} + \sum_{j > i} (-1)^i (-1)^{j-1} \sigma|_{[v_0, \dots, \widehat{v_i}, \dots, \widehat{v_j}, \dots, v_n]}.$$

Exchange  $i$  and  $j$  and the two terms cancel.  $\square$

**Definition.** The **singular homology** of  $X$  is

$$H_\bullet(X) = H_\bullet(X; \mathbb{Z}) = H_\bullet(C_\bullet(X), \partial).$$

Trivially this is a homeomorphism invariant of  $X$ , since we only used the notion of continuous maps to  $X$  to define it.

**Definition.** The **singular cochain complex**  $(C^\bullet(X), \partial^*)$  has

$$C^n(X) = \text{Hom}(C_n(X), \mathbb{Z}),$$

and

$$\begin{aligned} \partial^* : C^n(X) &\longrightarrow C^{n+1}(X) \\ \psi &\longmapsto (\sigma \mapsto \psi(\partial(\sigma))) \end{aligned}, \quad \sigma \in C_{n+1}(X),$$

which is adjoint to  $\partial$ .

Then  $\partial^*(\partial^*(\psi))(\sigma) = \partial^*(\psi)(\partial(\sigma)) = \psi(\partial(\partial(\sigma))) = 0$ , so  $(\partial^*)^2 = 0$  and this is a cochain complex.

**Definition.** The **singular cohomology** of  $X$  is

$$H^\bullet(X; \mathbb{Z}) = H^\bullet(C^\bullet(X), \partial^*).$$

The following is the rough idea.

- $\partial^2 = 0$  implies that the boundary of the boundary vanishes.
- $H_i(X)$  will probe  $i$ -dimensional holes or regions in  $X$ .
- $H^i(X)$  will be a rule associating an integer to an  $i$ -dimensional region of  $X$ .

Note that  $H^\bullet(X; \mathbb{Z}) \not\cong \text{Hom}(H_\bullet(X), \mathbb{Z})$  in general.

**Remark.** Let  $f : X \rightarrow Y$  be continuous. If  $\sigma : \Delta^n \rightarrow X$  then  $f \circ \sigma : \Delta^n \rightarrow Y$ , so  $f$  gives a homomorphism  $(f_\#)_n : C_n(X) \rightarrow C_n(Y)$ . Also  $f \circ (\sigma|_{\Delta_i^n}) \equiv (f \circ \sigma)|_{\Delta_i^n}$ , since  $f \circ (\sigma \circ \delta_i) = (f \circ \sigma) \circ \delta_i$ . Thus

$$\begin{aligned} f_\# : C_\bullet(X) &\longrightarrow C_\bullet(Y) \\ \sigma &\longmapsto f \circ \sigma \end{aligned}$$

is a chain map such that

$$\begin{array}{ccc} C_n(X) & \xrightarrow{\partial} & C_{n-1}(X) \\ (f_\#)_n \downarrow & & \downarrow (f_\#)_{n-1} \\ C_n(Y) & \xrightarrow{\partial} & C_{n-1}(Y) \end{array}$$

which gives homomorphisms

$$f_* : H_\bullet(X) \rightarrow H_\bullet(Y),$$

that is  $(f_*)_n : H_n(X) \rightarrow H_n(Y)$  for each  $n$ . By the exercise,

$$((f \circ g)_*)_n = (f_*)_n \circ (g_*)_n, \quad ((\text{id}_{C_\bullet(X)})_*)_n = \text{id}_{H_n(X)}.$$

Note that  $f : X \rightarrow Y$  induces a cochain map

$$\begin{aligned} f^\# : C^\bullet(Y) &\longrightarrow C^\bullet(X) \\ \psi &\longmapsto (\sigma \mapsto \psi(f \circ \sigma)) \end{aligned},$$

and homomorphisms

$$f^* : H^\bullet(Y) \rightarrow H^\bullet(X),$$

so cohomology is contravariant.

Lecture 3  
Wednesday  
14/10/20

### 1.3 Basic examples

What can we compute?

**Lemma 1.3.** *Let  $X$  be a point. Then*

$$H_i(\{\text{point}\}) = \begin{cases} \mathbb{Z} & i = 0 \\ 0 & \text{otherwise} \end{cases}.$$

*Proof.* For each  $n \geq 0$ , there exists a unique  $n$ -simplex  $\sigma_n : \Delta^n \rightarrow \{\text{point}\}$  in  $X$ , the constant map. Then  $\partial(\sigma_1) = \sigma_1 \circ \delta_0 - \sigma_1 \circ \delta_1 = \sigma_0 - \sigma_0 = 0$  and  $\partial(\sigma_2) = \sigma_2 \circ \delta_0 - \sigma_2 \circ \delta_1 + \sigma_2 \circ \delta_2 = \sigma_1 - \sigma_1 + \sigma_1 = \sigma_1$ , and

$$\partial(\sigma_n) = \begin{cases} \sigma_{n-1} & n \text{ even} \\ 0 & n \text{ odd} \end{cases}.$$

So  $C_\bullet(\{\text{point}\})$  is

$$\begin{array}{ccccccc} \dots & \longrightarrow & C_3(\{\text{point}\}) & \longrightarrow & C_2(\{\text{point}\}) & \longrightarrow & C_1(\{\text{point}\}) & \longrightarrow & C_0(\{\text{point}\}) \\ & & \parallel & & \parallel & & \parallel & & \parallel \\ \dots & \xrightarrow{1} & \mathbb{Z} & \xrightarrow{0} & \mathbb{Z} & \xrightarrow{1} & \mathbb{Z} & \xrightarrow{0} & \mathbb{Z} \end{array}.$$

Now check the result. □

**Exercise.**

$$H^i(\{\text{point}\}) \cong \begin{cases} \mathbb{Z} & i = 0 \\ 0 & \text{otherwise} \end{cases}.$$

There is basically only one other computation we can do from the definitions.

**Lemma 1.4.** *If  $X = \bigsqcup_{\alpha \in I} X_\alpha$  is a disjoint union of path-components,*

$$H_i(X) \cong \bigoplus_{\alpha \in I} H_i(X_\alpha).$$

*Proof.* Any continuous map  $\sigma : \Delta^i \rightarrow X$  has image in one  $X_\alpha$  and then all the faces of  $\sigma$  lie in the same  $X_\alpha$ , so

$$C_\bullet(X) = \bigoplus_{\alpha} C_\bullet(X_\alpha),$$

compatibly with the differential. □

**Lemma 1.5.** *If  $X$  is path-connected and non-empty,*

$$H_0(X) \cong \mathbb{Z}.$$

We sometimes write  $\pi_0(X)$  for the set of path-components of  $X$ .

*Proof.* Define the **augmentation**

$$\begin{aligned} \epsilon : C_0(X) &\longrightarrow \mathbb{Z} \\ \sum_i n_i \sigma_i &\longmapsto \sum_i n_i, \end{aligned}$$

where  $\sigma_i : \{\text{point}\} \rightarrow X$  are 0-simplices in  $X$ . Since  $X \neq \emptyset$ ,  $\epsilon$  is onto. If  $\tau = [v_0, v_1] : \Delta^1 \rightarrow X$ , then  $\epsilon(\partial(\tau)) = \epsilon(v_1 - v_0) = 0$ . So  $\text{im}(\partial : C_1(X) \rightarrow C_0(X)) \subseteq \ker \epsilon$ , so  $\epsilon$  defines  $H_0(X) = C_0(X) / \text{im } \partial \rightarrow \mathbb{Z}$ . So far we did not use path-connectivity. But suppose  $\sum_i n_i \sigma_i \in \ker \epsilon$ . Fix a basepoint  $p \in X$ . For all  $i$  pick

$$\begin{aligned} \tau_i : \Delta^i \cong [0, 1] &\longrightarrow X \\ 1 &\longmapsto \sigma_i \\ 0 &\longmapsto p \end{aligned}$$

Then  $\partial(\sum_i n_i \tau_i) = \sum_i n_i \sigma_i - (\sum_i n_i) p = \sum_i n_i \sigma_i$ , as  $\sum_i n_i \sigma_i \in \ker \epsilon$ , so  $\ker \epsilon \subseteq \text{im } \partial$  and  $\epsilon : H_0(X) \xrightarrow{\sim} \mathbb{Z}$ . □



## 1.4 Structural theorems

The following is an informal picture. Let  $X$  be an annulus, and let  $\sigma : \Delta^1 \rightarrow X$  be a 1-simplex, which happens to be a closed loop  $[0, 1] \rightarrow X$  going around the inner circle. Recall that  $\sigma$  has  $\partial(\sigma) = \sigma(1) - \sigma(0) = 0$ , so  $\sigma$  defines  $[\sigma] \in H_1(X)$ . We would hope this is non-zero, as we cannot see a way to fill in  $\sigma$  with 2-simplices, in contrast to a 1-simplex  $\tau : \Delta^1 \cong [0, 1] \rightarrow X$  away from the inner circle. But  $C_i(X)$  is uncountably generated for all  $i$  and very hard to control. A question is how do we rule out all configurations of 2-simplices, or other representatives for  $[\sigma] \in H_i(X)$ ? Informally, in the realm of nice spaces, there is nothing else you can compute from the definition. Homology and cohomology are rendered useful by a collection of structural theorems. We will state these, and see how to use them, and then return to prove them later.

**Theorem 1.6** (Homotopy invariance). *If  $f : X \rightarrow Y$  and  $g : X \rightarrow Y$  are homotopic, then*

$$f_* = g_* : H_\bullet(Y) \rightarrow H_\bullet(Y), \quad f^* = g^* : H^\bullet(Y) \rightarrow H^\bullet(Y).$$

**Corollary 1.7.** *If  $X \simeq Y$  then  $H_\bullet(X) \cong H_\bullet(Y)$  and  $H^\bullet(X) \cong H^\bullet(Y)$ .*

*Proof.* There exist  $f : X \rightarrow Y$  and  $g : Y \rightarrow X$  such that  $g \circ f \simeq \text{id}_X$  and  $f \circ g \simeq \text{id}_Y$ , so  $(f_*)^{-1} = g_*$  are isomorphisms.  $\square$

Thus homology and cohomology are insensitive to inessential deformations of a space.

**Corollary 1.8.** *For every  $n$ ,*

$$H_\bullet(\mathbb{R}^n) = \begin{cases} \mathbb{Z} & \bullet = 0 \\ 0 & \text{otherwise} \end{cases},$$

*and similarly for  $H^\bullet(\mathbb{R}^n)$ .*

**Definition.** An **exact sequence** is a chain or cochain complex with vanishing homology or cohomology, so

$$\cdots \rightarrow C_{n+1} \xrightarrow{\partial_{n+1}} C_n \xrightarrow{\partial_n} C_{n-1} \rightarrow \cdots,$$

such that  $\ker \partial_n = \text{im } \partial_{n+1}$  for all  $n$ .

- Given homomorphisms

$$A \xrightarrow{f} B \xrightarrow{g} C,$$

say this is **exact at  $B$**  if  $\ker g = \text{im } f$ .

- If

$$0 \rightarrow A \xrightarrow{f} B \rightarrow 0$$

is exact,  $A \cong_f B$ .

- A **short exact sequence** is one of shape

$$0 \rightarrow A \xrightarrow{f} B \xrightarrow{g} C \rightarrow 0.$$

**Example.** If

$$0 \rightarrow \mathbb{Z} \rightarrow A \rightarrow \mathbb{Z}/n \rightarrow 0,$$

possibly  $A = \mathbb{Z} \oplus \mathbb{Z}/n$ , and

$$0 \rightarrow \mathbb{Z} \xrightarrow{1 \mapsto (1,0)} \mathbb{Z} \oplus \mathbb{Z}/n \xrightarrow{(0,1) \mapsto 1} \mathbb{Z}/n \rightarrow 0$$

or  $A = \mathbb{Z}$ , and

$$0 \rightarrow \mathbb{Z} \xrightarrow{\cdot n} \mathbb{Z} \xrightarrow{p \mapsto p \bmod n} \mathbb{Z}/n \rightarrow 0.$$

See question sheet 1.

**Theorem 1.9** (Mayer-Vietoris). *If  $X = A \cup B$  with  $A$  and  $B$  open, there are **Mayer-Vietoris boundary homomorphisms**  $\partial_{MV} : H_{i+1}(X) \rightarrow H_i(A \cap B)$ , yielding a **long exact sequence***

$$\cdots \rightarrow H_{i+1}(X) \xrightarrow{\partial_{MV}} H_i(A \cap B) \xrightarrow{((i_A)_*, (i_B)_*)} H_i(A) \oplus H_i(B) \xrightarrow{(j_A)_* - (j_B)_*} H_i(X) \rightarrow \cdots,$$

where

$$\begin{array}{ccc} A \cap B & \xhookrightarrow{i_A} & A \\ i_B \downarrow & & \downarrow j_A \\ B & \xhookrightarrow{j_B} & X \end{array}.$$

The Mayer-Vietoris boundary homomorphism is defined algebraically and is not associated to a map of spaces.

**Remark.** Suppose  $\sigma \in C_{i+1}(X)$  is a cycle, so  $\partial(\sigma) = 0$ , and  $\sigma = \alpha + \beta$  for chains  $\alpha \in C_{i+1}(A)$  and  $\beta \in C_{i+1}(B)$ . Then  $\partial(\alpha) = -\partial(\beta)$  and  $\partial_{MV}([\sigma]) = [\partial(\alpha)]$ , since  $\partial(\alpha) \in A \cap B$ .

**Remark.** The Mayer-Vietoris sequence is natural, so if  $X = A \cup B$  and  $Y = C \cup D$  and  $f : X \rightarrow Y$  has  $f(A) \subseteq C$  and  $f(B) \subseteq D$  then there are homomorphisms of exact sequences

$$\begin{array}{ccccccc} \cdots & \longrightarrow & H_{i+1}(X) & \xrightarrow{\partial_{MV}} & H_i(A \cap B) & \longrightarrow & H_i(A) \oplus H_i(B) \longrightarrow H_i(X) \longrightarrow \cdots \\ & & \downarrow f_* & & \downarrow f_* & & \downarrow f_* \\ \cdots & \longrightarrow & H_{i+1}(Y) & \xrightarrow{\partial_{MV}} & H_i(C \cap D) & \longrightarrow & H_i(C) \oplus H_i(D) \longrightarrow H_i(Y) \longrightarrow \cdots \end{array},$$

such that all squares commute.

**Remark.** There is a Mayer-Vietoris sequence in cohomology, which is also natural. There are  $\partial_{MV}^* : H^i(A \cap B) \rightarrow H^{i+1}(X)$  such that

$$\cdots \rightarrow H^i(X) \xrightarrow{(j_A^*, j_B^*)} H^i(A) \oplus H^i(B) \xrightarrow{i_A^* - i_B^*} H^i(A \cap B) \xrightarrow{\partial_{MV}^*} H^{i+1}(X) \rightarrow \cdots$$

is exact, where

$$\begin{array}{ccc} A \cap B & \xhookrightarrow{i_A} & A \\ i_B \downarrow & & \downarrow j_A \\ B & \xhookrightarrow{j_B} & X \end{array}.$$

## 1.5 The sphere

**Proposition 1.10.**

$$H_i(S^1) \cong \begin{cases} \mathbb{Z} & i = 0, 1 \\ 0 & \text{otherwise} \end{cases}, \quad H^i(S^1) \cong \begin{cases} \mathbb{Z} & i = 0, 1 \\ 0 & \text{otherwise} \end{cases}.$$

*Proof.* Let  $S^1 = X = A \cup B$  where  $A$  and  $B$  are open intervals such that  $A \cap B$  are two disjoint open intervals, so  $A \simeq \{\text{point}\} \simeq B$  and  $A \cap B \simeq \{\text{point } p\} \sqcup \{\text{point } q\}$ . By homotopy invariance,

$$H_\bullet(\mathbb{R}) = \begin{cases} \mathbb{Z} & \bullet = 0 \\ 0 & \text{otherwise} \end{cases},$$

so we know  $H_\bullet(A)$ ,  $H_\bullet(B)$ , and  $H_\bullet(A \cap B)$ . Mayer-Vietoris for  $i \geq 2$  gives

$$\begin{array}{ccccc} H_i(A) \oplus H_i(B) & \longrightarrow & H_i(S^1) & \longrightarrow & H_{i-1}(A \cap B) \\ \parallel & & & & \parallel \\ 0 & & & & 0 \end{array}.$$

Check that  $H_i(S^1) = 0$ .<sup>2</sup> Mayer-Vietoris for  $i = 0, 1$  gives

$$\begin{array}{ccccccc} H_1(A) \oplus H_1(B) & \longrightarrow & H_1(S^1) & \longrightarrow & H_0(A \cap B) & \longrightarrow & H_0(A) \oplus H_0(B) \longrightarrow H_0(S^1) \\ \downarrow \cong & & & & \downarrow \cong & & \downarrow \cong \\ 0 & & & & \mathbb{Z} \oplus \mathbb{Z} & \xrightarrow{\alpha} & \mathbb{Z} \oplus \mathbb{Z} \xrightarrow{\beta} \mathbb{Z} \end{array}$$

Recall that  $H_0(Z)$  is free abelian on  $\pi_0(Z)$ , the set of path-components, and indeed is generated by  $\sigma : \{\text{point}\} \rightarrow Z$ , for any choice of point in each component. So

$$\alpha = ((i_A)_*, (i_B)_*) : \mathbb{Z}\langle p \rangle \oplus \mathbb{Z}\langle q \rangle \longrightarrow \mathbb{Z} \oplus \mathbb{Z} \\ (a, b) \longmapsto (a + b, a + b),$$

and

$$\beta = (j_A)_* - (j_B)_* : \mathbb{Z} \oplus \mathbb{Z} \longrightarrow \mathbb{Z} \\ (u, v) \longmapsto u - v.$$

By exactness,  $H_1(S^1) \cong \ker \alpha \cong \mathbb{Z}$ , generated by  $(1, -1) \equiv (p, -q) \in H_0(A) \oplus H_0(B)$ . □

The same method as for computing  $H_\bullet(S^1)$  shows the following.

**Proposition 1.11.**

$$H_j(S^n) \cong \begin{cases} \mathbb{Z} & j = 0, n \\ 0 & \text{otherwise} \end{cases}, \quad H^j(S^n) \cong \begin{cases} \mathbb{Z} & j = 0, n \\ 0 & \text{otherwise} \end{cases}.$$

This time let us do the cohomology computation.

*Proof.* Let  $S^n = A \cup B$  where  $A \cong B \cong \mathbb{R}^n$  and  $A \cap B \cong S^{n-1} \times (0, 1) \simeq S^{n-1}$ . By homotopy invariance and induction, we know  $H^\bullet(A)$ ,  $H^\bullet(B)$ , and  $H^\bullet(A \cap B)$ . Mayer-Vietoris now gives

$$\begin{array}{ccccccc} H^i(\mathbb{R}^n) \oplus H^i(\mathbb{R}^n) & \longrightarrow & H^i(S^{n-1}) & \longrightarrow & H^{i+1}(S^n) & \longrightarrow & H^{i+1}(\mathbb{R}^n) \oplus H^{i+1}(\mathbb{R}^n) \\ \downarrow \cong & & & & & & \downarrow \cong \\ 0 & & & & & & 0 \end{array},$$

so  $H^i(S^{n-1}) \xrightarrow{\sim} H^{i+1}(S^n)$  for all  $i > 0$ . For  $i = 0, 1$ ,

$$\begin{array}{ccccccc} H^0(S^n) & \longrightarrow & H^0(\mathbb{R}^n) \oplus H^0(\mathbb{R}^n) & \longrightarrow & H^0(S^{n-1}) & \longrightarrow & H^1(S^n) \longrightarrow H^1(\mathbb{R}^n) \oplus H^1(\mathbb{R}^n) \\ & & & & & & \downarrow \cong \\ & & & & & & 0 \end{array}.$$

We showed before that for path-connected  $X$ ,  $H_0(X) \cong \mathbb{Z}$  is generated by  $\sigma : \{\text{point}\} \rightarrow X \in C_0(X)$ . By question sheet 1,  $H^0(X) \cong \mathbb{Z}$  is generated by

$$\psi : C_0(X) \longrightarrow \mathbb{Z} \\ \sigma \longmapsto 1, \quad \sigma : \{\text{point}\} \rightarrow X.$$

If  $n > 1$ ,  $S^{n-1}$  is connected. So

$$\begin{array}{ccccccc} H^0(S^n) & \longrightarrow & H^0(\mathbb{R}^n) \oplus H^0(\mathbb{R}^n) & \longrightarrow & H^0(S^{n-1}) & \longrightarrow & H^1(S^n) \longrightarrow H^1(\mathbb{R}^n) \oplus H^1(\mathbb{R}^n) \\ \downarrow \cong & & \downarrow \cong & & \downarrow \cong & & \downarrow \cong \\ \mathbb{Z} & \xrightarrow{\alpha} & \mathbb{Z} \oplus \mathbb{Z} & \xrightarrow{\alpha} & \mathbb{Z} & & 0 \end{array},$$

where  $\alpha(p, q) = p + q$  is onto, so  $H^1(S^n) = 0$ , and now we have computed enough to complete the induction. □

**Corollary 1.12.**  $\mathbb{R}^m \cong \mathbb{R}^n$  if and only if  $m = n$ .

*Proof.* If  $\mathbb{R}^m \cong \mathbb{R}^n$ , then  $S^{m-1} \simeq \mathbb{R}^m \setminus \{0\} \cong \mathbb{R}^n \setminus \{0\} \simeq S^{n-1}$ , so  $S^{m-1} \simeq S^{n-1}$ . Thus  $H_\bullet(S^{m-1}) \cong H_\bullet(S^{n-1})$ , so  $m = n$ . □

This homeomorphism invariance of dimension was an early success of the subject. Recall there are space-filling curves  $\phi : [0, 1] \rightarrow [0, 1]^2$  that are continuous and surjective.

<sup>2</sup>Exercise

## 1.6 Degrees

**Lemma 1.13.** Assume  $n > 0$ . A map  $f : S^n \rightarrow S^n$  has a **degree**  $\deg f \in \mathbb{Z}$  and if  $g \simeq f$ , then  $\deg g = \deg f$ .

*Proof.*  $f$  induces  $(f_*)_n : H_n(S^n) \cong \mathbb{Z} \rightarrow H_n(S^n) \cong \mathbb{Z}$ , which is multiplication by an integer. This defines  $\deg f$ . If  $g \simeq f$ , then  $g_* = f_*$ . A caveat is to use the same isomorphism on both sides and make sure  $\deg f$  is defined and not just up to sign.  $\square$

**Exercise.** Check that  $\deg(f \circ g) = \deg f \cdot \deg g$ .

**Example.**  $\deg \text{id} = 1$ , so if  $f$  is a homeomorphism,  $\deg f \in \{\pm 1\}$ .

**Example.** The degree of the constant map is zero, since the constant map

$$\begin{array}{ccc} f & : & S^n \longrightarrow S^n \\ & & x \longmapsto p \end{array}$$

factorises as  $S^n \rightarrow \{\text{point}\} \rightarrow S^n$ , so

$$\begin{array}{ccccc} H_n(S^n) & \longrightarrow & H_n(\{\text{point}\}) & \longrightarrow & H_n(S^n) \\ \cong & & \cong & & \cong \\ \mathbb{Z} & \xrightarrow{\quad \quad \quad} & 0 & \xrightarrow{\quad \quad \quad} & \mathbb{Z} \end{array}$$

factorises through the zero group.

Note that combining with  $S^{n-1} \simeq \mathbb{R}^n \setminus \{0\}$ , this fills in details, modulo homotopy invariance and Mayer-Vietoris, for results from the first lecture on Brouwer's theorem.

**Lemma 1.14.** Let  $O(k) = \{A \in \text{Mat}_k \mathbb{R} \mid AA^T = I\}$ . A matrix  $A \in O(n+1)$ , which acts on  $S^n \subseteq \mathbb{R}^{n+1}$ , acts on  $H_n(S^n)$  by multiplication by  $\det A$ .

*Proof.*  $O(n+1)$  has two path-connected components, so by homotopy invariance of degree, it suffices to show reflection in a hyperplane has degree  $-1$ . Let  $H = S^{n-1}$  be a hyperplane, let  $L$  be an invariant hemisphere, and let  $H' = \partial L \cap H$ . Note that a reflection  $r_H : S^n \rightarrow S^n$  in  $H$  induces a reflection  $r_{H'} : \partial L = S^{n-1} \rightarrow \partial L = S^{n-1}$  in  $H'$ . We computed  $H_\bullet(S^n)$  by Mayer-Vietoris, using the decomposition which is  $r_H$ -invariant. By the naturality of Mayer-Vietoris,

$$\begin{array}{ccccccc} 0 & \longrightarrow & H_n(S^n) & \xrightarrow{\sim} & H_{n-1}(S^{n-1}) & \longrightarrow & 0 \\ & & \downarrow r_H & & \downarrow r_{H'} & & \\ 0 & \longrightarrow & H_n(S^n) & \xrightarrow{\sim} & H_{n-1}(S^{n-1}) & \longrightarrow & 0 \end{array},$$

so inductively, it suffices to treat the case  $n = 1$ . So consider a circle  $S^1 = A \cup B$  where  $p, q \in A \cap B$ . Our former Mayer-Vietoris computation of  $H_\bullet(S^1)$  gave

$$\begin{array}{ccccccc} 0 & \longrightarrow & H_1(S^1) & \longrightarrow & H_0(p \sqcup q) & \longrightarrow & H_0(A) \oplus H_0(B) \\ & & & & \downarrow \cong & & \downarrow \cong \\ & & & & \mathbb{Z} \oplus \mathbb{Z} & \xrightarrow{\quad \quad \quad} & \mathbb{Z} \oplus \mathbb{Z} \end{array},$$

and  $H_1(S^1) = \ker \alpha \cong \mathbb{Z} \langle (1, -1) \rangle$  is generated by  $p - q$ . So as  $r_H$  exchanges  $p$  and  $q$  it acts on  $H_1(S^1)$  by  $-1$ .  $\square$

**Corollary 1.15.**

1. The antipodal map

$$\begin{array}{ccc} a_n & : & S^n \longrightarrow S^n \\ & & x \longmapsto -x \end{array}$$

has degree  $(-1)^{n+1}$ .

2. If  $f : S^n \rightarrow S^n$  has no fixed point,  $f \simeq a_n$ .

3. If  $G$  acts freely on  $S^{2k}$ , then  $G \leq \mathbb{Z}/2$ .

*Proof.*

1.  $a_n : S^n \rightarrow S^n$  is a composition of  $n + 1$  reflections  $\mathbb{R}^{n+1} \rightarrow \mathbb{R}^{n+1}$ .
2. We will show if  $f(x) \neq g(x)$  for all  $x$ , then  $f \simeq a_n \circ g$ . Consider

$$\phi_t : x \mapsto \frac{tf(x) - (1-t)g(x)}{\|tf(x) - (1-t)g(x)\|}, \quad 0 \leq t \leq 1.$$

Note that  $tf(x) + (1-t)g(x) \neq 0$  or  $t = \frac{1}{2}$  and  $f(x) = g(x)$ , a contradiction. So  $f = \phi_1 \simeq \phi_0 = a_n \circ g$ .

3. Question sheet 1.

□

We borrow a concept from differential topology. A **vector field** on  $S^n$  is a map  $v : S^n \rightarrow \mathbb{R}^{n+1}$  such that for all  $x \in S^n$ , the Euclidean inner product on  $\mathbb{R}^{n+1}$  has  $\langle x, v(x) \rangle = 0$ . Note that this is a global section of the tangent bundle  $TS^n \rightarrow S^n$ .

**Proposition 1.16** (Hairy ball theorem).  $S^n$  has a nowhere-vanishing vector field if and only if  $n$  is odd.

*Proof.* If  $n = 2k - 1$ , set

$$v(x_1, y_1, \dots, x_k, y_k) = (-y_1, x_1, \dots, -y_k, x_k).$$

Suppose  $n$  is even, and for contradiction that such  $v$  exists. So  $v/\|v\| : S^n \rightarrow S^n$ . Consider

$$v_t(x) = (\cos t)x + (\sin t)\frac{v}{\|v\|}(x).$$

Then  $|v_t(x)| = 1$  for all  $t$ , and  $v_0 = \text{id}$  and  $v_\pi = -\text{id} = a_n$ , so  $\text{id}_{S^n} \simeq a_n$ . Thus  $\deg \text{id} = \deg a_n$ , so  $1 = (-1)^{n+1}$ . □

## 1.7 The Klein bottle


We should do one computation which involves knowing the maps, not just on  $H_0(X)$ , in an exact sequence, and not just that the sequence is exact. The **Klein bottle**  $K$  is obtained from gluing two Möbius bands together.

Lecture 5  
Monday  
19/10/20


**Lemma 1.17.**

$$H_j(K; \mathbb{Z}) \cong \begin{cases} \mathbb{Z} & j = 0 \\ \mathbb{Z} \oplus \mathbb{Z}/2 & j = 1 \\ 0 & \text{otherwise} \end{cases}.$$


*Proof.* Apply Mayer-Vietoris to  $K$



where  $A \simeq S^1$  is a Möbius band



and  $B \simeq S^1$  is a similar Möbius band, such that  $A \cap B \simeq S^1$  is



The essential part of the long exact sequence is

$$\begin{array}{ccccccc}
 0 \longrightarrow H_2(K) \longrightarrow H_1(A \cap B) & \xrightarrow{\psi} & H_1(A) \oplus H_1(B) & \longrightarrow & H_1(K) & \xrightarrow{0} & H_0(A \cap B) \longrightarrow H_0(A) \oplus H_0(B) \\
 \parallel & & \parallel & & & & \parallel \\
 \mathbb{Z} & \xrightarrow{\quad \quad \quad} & \mathbb{Z} \oplus \mathbb{Z} & & & & \mathbb{Z} \xrightarrow[p \mapsto (p,p)]{\quad \quad \quad} \mathbb{Z} \oplus \mathbb{Z}
 \end{array}$$

By exactness,  $H_1(K) = (\mathbb{Z} \oplus \mathbb{Z}) / \text{im } \psi$  and  $H_2(K) \cong \ker \psi$ . The key claim is that  $\psi(1) = (2, 2)$  and note  $(\mathbb{Z} \oplus \mathbb{Z}) / \langle 2, 2 \rangle \cong \mathbb{Z} \oplus \mathbb{Z}/2$ . For this,  $A \cap B$  is homotopy equivalent to the boundary circle of the central Möbius band, so  $H_1(A \cap B) = \mathbb{Z} \langle v + w \rangle$ , and  $A$  is homotopy equivalent to the core circle of the central Möbius band, so  $H_1(A) = \mathbb{Z} \langle \sigma_1 + \sigma_2 \rangle$ . Thus  $\psi : v \mapsto \sigma_1 + \sigma_2$  and  $\psi : w \mapsto \sigma_1 + \sigma_2$ .  $\square$

**Remark.** We could define

$$C_k(X; G) = \left\{ \sum_i a_i \sigma_i \mid a_i \in G, \sigma_i : \Delta^k \rightarrow X \right\},$$

for any abelian group  $G$ , with the same differential  $\partial$ , which gives  $H_\bullet(X; G)$ , the **singular homology with coefficients in  $G$** .

**Example.**

$$H_j(S^1; \mathbb{Z}/2) \cong \begin{cases} \mathbb{Z}/2 & j = 0, 1 \\ 0 & \text{otherwise} \end{cases}, \quad H_i(\{\text{point}\}; \mathbb{Z}/2) \cong \begin{cases} \mathbb{Z}/2 & i = 0 \\ 0 & \text{otherwise} \end{cases}.$$

In the previous sequence, if we compute  $H_\bullet(K; \mathbb{Z}/2)$ , get

$$\begin{array}{ccccccc}
 0 \longrightarrow H_2(K; \mathbb{Z}/2) \longrightarrow H_1(A \cap B; \mathbb{Z}/2) & \xrightarrow{\psi} & H_1(A; \mathbb{Z}/2) \oplus H_1(B; \mathbb{Z}/2) & & & & \\
 \parallel & & \parallel & & & & \\
 \mathbb{Z}/2 & \xrightarrow[1 \mapsto (2,2) \equiv (0,0)]{\quad \quad \quad} & \mathbb{Z}/2 \oplus \mathbb{Z}/2 & & & & 
 \end{array}$$

so  $\psi$  vanishes for  $H_\bullet(-; \mathbb{Z}/2)$  and

$$H_i(K; \mathbb{Z}/2) \cong \begin{cases} \mathbb{Z}/2 & i = 0 \\ \mathbb{Z}/2 \oplus \mathbb{Z}/2 & i = 1 \\ \mathbb{Z}/2 & i = 2 \\ 0 & \text{otherwise} \end{cases}.$$

It is also instructive to think about cohomology in this example, where  $K = A \cup B$  for  $A, B \simeq S^1$  and  $A \cap B \simeq S^1$  as before. So the interesting parts of the cohomology Mayer-Vietoris sequences look like

$$\begin{array}{ccccccc}
 H^1(K) & \xrightarrow{(j_A^*, j_B^*)} & H^1(A) \oplus H^1(B) & \xrightarrow{i_A^* - i_B^*} & H^1(A \cap B) & \longrightarrow & H^2(K) \longrightarrow 0 \\
 \parallel & & \parallel & & \parallel & & \\
 \mathbb{Z} \oplus \mathbb{Z} & \xrightarrow[\psi]{\quad \quad \quad} & \mathbb{Z} & & & & 
 \end{array}$$

Check that this  $\psi$  is  $(a, b) \mapsto 2(a - b)$ .<sup>3</sup> So  $H^2(K) \cong \mathbb{Z}/2$ . For contrast,  $H_2(K) = 0$  if we use  $\mathbb{Z}$  coefficients.

**Remark.** There were many ways we could have cut up  $K$ . In some cases, some decompositions will give easier algebra than others.

<sup>3</sup>Exercise

## 2 Structural theorems

Now we should pay some debts.

### 2.1 Chain homotopy

Let  $C_\bullet$  and  $D_\bullet$  be chain complexes.

**Definition.** Chain maps  $f : C_\bullet \rightarrow D_\bullet$  and  $g : C_\bullet \rightarrow D_\bullet$  are **chain homotopic** if there exist  $P_n : C_n \rightarrow D_{n+1}$  such that

$$P_{n-1} \circ \partial_n^{C_\bullet} \pm \partial_{n+1}^{D_\bullet} \circ P_n = f_n - g_n,$$

so

$$\begin{array}{ccccccc} \cdots & \longrightarrow & C_n & \xrightarrow{\partial} & C_{n-1} & \longrightarrow & \cdots \\ & & \searrow P_n & \downarrow & \swarrow P_{n-1} & & \\ \cdots & \longrightarrow & D_{n+1} & \xrightarrow{\partial} & D_n & \longrightarrow & \cdots \end{array}.$$

**Lemma 2.1.** If  $f : C_\bullet \rightarrow D_\bullet$  and  $g : C_\bullet \rightarrow D_\bullet$  are chain homotopic, then

$$(f_*)_i = (g_*)_i : H_i(C_\bullet, \partial) \rightarrow H_i(D_\bullet, \partial),$$

for all  $i$ , that is chain homotopic maps induce the same map on homology.

Recall we are trying to prove if  $f \simeq g : X \rightarrow Y$ , then  $f_* = g_* : H_\bullet(X) \rightarrow H_\bullet(Y)$ . So it will be sufficient to show  $f_\#, g_\# : C_\bullet(X) \rightarrow C_\bullet(Y)$  are chain homotopic.

*Proof.* Let

$$\begin{array}{ccc} & C_n & \xrightarrow{\partial} C_{n-1} \\ & \searrow P_n & \downarrow \\ D_{n+1} & \xrightarrow{\partial} & D_n \end{array},$$

such that  $P_{n-1} \circ \partial \pm \partial \circ P_n = f_n - g_n$ . Let  $\alpha \in C_n$  be a cycle, so  $\partial(\alpha) = 0$ . So  $\partial(f_n(\alpha)) = f_{n-1}(\partial(\alpha)) = 0$ , so  $(f_*)_n([\alpha]) = [f_n(\alpha)]$ . So

$$f_n(\alpha) - g_n(\alpha) = (f_n - g_n)(\alpha) = P_{n-1}(\partial(\alpha)) \pm \partial(P_n(\alpha)) = \partial(P_n(\alpha)) \in \text{im } \partial,$$

so  $[f_n(\alpha)] = [g_n(\alpha)] \in H_n(D_\bullet)$ . □

**Exercise.** Chain homotopy is an equivalence relation on chain complexes and chain maps.

### 2.2 Homotopy invariance

**Theorem 2.2** (Homotopy invariance, version 2). If  $f \simeq g : X \rightarrow Y$  then

$$f_\# \simeq g_\# : (C_\bullet(X), \partial) \rightarrow (C_\bullet(Y), \partial)$$

are chain homotopic.

*Proof.* If  $f \simeq g$ , then there exists  $F : X \times [0, 1] \rightarrow Y$  such that  $F|_{X \times \{0\}} = f$  and  $F|_{X \times \{1\}} = g$ . So if

$$\begin{array}{ccc} \iota_0 : X & \longrightarrow & X \times [0, 1] \\ x & \longmapsto & (x, 0) \end{array}, \quad \begin{array}{ccc} \iota_1 : X & \longrightarrow & X \times [0, 1] \\ x & \longmapsto & (x, 1) \end{array},$$

then  $f = F \circ \iota_0$  and  $g = F \circ \iota_1$ , so  $f_\# = g_\#$  if  $(\iota_0)_\# = (\iota_1)_\#$  and it suffices to prove that  $(\iota_0)_\# \simeq (\iota_1)_\# : C_\bullet(X) \rightarrow C_\bullet(X \times [0, 1])$ , so  $Y$  is out of the picture. So want  $P_n : C_n(X) \rightarrow C_{n+1}(X \times [0, 1])$ . The idea is that  $P_n$  is a **prism operator**

$$\begin{array}{ccc} C_n(X) & \longrightarrow & C_{n+1}(X \times [0, 1]) \\ \sigma : \Delta^n \rightarrow X & \longmapsto & \text{linear combination of simplices for } \sigma \times \text{id} : \Delta^n \times [0, 1] \rightarrow X \times [0, 1] \end{array}.$$

It gives an universal way of cutting up  $\Delta^n \times [0, 1]$  into  $(n + 1)$ -simplices. The equation

$$\partial \circ P \pm P \circ \partial = (\iota_1)_\# - (\iota_0)_\#$$

says that the boundary of the prism is the prism on the boundary plus the top minus the bottom. The details of the proof are not very illuminating, so we will be quite terse. Label the base of the prism by  $[v_0, \dots, v_n]$  and the top  $[w_0, \dots, w_n]$ . Claim that  $\sigma_{n+1}^i = [v_0, \dots, v_i, w_i, \dots, w_n]$  is an  $(n + 1)$ -simplex, and

$$\Delta^n \times [0, 1] = \bigcup_{i=0}^n \sigma_{n+1}^i.$$

We will not prove this, so see Hatcher. Define

$$\begin{aligned} P_n : C_n(X) &\longrightarrow C_{n+1}(X \times [0, 1]) \\ \sigma &\longmapsto \sum_{i=0}^n (-1)^i (\sigma \times \text{id})|_{[v_0, \dots, v_i, w_i, \dots, w_n]} = \sum_{i=0}^n (-1)^i ((\sigma \times \text{id}) \circ \sigma_{n+1}^i) . \end{aligned}$$

Claim that  $\partial \circ P + P \circ \partial = (\iota_1)_\# - (\iota_0)_\#$ . Well,

$$\begin{aligned} \partial(P_n(\sigma)) &= \sum_{j \leq i} (-1)^i (-1)^j (\sigma \times \text{id})|_{[v_0, \dots, \widehat{v_j}, \dots, v_i, w_i, \dots, w_n]} \\ &\quad + \sum_{j \geq i} (-1)^i (-1)^{j+1} (\sigma \times \text{id})|_{[v_0, \dots, v_i, w_i, \dots, \widehat{w_j}, \dots, w_n]} \\ &= (\sigma \times \text{id})|_{[\widehat{v_0}, w_0, \dots, w_n]} - (\sigma \times \text{id})|_{[v_0, \dots, v_n, \widehat{w_n}]} \\ &\quad + \sum_{j < i} (-1)^i (-1)^j (\sigma \times \text{id})|_{[v_0, \dots, \widehat{v_j}, \dots, v_i, w_i, \dots, w_n]} \\ &\quad + \sum_{j > i} (-1)^i (-1)^{j+1} (\sigma \times \text{id})|_{[v_0, \dots, v_i, w_i, \dots, \widehat{w_j}, \dots, w_n]} , \end{aligned}$$

since the  $i = j$  terms cancel in pairs except for  $i = j = 0$ , the top, and  $i = j = n$ , the bottom. Check that the latter sums are  $-P_n(\partial(\sigma))$ ,<sup>4</sup> which is routine but unenlightening.  $\square$

**Remark.** If  $C^\bullet$  and  $D^\bullet$  are cochain complexes,  $f \simeq g$  are **cochain homotopic** if there exist  $P^i : C^i \rightarrow D^{i-1}$  such that

$$\partial^* \circ P \pm P \circ \partial^* = f - g,$$

so

$$\begin{array}{ccccccc} \dots & \xrightarrow{\quad} & C^i & \xrightarrow{\partial^i} & C^{i+1} & \xrightarrow{\quad} & \dots \\ & & \swarrow P^i & \downarrow & \swarrow P^{i+1} & & \\ \dots & \xrightarrow{\quad} & D^{i-1} & \xrightarrow{\partial^{i+1}} & D^i & \xrightarrow{\quad} & \dots \end{array}$$

Check that<sup>5</sup>

$$f^* = g^* : H^\bullet(C^\bullet) \rightarrow H^\bullet(D^\bullet).$$

Then  $P_n : C_n(X) \rightarrow C_{n+1}(X \times [0, 1])$  has dual

$$P^n : \text{Hom}(C_{n+1}(X \times [0, 1]), \mathbb{Z}) = C^{n+1}(X \times [0, 1]) \rightarrow \text{Hom}(C_n(X), \mathbb{Z}) = C^n(X),$$

and  $\partial \circ P + P \circ \partial = (\iota_1)_\# - (\iota_0)_\#$  implies that

$$\partial^* \circ P + P \circ \partial^* = \iota_1^\# - \iota_0^\#,$$

so cohomology is also homotopy invariant.

---

<sup>4</sup>Exercise

<sup>5</sup>Exercise



### 2.3 The long exact sequence

We have made various computations using homotopy invariance, which we have proved, and Mayer-Vietoris, which we have not. Before addressing that, we need some more algebra. Recall that a short exact sequence is an exact sequence of the shape

$$0 \rightarrow A \xrightarrow{\alpha} B \xrightarrow{\beta} C \rightarrow 0, \quad \text{im } \alpha = \ker \beta.$$

**Definition.** A short exact sequence of chain complexes is a diagram

$$\begin{array}{ccccccc} & & \vdots & & \vdots & & \vdots \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & A_{n+1} & \xrightarrow{\alpha} & B_{n+1} & \xrightarrow{\beta} & C_{n+1} \longrightarrow 0 \\ & & \downarrow \partial & & \downarrow \partial & & \downarrow \partial \\ 0 & \longrightarrow & A_n & \xrightarrow{\alpha} & B_n & \xrightarrow{\beta} & C_n \longrightarrow 0, \\ & & \downarrow \partial & & \downarrow \partial & & \downarrow \partial \\ 0 & \longrightarrow & A_{n-1} & \xrightarrow{\alpha} & B_{n-1} & \xrightarrow{\beta} & C_{n-1} \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ & & \vdots & & \vdots & & \vdots \end{array}$$

such that all squares commute, and the columns are chain complexes and the rows are exact, so  $\text{im } \alpha = \ker \beta$  and  $\partial^2 = 0$ . Write

$$0 \rightarrow A_\bullet \rightarrow B_\bullet \rightarrow C_\bullet \rightarrow 0.$$

**Proposition 2.3.** *If*

$$0 \rightarrow A_\bullet \rightarrow B_\bullet \rightarrow C_\bullet \rightarrow 0$$

*is a short exact sequence of chain complexes, there is a boundary map  $\delta : H_n(C_\bullet) \rightarrow H_{n-1}(A_\bullet)$  fitting into a long exact sequence on homology*

$$\cdots \rightarrow H_n(A_\bullet) \xrightarrow{(\alpha_*)_n} H_n(B_\bullet) \xrightarrow{(\beta_*)_n} H_n(C_\bullet) \xrightarrow{\delta} H_{n-1}(A_\bullet) \rightarrow \cdots$$

*Proof.* By diagram chasing, we will construct  $\delta$ , and the proof of exactness is relegated to question sheet 1. Let

$$\begin{array}{ccccccc} & & \vdots & & \vdots & & \vdots \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & A_n & \xrightarrow{\alpha} & B_n & \xrightarrow{\beta} & C_n \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & A_{n-1} & \xrightarrow{\alpha} & B_{n-1} & \xrightarrow{\beta} & C_{n-1} \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & A_{n-2} & \xrightarrow{\alpha} & B_{n-2} & \xrightarrow{\beta} & C_{n-2} \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ & & \vdots & & \vdots & & \vdots \end{array}$$

(Note: A dashed arrow labeled  $\alpha$  points from  $B_n$  to  $A_{n-1}$  in the original image.)

Let  $c_n \in C_n$  be a cycle, so  $\partial(c_n) = 0$ , representing  $[c_n] \in H_n(C_\bullet)$ . Since  $\beta$  is onto, there exists  $b_n \in B_n$  such that  $\beta(b_n) = c_n$ . Since the top right square commutes,  $\beta(\partial(b_n)) = \partial(\beta(b_n)) = \partial(c_n) = 0$ . Since the middle sequence is exact,  $\partial(b_n) \in \ker \beta = \text{im } \alpha$ , so  $\partial(b_n) = \alpha(a_{n-1})$ . Since the bottom left square commutes,  $\alpha(\partial(a_{n-1})) = \partial(\alpha(a_{n-1})) = \partial^2(b_n) = 0$ . Then  $\alpha$  is one-to-one, so  $\alpha(\partial(a_{n-1})) = 0$  implies that  $\partial(a_{n-1}) = 0$ , and set

$$\delta([c_n]) = [a_{n-1}].$$

Check  $\delta$  is well-defined.

- Given  $c_n$ , we chose  $b_n$ . If  $\beta(b'_n) = c_n$ ,  $b_n - b'_n \in \ker \beta = \text{im } \alpha$ , so  $b'_n = b_n + \alpha(a_n)$  for some  $a_n \in A_n$ , and  $\partial(b'_n) = \partial(b_n) + \partial(\alpha(a_n)) = \alpha(a_{n-1} + \partial(a_n))$ , so  $[a_{n-1}] \in H_{n-1}(A_\bullet)$  is unchanged.
- If  $[c_n] = [c'_n]$ , then  $c_n - c'_n \in \text{im } \partial$ , say  $c'_n = c_n + \partial(c_{n+1})$ . Pick  $b_{n+1}$  such that  $\beta(b_{n+1}) = c_{n+1}$  and then  $b_n \mapsto b_n + \partial(b_{n+1})$  and  $\partial(b_n)$  is unchanged, so get the same  $a_{n-1}$ .

So  $\delta$  is well-defined and it is easy to see it is a homomorphism. In the resulting

$$\cdots \rightarrow H_n(A_\bullet) \xrightarrow{(\alpha_*)_n} H_n(B_\bullet) \xrightarrow{(\beta_*)_n} H_n(C_\bullet) \xrightarrow{\delta} H_{n-1}(A_\bullet) \rightarrow \cdots,$$

should check exactness at all three kinds of terms, that is  $\text{im } \beta_* \subseteq \ker \delta$  and  $\ker \delta \subseteq \text{im } \beta_*$ , etc, so six inclusions in total. <sup>6</sup>  $\square$

For this piece of algebra to be useful, we need a source of short exact sequences of chain complexes.

**Example.** Recall if  $G$  is an abelian group,

$$C_k(X; G) = \left\{ \sum_i a_i \sigma_i \mid a_i \in G, \sigma_i : \Delta^k \rightarrow X \right\},$$

which gives  $H_\bullet(X; G)$ , the singular homology with coefficients in  $G$ . Note that if

$$0 \rightarrow G_1 \rightarrow G_2 \rightarrow G_3 \rightarrow 0$$

is a short exact sequence of groups,

$$0 \rightarrow C_\bullet(X; G_1) \rightarrow C_\bullet(X; G_2) \rightarrow C_\bullet(X; G_3) \rightarrow 0$$

is a short exact sequence of chain complexes. The resulting  $\delta : H_n(X; G_3) \rightarrow H_{n-1}(X; G_1)$  is a **Bockstein homomorphism**. For example,

$$0 \rightarrow \mathbb{Z} \xrightarrow{\cdot n} \mathbb{Z} \xrightarrow{p \mapsto p \bmod n} \mathbb{Z}/n \rightarrow 0, \quad 0 \rightarrow \mathbb{Z}/n \xrightarrow{\cdot n} \mathbb{Z}/n^2 \xrightarrow{p \mapsto p \bmod n} \mathbb{Z}/n \rightarrow 0$$

give the **classical Bockstein homomorphisms**

$$H_p(X; \mathbb{Z}/n) \rightarrow H_{p-1}(X; \mathbb{Z}), \quad H_p(X; \mathbb{Z}/n) \rightarrow H_{p-1}(X; \mathbb{Z}/n).$$

We will revisit these later, probably.

## 2.4 Relative homology

**Example.** Let  $A \subseteq X$  be a subspace. We have an inclusion  $C_\bullet(A) \hookrightarrow C_\bullet(X)$  compatible with boundary maps, since if  $\sigma : \Delta^i \rightarrow A \subseteq X$ , then  $\sigma \circ \delta_i : \Delta^{i-1} \rightarrow A$  too. Define

$$C_\bullet(X, A) = C_\bullet(X) / C_\bullet(A),$$

so

$$0 \rightarrow C_\bullet(A) \rightarrow C_\bullet(X) \rightarrow C_\bullet(X, A) \rightarrow 0$$

is a short exact sequence of chain complexes.

**Definition.**  $H_\bullet(C_\bullet(X, A), \partial)$  is denoted  $H_\bullet(X, A)$ , or  $H_\bullet(X, A; G)$ , the **relative homology** of  $(X, A)$ .

---

<sup>6</sup>Exercise: do this

**Lemma 2.4.** *If  $f : (X, A) \rightarrow (Y, B)$  is a map of pairs, that is  $f : X \rightarrow Y$  satisfies  $f(A) \subseteq B$ , then  $f$  induces  $(f_*)_i : H_i(X, A) \rightarrow H_i(Y, B)$  for all  $i$ .*

*Proof.* Elementary. □

The long exact sequence

$$\cdots \rightarrow H_i(A) \rightarrow H_i(X) \rightarrow H_i(X, A) \rightarrow H_{i-1}(A) \rightarrow \cdots$$

is called the **long exact sequence of the pair**  $(X, A)$ .

**Remark.**

- Cycles in  $C_\bullet(X, A)$  are chains in  $X$  whose boundary lies in  $A$ .
- You might expect that things in  $A$  do not matter for  $C_\bullet(X, A)$ , as we quotient all simplices in  $A$ . A precise version of that intuition is excision.

**Theorem 2.5** (Excision). *Let  $X$  be a space,  $A \subseteq X$  a subspace, and  $Z$  a subspace such that  $\bar{Z} \subseteq \mathring{A}$ . Then the inclusion  $\iota : (X \setminus Z, A \setminus Z) \hookrightarrow (X, A)$  is an isomorphism on relative homology, so for all  $n$ ,*

$$(\iota_*)_n : H_n(X \setminus Z, A \setminus Z) \xrightarrow{\sim} H_n(X, A).$$

We will prove excision and Mayer-Vietoris together next time. For now, let us see how this helps us understand relative homology.

**Remark.** Naturality under maps, homotopy invariance, the relative homology long exact sequence, and excision are the key tools of homology and cohomology. Much of what we will do will be built from these.

**Lemma 2.6** (5-lemma). *Suppose*

$$\begin{array}{ccccccccc} A & \xrightarrow{\partial} & B & \xrightarrow{\partial} & C & \xrightarrow{\partial} & D & \xrightarrow{\partial} & E \\ \downarrow \alpha & & \downarrow \beta & & \downarrow \gamma & & \downarrow \delta & & \downarrow \epsilon \\ A' & \xrightarrow{\partial'} & B' & \xrightarrow{\partial'} & C' & \xrightarrow{\partial'} & D' & \xrightarrow{\partial'} & E' \end{array}$$

*is a commuting diagram of abelian groups with exact rows. If  $\alpha, \beta, \delta, \epsilon$  are isomorphisms, then so is  $\gamma$ .*

*Proof.* More diagram chasing. We will show  $\gamma$  is one-to-one, and you check it is onto. Let  $c \in C$  have  $\gamma(c) = 0$ . Then  $\delta(\partial(c)) = \partial'(\gamma(c)) = 0$  so  $\partial(c) \in \ker \delta$ , and  $\delta$  is an isomorphism so  $\partial(c) = 0$ . Since the rows are exact,  $c \in \ker \partial = \text{im } \partial$ , so  $c = \partial(b)$  for  $b \in B$ . Then  $\partial'(\beta(b)) = \gamma(\partial(b)) = \gamma(c) = 0$ , so  $\beta(b) \in \ker \partial' = \text{im } \partial'$ , and  $\beta(b) = \partial'(a')$ . Since  $\alpha$  is an isomorphism, there exists  $a \in A$  such that  $\alpha(a) = a'$ . Now  $\beta(\partial(a)) = \partial'(\alpha(a)) = \partial'(a') = \beta(b)$  so  $\partial(a) - b \in \ker \beta$ , and  $\beta$  is an isomorphism so  $b = \partial(a)$ . Thus  $c = \partial(b) = \partial^2(a) = 0$  and  $c$  is one-to-one. □

**Corollary 2.7.** *If  $f : (X, A) \rightarrow (Y, B)$  is a map of pairs, and any two of the induced homomorphisms*

$$H_\bullet(X) \rightarrow H_\bullet(Y), \quad H_\bullet(A) \rightarrow H_\bullet(B), \quad H_\bullet(X, A) \rightarrow H_\bullet(Y, B)$$

*are isomorphisms, then so is the third.*

*Proof.* Apply the 5-lemma to

$$\begin{array}{ccccccccccc} \cdots & \longrightarrow & H_i(A) & \longrightarrow & H_i(X) & \longrightarrow & H_i(X, A) & \longrightarrow & H_{i-1}(A) & \longrightarrow & H_{i-1}(X) & \longrightarrow & \cdots \\ & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \\ \cdots & \longrightarrow & H_i(B) & \longrightarrow & H_i(Y) & \longrightarrow & H_i(Y, B) & \longrightarrow & H_{i-1}(B) & \longrightarrow & H_{i-1}(Y) & \longrightarrow & \cdots \end{array}$$

□

## 2.5 Reduced homology and good pairs

We need two definitions to proceed. The first looks a bit odd, but be patient.

**Definition.** If  $X$  is a space, and  $x_0 \in X$  is a basepoint, the **reduced homology** is

$$\widetilde{H}_i(X) = H_i(X, x_0).$$

**Exercise.** The long exact sequence of a pair shows

$$\widetilde{H}_0(X) \oplus \mathbb{Z} \cong H_0(X), \quad \widetilde{H}_i(X) \cong H_i(X), \quad i > 0.$$

**Definition.** A pair  $(X, A)$  is **good** if  $A \subseteq X$  is closed and is a deformation retract of an open neighbourhood  $A \subseteq U \subseteq X$ , that is there exists  $H : [0, 1] \times U \rightarrow U$  such that

- $H|_{\{0\} \times U} = \text{id}$  and  $H|_{\{1\} \times U}$  has image in  $A$ , and
- $H$  is fixed on  $A$ , so for all  $t \in [0, 1]$  and  $a \in A$ ,  $H(t, a) = a$ .

So you can squeeze  $U$  back onto  $A$  without moving  $A$ . If  $X$ , and hence  $U$ , is Hausdorff,  $A$  is automatically closed.

**Proposition 2.8.** *If  $(X, A)$  is good, the natural map  $(X, A) \rightarrow (X/A, A/A)$  induces isomorphisms*

$$H_\bullet(X, A) \xrightarrow{\sim} \widetilde{H}_\bullet(X/A).$$

*Proof.* Note that homotopy invariance and the 5-lemma show inclusion defines isomorphisms

$$H_\bullet(A) \xrightarrow{\sim} H_\bullet(U), \quad H_\bullet(X, A) \xrightarrow{\sim} H_\bullet(X, U).$$

The inclusion  $A/A = \{\text{point}\} \hookrightarrow U/A$  is a deformation retract and in particular a homotopy equivalence, so

$$H_\bullet(X/A, A/A) \xrightarrow{\sim} H_\bullet(X/A, U/A)$$

is also an isomorphism by the 5-lemma. Consider

$$\begin{array}{ccccc} H_\bullet(X, A) & \xrightarrow[\sim]{\text{Homotopy}} & H_\bullet(X, U) & \xleftarrow[\sim]{\text{Excision}} & H_\bullet(X \setminus A, U \setminus A) \\ \downarrow & & & & \downarrow \\ H_\bullet(X/A, A/A) & \xrightarrow[\text{Homotopy}]{\sim} & H_\bullet(X/A, U/A) & \xleftarrow[\text{Excision}]{\sim} & H_\bullet((X/A) \setminus (A/A), (U/A) \setminus (A/A)) \end{array},$$

where the vertical maps collapse  $A$ . Then the right vertical map is a homeomorphism of pairs, since  $X \setminus A \cong (X/A) \setminus (A/A)$ . So the right vertical map is an isomorphism and hence the left vertical map is an isomorphism.  $\square$

**Remark.** The **tubular neighbourhood theorem** of differential topology, which we will discuss more later, implies that if  $X$  is a smooth manifold and  $A \subseteq X$  is a compact smooth submanifold,  $(X, A)$  is a good pair.

**Example.**

$$H_j(D^n, \partial D^n) \cong \widetilde{H}_j(D^n / \partial D^n) = \widetilde{H}_j(S^n) = \begin{cases} \mathbb{Z} & j = n \\ 0 & \text{otherwise} \end{cases}.$$

**Example.** Let  $S^1$  be the equator. Then

$$H_j(S^2, S^1) \cong \widetilde{H}_j(S^2 \vee S^2) \cong \begin{cases} \mathbb{Z} \oplus \mathbb{Z} & j = 2 \\ 0 & \text{otherwise} \end{cases}.$$

**Remark.** If  $M$  is a manifold and  $x \in M$ , by excision with  $Z = M \setminus \{\text{open disc neighbourhood of } x\}$  and homotopy invariance or directly from the long exact sequence of a pair,

$$H_j(M, M \setminus \{x\}) \cong H_j(\mathbb{R}^n, \mathbb{R}^n \setminus \{0\}) \cong H_j(D^n, \partial D^n) \cong \begin{cases} \mathbb{Z} & j = n = \dim_{\mathbb{R}} M \\ 0 & \text{otherwise} \end{cases}.$$

## 2.6 Mayer-Vietoris and excision

We have stated two major properties of homology and cohomology without proof, Mayer-Vietoris and excision. Recall that we also saw if

$$0 \rightarrow A_\bullet \rightarrow B_\bullet \rightarrow C_\bullet \rightarrow 0$$

is a short exact sequence of chain complexes, then there exists a long exact sequence in homology

$$\cdots \rightarrow H_i(A_\bullet) \rightarrow H_i(B_\bullet) \rightarrow H_i(C_\bullet) \rightarrow H_{i-1}(A_\bullet) \rightarrow \cdots$$

Mayer-Vietoris will be a consequence of this.

**Definition.** Let  $\mathcal{U} = \{U_\alpha\}_{\alpha \in I}$  be a collection of subsets of  $X$  with the property that  $X = \bigcup_{\alpha \in I} U_\alpha$ , such as an open cover. Then

$$C_j^\mathcal{U}(X) = \left\{ \sum_i a_i \sigma_i \mid a_i \in \mathbb{Z}, \sigma_i : \Delta^j \rightarrow X, \exists \alpha(i) \in I, \text{im } \sigma_i \subseteq U_{\alpha(i)} \right\}$$

is the **subcomplex** of  $(C_\bullet(X), \partial)$  generated by simplices each of which lie wholly inside some set in  $\mathcal{U}$ .

Note that

$$\begin{array}{ccc} C_\bullet(X) & \longrightarrow & C_{\bullet-1}(X) \\ \cup & & \cup \\ C_\bullet^\mathcal{U}(X) & \longrightarrow & C_{\bullet-1}^\mathcal{U}(X) \end{array},$$

since  $C_\bullet^\mathcal{U}(X)$  is preserved by  $\partial$  so is a subcomplex.

**Proposition 2.9** (Small simplices theorem). *The inclusion  $C_\bullet^\mathcal{U}(X) \hookrightarrow C_\bullet(X)$  induces an isomorphism on homology.*

**Remark.** Suppose  $f : X \rightarrow Y$  sends each element of  $\mathcal{U}$  into some element of  $\mathcal{V}$ , the corresponding cover of  $Y$ . Then  $f$  induces  $f_\# : C_\bullet^\mathcal{U}(X) \rightarrow C_\bullet^\mathcal{V}(Y)$ .

**Example** (Mayer-Vietoris). Let  $\mathcal{U} = \{A, B\}$  for  $A, B \subseteq X$  open. Then there is an obvious short exact sequence of chain complexes

$$0 \rightarrow C_\bullet(A \cap B) \xrightarrow{\sigma \mapsto (\sigma, \sigma)} C_\bullet(A) \oplus C_\bullet(B) \xrightarrow{(u, v) \mapsto u - v} C_\bullet^\mathcal{U}(X) \rightarrow 0,$$

which is onto since  $C_\bullet^\mathcal{U}(X)$  only contains simplices lying in  $A$  or  $B$ . The associated long exact sequence is the Mayer-Vietoris sequence, using small simplices to identify  $H_\bullet(C_\bullet^\mathcal{U}(X)) \xrightarrow{\sim} H_\bullet(C_\bullet(X))$ . Note also the construction of the  $\partial$  map in the long exact sequence associated to a short exact sequence of complexes does reproduce our earlier description of  $\partial_{MV}$ . Also the naturality of Mayer-Vietoris under maps  $f : X \rightarrow Y$  such that  $f(A) \subseteq C$  and  $f(B) \subseteq D$  is just the naturality of  $C_\bullet^\mathcal{U}(X) \rightarrow C_\bullet^\mathcal{V}(Y)$ .

**Example** (Excision). Recall we have  $Z, A \subseteq X$  and  $\bar{Z} \subseteq \mathring{A}$ . Let  $B = X \setminus Z$  and let  $\mathcal{U} = \{A, B\}$ , so the interiors of  $A$  and  $B$  do cover  $X$ . Note that

$$C_n^\mathcal{U}(X) / C_n(A) \cong C_n(B) / C_n(A \cap B)$$

is the free abelian group on simplices in  $B$  not wholly contained in  $A$ . The short exact sequences of chain complexes

$$\begin{array}{ccccccc} 0 & \longrightarrow & C_\bullet(A) & \longrightarrow & C_\bullet^\mathcal{U}(X) & \longrightarrow & C_\bullet^\mathcal{U}(X) / C_\bullet(A) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & C_\bullet(A) & \longrightarrow & C_\bullet(X) & \longrightarrow & C_\bullet(X) / C_\bullet(A) \longrightarrow 0 \end{array},$$

and the natural map of short exact sequences give a map of long exact sequences

$$\begin{array}{ccccccccccc} \cdots & \rightarrow & H_i(A) & \rightarrow & H_i(C_\bullet^\mathcal{U}(X)) & \rightarrow & H_i(C_\bullet^\mathcal{U}(X) / C_\bullet(A)) & \rightarrow & H_{i-1}(A) & \rightarrow & H_{i-1}(C_\bullet^\mathcal{U}(X)) & \rightarrow \cdots \\ & & \downarrow = & & \downarrow \sim \text{ss} & & \downarrow \phi & & \downarrow \sim \text{ss} & & \downarrow = & \\ \cdots & \rightarrow & H_i(A) & \longrightarrow & H_i(X) & \longrightarrow & H_i(X, A) & \longrightarrow & H_{i-1}(A) & \longrightarrow & H_{i-1}(X) & \longrightarrow \cdots \end{array}.$$

Lecture 7  
Friday  
23/10/20

So by the 5-lemma,  $\phi$  is an isomorphism, so

$$C_{\bullet}^{\mathcal{U}}(X)/C_{\bullet}(A) \hookrightarrow C_{\bullet}(X)/C_{\bullet}(A)$$

is an isomorphism on homology. So

$$\begin{aligned} H_{\bullet}(X, A) &= H_{\bullet}(C_{\bullet}(X)/C_{\bullet}(A)) \cong H_{\bullet}(C_{\bullet}^{\mathcal{U}}(X)/C_{\bullet}(A)) \\ &\cong H_{\bullet}(C_{\bullet}(B)/C_{\bullet}(A \cap B)) = H_{\bullet}(B, A \cap B) = H_{\bullet}(X \setminus Z, A \setminus Z), \end{aligned}$$

proving excision.

## 2.7 Small simplices theorem

So it just remains to prove the small simplices theorem that  $C_{\bullet}^{\mathcal{U}}(X) \hookrightarrow C_{\bullet}(X)$  is an isomorphism on homology. The key geometric ingredient is to divide simplices into smaller simplices.

**Definition.** The **barycentre**, or centre of mass, of  $\Delta^n$  is

$$b_n = \frac{(1, \dots, 1)}{n+1}.$$

A **barycentric subdivision** is the following three-step procedure.

- Subdivide the boundary.
- Add the barycentre.
- Cone off from the barycentre to the subdivided boundary.

**Definition.** If  $\sigma : \Delta^i \rightarrow \Delta^n \in C_i(\Delta^n)$ ,

$$\begin{aligned} \text{Cone}_i^{\Delta^n}(\sigma) : \quad \Delta^{i+1} &\longrightarrow \Delta^n \\ (t_0, \dots, t_{i+1}) &\longmapsto t_0 b_n + (1 - t_0) \sigma \left( \frac{(t_1, \dots, t_{i+1})}{1 - t_0} \right). \end{aligned}$$

So, extended linearly,  $\text{Cone}_i^{\Delta^n} : C_i(\Delta^n) \rightarrow C_{i+1}(\Delta^n)$ .

**Exercise.**

$$\partial \left( \text{Cone}_i^{\Delta^n}(\sigma) \right) = \begin{cases} \sigma - \text{Cone}_{i-1}^{\Delta^n}(\partial(\sigma)) & i > 0 \\ \sigma - \epsilon(\sigma) b_n & i = 0 \end{cases},$$

where

$$\begin{aligned} \epsilon : C_0(\Delta^n) &\longrightarrow \mathbb{Z} \\ \sum_i n_i p_i &\longmapsto \sum_i n_i \end{aligned}$$

is the augmentation.

**Definition.** Define

$$\begin{aligned} c : C_{\bullet}(\Delta^n) &\longrightarrow C_{\bullet}(\Delta^n) \\ \sigma &\longmapsto \begin{cases} \epsilon(\sigma) b_n & \text{on } C_0(\Delta^n) \\ 0 & \text{on } C_i(\Delta^n), i > 0 \end{cases}. \end{aligned}$$

Then

$$\partial \circ \text{Cone}^{\Delta^n} + \text{Cone}^{\Delta^n} \circ \partial = \text{id}_{C_{\bullet}(\Delta^n)} - c.$$

**Definition.** A collection of chain maps  $\phi^X : C_{\bullet}(X) \rightarrow C_{\bullet}(X)$ , defined for all spaces  $X$ , is **natural** if for all  $f : X \rightarrow Y$ ,

$$f_{\#} \circ \phi^X = \phi^Y \circ f_{\#}.$$

Similarly for a collection  $P : C_{\bullet}(X) \rightarrow C_{\bullet+1}(X)$  of chain homotopies between natural  $\phi^X$  and  $\psi^X$ .

**Definition.** Define

$$\begin{aligned} \phi_0^X &= \text{id}_{C_0(X)}, & \phi_n^X &: C_n(X) \longrightarrow C_n(X) \\ \sigma &\longmapsto \sigma_{\#} \left( \text{Cone}_{n-1}^{\Delta^n} (\phi_{n-1}^{\Delta^n} (\partial(\iota_n))) \right), \end{aligned}$$

where  $\iota_n : \Delta^n \rightarrow \Delta^n \in C_n(\Delta^n)$  is the identity, so  $\partial(\iota_n) \in C_{n-1}(\Delta^n)$ .

Since  $\sigma : \Delta^n \rightarrow X$  is  $\sigma \circ \iota_n : \Delta^n \rightarrow \Delta^n \rightarrow X$ , this is natural, since

$$\phi_n^X(\sigma) = \phi_n^X(\sigma_{\#}(\iota_n)) = \sigma_{\#}(\phi_n^{\Delta^n}(\iota_n)).$$

The idea is that we know how to subdivide  $\Delta^n$ , so know how to subdivide any simplex in  $X$ .

**Definition.** Similarly, define

$$\begin{aligned} P_n^X &: C_n(X) \longrightarrow C_{n+1}(X) \\ \sigma &\longmapsto \sigma_{\#} \left( \text{Cone}_n^{\Delta^n} (\phi_n^{\Delta^n}(\iota_n) - \iota_n - P_{n-1}^{\Delta^n}(\partial(\iota_n))) \right). \end{aligned}$$

This decomposes the prism  $\Delta^n \times [0, 1]$  by joining  $\Delta^n \times \{0\}$  and  $\Delta^n \times \{1\}$  to the barycentre of  $\Delta^n \times \{1\}$ .

**Fact.**  $\phi^X : C_{\bullet}(X) \rightarrow C_{\bullet}(X)$  is a natural chain map, and  $P^X : C_{\bullet}(X) \rightarrow C_{\bullet+1}(X)$  is a natural chain homotopy from  $\phi^X$  to the identity, that is

$$\partial \circ P_n^X + P_{n-1}^X \circ \partial = \phi_n^X - \text{id}_{C_n(X)}.$$

We will not prove this.

Ok, now we know how to divide simplices.

**Lemma 2.10.** *If  $[v_0, \dots, v_n] \subseteq \mathbb{R}^{n+1}$  is a simplex, then each simplex of its barycentric division has Euclidean diameter at most  $n/(n+1)$  the Euclidean diameter of  $[v_0, \dots, v_n]$ .*

**Corollary 2.11.**

1. If  $\sigma \in C_n^{\mathcal{U}}(X)$ ,  $\phi_n^X(\sigma) \in C_n^{\mathcal{U}}(X)$ .
2. If  $\sigma \in C_n(X)$ , there exists  $k \gg 0$  such that  $(\phi_n^X)^k(\sigma) \in C_n^{\mathcal{U}}(X)$ .

*Proof.*

1. Obvious.
2.  $\sigma$  is a finite sum of simplices, so it suffices to prove the result for one  $\sigma : \Delta^n \rightarrow X$ . Let  $\mathcal{U} = \{U_{\alpha}\}_{\alpha \in I}$ . Now  $\left\{ \sigma^{-1} \left( U_{\alpha}^{\circ} \right) \right\}_{\alpha \in I}$  is an open cover of  $\Delta^n$ , so has a Lebesgue number, that is there exists  $\epsilon > 0$  such that any open  $\epsilon$ -ball in  $\Delta^n$  lies in some  $\sigma^{-1}(U_{\alpha})$ . Now pick  $k \gg 0$  such that  $(n/(n+1))^k \ll \epsilon$ .

□

*Proof of Proposition 2.9.* Let  $U : H_{\bullet}(C_{\bullet}^{\mathcal{U}}(X)) \rightarrow H_{\bullet}(X)$  be the natural map.

- If  $[c] \in H_n(X)$ , there exists  $k$  such that  $(\phi_n^X)^k(c) \in C_n^{\mathcal{U}}(X)$ . Since  $\phi^X \simeq \text{id}$ ,  $(\phi^X)^k \simeq \text{id}$ , so there exists  $F$  such that  $\partial \circ F + F \circ \partial = (\phi^X)^k - \text{id}$ . Then  $(\phi^X)^k(c) = c + \text{im } \partial$ , so  $U$  is onto.
- If  $U([c]) = 0$  for  $[c] \in H_n(C_{\bullet}^{\mathcal{U}}(X))$  and  $z \in C_{n+1}(X)$  has  $\partial(z) = c$ , there exists  $k$  such that  $(\phi_{n+1}^X)^k(z) \in C_{n+1}^{\mathcal{U}}(X)$  and  $(\phi_{n+1}^X)^k(z) - z = (\partial \circ F + F \circ \partial)(z)$ , so

$$c = \partial(z) = \partial \left( (\phi_{n+1}^X)^k(z) \right) - \partial(F(\partial(z))) \in C_{n+1}^{\mathcal{U}}(X),$$

since  $\partial(z) \in C_n^{\mathcal{U}}(X)$  and  $F$  is natural. Then  $c \in \text{im}(\partial : C_{n+1}^{\mathcal{U}}(X) \rightarrow C_n^{\mathcal{U}}(X))$ , so  $[c] = 0$  and  $U$  is one-to-one.

□

### 3 Cellular homology and cohomology

Singular homology and cohomology are defined for all topological spaces, but we are mostly interested in nice spaces. In particular, we have seen  $H_\bullet(S^n)$ ,  $H_\bullet(\text{Klein})$ ,  $H_\bullet(\Sigma_g)$ , and  $H^\bullet(\mathbb{CP}^n)$  are all finite rank, even though  $C_\bullet(X)$  is vast in each case. Our next goal is to develop a computational shortcut which makes this manifest.

#### 3.1 Cell complexes

**Definition.** A **cell complex**, or **CW complex**, is a space obtained inductively as follows.

- $X_0$  is a discrete set, such as a finite set.
- Given  $X_{k-1}$ ,

$$X_k = X_{k-1} \cup \bigcup_{i \in I_k} D_i^k,$$

for  $I_k$  an indexing set and  $D_i^k = \{x \in \mathbb{R}^k \mid \|x\| \leq 1\}$  a closed disc, called  **$k$ -cells**, attached via  $\partial D_i^k = S^{k-1} \rightarrow X_{k-1}$ , so  $X_{k-1} \sqcup \bigsqcup_{i \in I_k} D_i^k \rightarrow X_k$  is the quotient map identifying  $\partial D_i^k$  and its image.

- $X = \bigcup_{k \geq 0} X_k$  with the **weak** topology, where  $U \subseteq X$  is open if and only if  $U \cap X_k$  is open in  $X_k$  for all  $k$ .

**Example.**

- $S^n = \{\text{point}\} \cup D^n$  attached via the constant map  $\partial D^n \rightarrow \{\text{point}\}$ .
- $S^n = \{\text{point}\} \cup \{\text{point}\} \cup D_{\alpha_1}^1 \cup D_{\alpha_2}^1 \cup D_{\beta_1}^2 \cup D_{\beta_2}^2$ .
- $T^2$  has one 0-cell, two 1-cells, and one 2-cell.
- $\Sigma_2$  has one 0-cell, four 1-cells, and one 2-cell.
- The **wedge product**. If  $X$  and  $Y$  are cell complexes,  $X \vee Y = \langle X \sqcup Y \rangle / x_0 \sim y_0$  where  $x_0 \in X_0$  and  $y_0 \in Y_0$ .

**Notation.** Let  $X$  be a cell complex. The  $D_i^k$  are  $k$ -cells.

- $X_k$  is the  **$k$ -skeleton** of  $X$ .
- If there exists  $N$  such that  $X = X_N$ ,  $X$  is a **finite-dimensional** cell complex.
- If  $X = X_N$  and  $I_j < \infty$  for all  $j$ ,  $X$  is a **finite** cell complex. Then  $X$  is compact.
- $X = \bigsqcup_{k \geq 0} \mathring{D}_\alpha^k$  is the disjoint union of its open cells  $C_\alpha^k$  as attaching maps take  $\partial D_\alpha^k$  to  $X_{k-1}$ .
- A **subcomplex**  $A \subseteq X$  is a closed subspace which is a union of cells of  $X$ . Note that given a cell complex  $X$ , you cannot throw out a random bunch of cells to get a subcomplex. There may be later cells that try to attach to things you are throwing out.

#### 3.2 Point-set digression

Let  $X = \bigcup_{n \geq 0} X_n$  be a cell complex.

**Exercise.**  $A \subseteq X$  is open, or closed, if and only if  $(\phi_\alpha^n)^{-1}(A) \subseteq D_\alpha^n$  is open, or closed, for all  $\alpha$ , where

$$\phi_\alpha^n : D_\alpha^n \hookrightarrow X_{n-1} \sqcup \bigsqcup_{\alpha} D_\alpha^n \twoheadrightarrow X_n \hookrightarrow X$$

is the **characteristic** map of the cell, so  $\phi_\alpha^n|_{\partial D_\alpha^n}$  is the attaching map.



Let  $A \subseteq X$ . We build an open neighbourhood  $N_\epsilon(A)$  of  $A$  inductively. Let  $N_\epsilon^0(A) = A \cap X_0$ . Given  $N_\epsilon^n(A) \subseteq X_n$  an open neighbourhood of  $A \cap X_n$ , define  $N_\epsilon^{n+1}(A)$  by specifying

$$\begin{aligned} (\phi_\alpha^{n+1})^{-1}(N_\epsilon^{n+1}(A)) = & \left( \text{open } \epsilon\text{-neighbourhood of } (\phi_\alpha^n)^{-1}(A) \setminus \partial D_\alpha^{n+1} \subseteq D_\alpha^{n+1} \setminus \partial D_\alpha^{n+1} \right) \\ & \cup \left( (1 - \epsilon, 1] \times (\phi_\alpha^n)^{-1}(N_\epsilon^n(A)) \right), \end{aligned}$$

where  $\epsilon$  depends on  $\alpha$ , and  $(1 - \epsilon, 1]$  is the radial spherical coordinate on  $D_\alpha^{n+1}$  and  $(\phi_\alpha^n)^{-1}(N_\epsilon^n(A))$  is the angular coordinate on  $\partial D_\alpha^{n+1}$ . Then  $N_\epsilon(A) = \bigcup_{n \geq 0} N_\epsilon^n(A)$  is open, as it is open in every cell.

**Proposition 3.1.** *Cell complexes are Hausdorff and locally contractible. So connected if and only if path-connected.*

*Proof.* For a proof, see Hatcher, appendix A. □

**Fact.** A compact smooth manifold, perhaps with boundary, is homotopy equivalent to a finite cell complex. And given  $N \subseteq M$  a properly embedded submanifold, there exists a cell structure on  $M$  making  $N$  a subcomplex. Can drop smoothness, but there are nice proofs using Morse theory if you have it.

**Lemma 3.2.** *If  $X$  is a cell complex and  $A \subseteq X$  is a subcomplex, then  $(X, A)$  is a good pair.*

*Proof.* See Hatcher. Again, point-set rather than algebraic topology. □

**Corollary 3.3.**  $H_\bullet(X, A) \cong \widetilde{H}_\bullet(X/A)$ .

**Corollary 3.4.** *If  $X = \bigcup_{k \geq 0} X_k$  is a cell complex,*

$$H_i(X_k, X_{k-1}) = \begin{cases} \bigoplus_{\alpha \in I_k} \mathbb{Z} & i = k \\ 0 & \text{otherwise} \end{cases}$$

*is free abelian on the set of  $k$ -cells in  $X$ .*

*Proof.*  $X_{k-1} \subseteq X_k$  is a subcomplex, so

$$H_\bullet(X_k, X_{k-1}) \cong \widetilde{H}_\bullet(X_k/X_{k-1}) \cong \widetilde{H}_\bullet\left(\bigvee_{\alpha \in I_k} S_\alpha^k\right),$$

as  $\partial D_\alpha^k \rightarrow X_{k-1}$  for all  $k$ -cells and  $X_{k-1}$  is collapsed to a point. Choose  $x_\alpha \in S_\alpha^k$  for all  $\alpha$ . Then  $(\bigsqcup_\alpha S_\alpha^k, \bigsqcup_\alpha \{x_\alpha\})$  is a good pair and  $\bigsqcup_\alpha S_\alpha^k / \bigsqcup_\alpha \{x_\alpha\} = \bigvee_\alpha S_\alpha^k$ , so

$$H_\bullet(X_k, X_{k-1}) \cong H_\bullet\left(\bigsqcup_\alpha S_\alpha^k, \bigsqcup_\alpha \{x_\alpha\}\right) = \bigoplus_\alpha H_\bullet(S_\alpha^k, \{x_\alpha\}) = \bigoplus_\alpha \widetilde{H}_\bullet(S_\alpha^k).$$

□

**Proposition 3.5.** *If  $Z \subseteq X$  is compact, there exists  $N$  such that  $Z \subseteq X_N$ .*

*Proof.* We will show  $Z$  meets only finitely many cells of  $X$ . Suppose for contradiction there exists  $S = \{x_0, x_1, \dots\} \subseteq Z$  such that  $x_i \in e_i$  and the cells  $\{e_i\}$  are pairwise distinct. Claim that  $S$  is closed in  $X$ . Well,  $S \cap X_0$  is closed in  $X_0$ , a discrete space. Inductively, if  $S \cap X_{n-1}$  is closed in  $X_{n-1}$  and  $\phi_\alpha^n : D_\alpha^n \rightarrow X_n$  is an  $n$ -cell,  $(\phi_\alpha^n|_{\partial D_\alpha^n})^{-1}(S) \subseteq S_\alpha^{n-1}$  is closed. Then

$$(\phi_\alpha^n)^{-1}(S) = (\phi_\alpha^n|_{\partial D_\alpha^n})^{-1}(S) \cup \{\text{at most one point}\} \subseteq D_\alpha^n$$

is closed, since  $X$  is the disjoint union of interiors of cells, so  $S$  meets each cell of  $X_n$  in a closed set, so  $S \subseteq X_n$  is closed. Same for  $S' \subseteq S$ , so  $S$  is discrete. Thus  $S$  is finite. □

**Proposition 3.6.** Let  $X = \bigcup_{k \geq 0} X_k$  be a cell complex.

1.  $H_k(X_n) = 0$  for all  $k > n$ .
2. The inclusion  $X_n \hookrightarrow X$  induces an isomorphism  $H_j(X_n) \xrightarrow{\sim} H_j(X)$  for all  $j < n$ .

*Proof.*

1. If  $k > n$ , the long exact sequence of a pair  $(X_n, X_{n-1})$  gives

$$\begin{array}{ccccccc} H_{k+1}(X_n, X_{n-1}) & \longrightarrow & H_k(X_{n-1}) & \longrightarrow & H_k(X_n) & \longrightarrow & H_k(X_n, X_{n-1}) \\ \parallel & & & & & & \parallel \\ 0 & & & & & & 0 \end{array},$$

so  $H_k(X_n) \cong \dots \cong H_k(X_0) = 0$ , since  $X_0$  is a discrete set.

2. The same sequence as before

$$H_{k+1}(X_n, X_{n-1}) \rightarrow H_k(X_{n-1}) \rightarrow H_k(X_n) \rightarrow H_k(X_n, X_{n-1})$$

with  $k < n - 1$  shows  $H_k(X_{n-1}) \cong \dots \cong H_k(X_N)$  for all  $N > n - 1$ . If  $X$  is finite-dimensional, we are done. In general, if  $\alpha \in H_k(X)$ ,  $\alpha$  is represented by a finite union of simplices, which is compact. If  $Z \subseteq X$  is compact, there exists  $N$  such that  $Z \subseteq X_N$ . So  $\alpha \in \text{im}(\phi_N : H_k(X_N) \rightarrow H_k(X))$  for all  $N \gg 0$ . Similarly, if a cycle  $\alpha = \sum_i a_i \sigma_i \in H_k(X)$  bounds a  $(k+1)$ -chain in  $X$ , that  $(k+1)$ -chain lives in some  $X_{N'}$  for  $N' \gg 0$ , so  $[\alpha] = 0 \in H_k(X_{N'})$ , that is  $\phi_N$  is one-to-one for  $N \gg 0$ . □

**Corollary 3.7.** Let  $X$  be a finite-dimensional cell complex of dimension  $n$ . Then

$$H_j(X) = 0, \quad j > n.$$


### 3.3 Cellular homology

We are still computing from the huge chain groups  $C_\bullet(X)$ . The goal is to describe a model for  $H_\bullet(X)$  starting from a much smaller chain complex.

**Definition.** Let  $X = \bigcup_{n \geq 0} X_n$  be a cell complex. Let

$$C_k^{\text{cell}}(X) = H_k(X_k, X_{k-1}).$$

This is free abelian on the  $k$ -cells. Then



Observe that  $\partial_k^{\text{cell}} \circ \partial_{k+1}^{\text{cell}} = 0$  since we have two consecutive maps from one long exact sequence. This is the **cellular chain complex** of  $X = \bigcup_{n \geq 0} X_n$ . Note that it depends on the chosen cell structure. Write  $H_\bullet^{\text{cell}}(X)$  for  $H_\bullet(C_\bullet^{\text{cell}}(X), \partial_\bullet^{\text{cell}})$ .

Lecture 9  
Wednesday  
28/10/20

**Proposition 3.8.**

$$H_{\bullet}^{\text{cell}}(X) \cong H_{\bullet}(X).$$

*Proof.* Recall that  $H_j(X_k) = 0$  if  $j > k$  and  $X_k \hookrightarrow X$  is an isomorphism on homology for  $j < k$ . Then

$$\begin{array}{ccccccc}
 & & & & H_k(X_{k+1}, X_k) = 0 & & \\
 & & & & \nearrow & & \\
 & & H_k(X_{k+1}) \cong H_k(X) & & & & \\
 & \nearrow & & \searrow & & & \\
 0 = H_k(X_{k-1}) & & H_k(X_k) & & & & \\
 & \nearrow \partial_{k+1} & & \searrow (i_k)_* & & & \\
 H_{k+1}(X_{k+1}, X_k) & \xrightarrow{\partial_{k+1}^{\text{cell}}} & H_k(X_k, X_{k-1}) & \xrightarrow{\partial_k^{\text{cell}}} & H_{k-1}(X_{k-1}, X_{k-2}) & & \\
 & & \searrow \partial_k & & \nearrow (i_{k-1})_* & & \\
 & & H_{k-1}(X_{k-1}) & & & & \\
 & & \nearrow & & & & \\
 & & 0 = H_{k-1}(X_{k-2}) & & & & 
 \end{array}$$

so

$$\begin{aligned}
 H_k(X) &\cong H_k(X_{k+1}) \cong H_k(X_k) / \text{im } \partial_{k+1} \cong \text{im } (i_k)_* / \text{im } ((i_k)_* \circ \partial_{k+1}) && \text{since } (i_k)_* \text{ is injective} \\
 &\cong \ker \partial_k / \text{im } \partial_{k+1}^{\text{cell}} \cong \ker ((i_{k-1})_* \circ \partial_k) / \text{im } \partial_{k+1}^{\text{cell}} && \text{since } (i_{k-1})_* \text{ is injective} \\
 &= \ker \partial_k^{\text{cell}} / \text{im } \partial_{k+1}^{\text{cell}} = H_k^{\text{cell}}(X).
 \end{aligned}$$

□

**Remark.** If  $X$  and  $Y$  are cell complexes and  $f : X \rightarrow Y$  is a map, in general  $f$  does not induce maps  $C_{\bullet}^{\text{cell}}(X) \rightarrow C_{\bullet}^{\text{cell}}(Y)$ . Ok if  $f$  is **cellular**, so  $f$  takes a  $k$ -skeleton  $X_k$  into a  $k$ -skeleton  $Y_k$ , for all  $k$ .

The following are immediate.

**Corollary 3.9.** *Let  $X$  be a finite cell complex.*

- $H_k(X)$  is a finitely generated abelian group of rank at most  $n_k$ , the number of  $k$ -cells.
- If  $H_k(X) \neq 0$ , every cell structure on  $X$  must have at least  $\text{rk } H_k(X)$  distinct  $k$ -cells.
- If  $X$  admits a cell structure with only even-dimensional cells,  $H_{\bullet}(X) \cong C_{\bullet}^{\text{cell}}(X)$  for this cell structure.
- $H_{\bullet}(X; \mathbb{F})$  is a finite-dimensional vector space over the field  $\mathbb{F}$ , such as  $\mathbb{Q}$ .

**Example.** Let

$$\begin{aligned}
 \mathbb{CP}^n &= \{\text{lines in } \mathbb{C}^{n+1}\} = (\mathbb{C}^{n+1} \setminus \{0\}) / \mathbb{C}^* = S^{2n+1} / S^1 \\
 &= \{[z_0 : \dots : z_n] \mid (z_0, \dots, z_n) \in \mathbb{C}^{n+1} \setminus \{0\}, \forall \lambda \in \mathbb{C}^*, (z_0, \dots, z_n) \sim (\lambda z_0, \dots, \lambda z_n)\} \\
 &= \{z_0 = 0\} \cup \{[1 : z_1 : \dots : z_n] \mid (z_1, \dots, z_n) \in \mathbb{C}^n\} \cong \mathbb{CP}^{n-1} \cup \mathbb{C}^n,
 \end{aligned}$$

where  $S^{2n+1} \subseteq \mathbb{C}^{n+1}$  is the unit sphere and  $S^1 : (z_1, \dots, z_{n+1}) \mapsto (\lambda z_1, \dots, \lambda z_{n+1})$  for  $\lambda \in S^1$ . The attaching map is

$$\begin{aligned}
 S^{2n+1} &\longrightarrow \mathbb{CP}^n \\
 (z_1, \dots, z_{n+1}) &\longmapsto [z_1 : \dots : z_{n+1}].
 \end{aligned}$$

So, inductively in  $n$ ,  $\mathbb{CP}^n$  has a cell structure with one  $2n$ -cell for all  $n$ , so

$$H_{\bullet}(\mathbb{CP}^n) \cong \begin{cases} \mathbb{Z} & \bullet = 0, 2, \dots, 2n-2, 2n \\ 0 & \text{otherwise} \end{cases}.$$

See example sheet 1.

**Remark. Grassmannians**

$$\mathrm{Gr}(k; \mathbb{C}^n) = \{k\text{-dimensional linear subspaces of } \mathbb{C}^n\}$$

also have cell structures with only even-dimensional cells.

**Exercise.**  $\mathbb{RP}^n = S^n \setminus \{\pm 1\} = \mathbb{RP}^{n-1} \cup D^n$  has a cell structure with one cell in each degree  $0 \leq i \leq n$ .

### 3.4 Degrees

How do we compute  $\partial_n^{\mathrm{cell}} : \bigoplus_{\alpha \in I_n} \mathbb{Z} \rightarrow \bigoplus_{\beta \in I_{n-1}} \mathbb{Z}$ ? That is, want values  $d_{\alpha\beta} \in \mathbb{Z}$  such that

$$\partial_n^{\mathrm{cell}}(e_\alpha^n) = \sum_{\beta} d_{\alpha\beta} e_\beta^{n-1}.$$

Note we have an attaching map  $\phi_\alpha^n : \partial(e_\alpha^n) = S^{n-1} \rightarrow X_{n-1}$ .

**Lemma 3.10.**  $d_{\alpha\beta}$  is the degree of

$$f_{\alpha\beta} : S_\alpha^{n-1} \xrightarrow{\phi_\alpha^n} X_{n-1} \rightarrow X_{n-1}/X_{n-2} \cong \bigvee_{I_{n-1}} S^{n-1} \xrightarrow{\text{collapse}} S_\beta^{n-1}.$$

**Remark.** For this degree to be well-defined, not just up to sign, need generators for  $H_{n-1}(S_\alpha^{n-1})$  and  $H_{n-1}(S_\beta^{n-1})$ , such as identifying  $S^{n-1} = \partial D^n \subseteq \mathbb{R}^n$ .

*Proof.* By chasing,

$$\begin{array}{ccc} H_n(D_\alpha^n, \partial D_\alpha^n) & \xrightarrow[\sim]{\text{LES}} H_{n-1}(\partial D_\alpha^n) & \xrightarrow{\deg f_{\alpha\beta}} \widetilde{H}_{n-1}(S_\beta^{n-1}) \\ \downarrow (\phi_\alpha^n)_* & \downarrow (\phi_\alpha^n|_{\partial D_\alpha^n})_* & \uparrow \text{collapse} \\ H_n(X_n, X_{n-1}) & \xrightarrow{\partial_n} H_{n-1}(X_{n-1}) & \widetilde{H}_{n-1}(\bigvee_\gamma S_\gamma^{n-1}) \\ & \searrow \partial_n^{\mathrm{cell}} & \uparrow \mathbb{R} \\ & H_{n-1}(X_{n-1}, X_{n-2}) & \xrightarrow[\text{Excision}]{\sim} \widetilde{H}_{n-1}(X_{n-1}/X_{n-2}) \end{array} \quad , \quad \begin{array}{ccccc} 1 & \longrightarrow & 1 & \longrightarrow & d_{\alpha\beta} \\ \downarrow e_\alpha & & & & \uparrow \\ & \searrow & \sum_\gamma d_{\alpha\gamma} e_\gamma & \longrightarrow & \sum_\gamma d_{\alpha\gamma} e_\gamma \end{array} ,$$

so  $d_{\alpha\beta} = \deg f_{\alpha\beta}$  as claimed.  $\square$

For this to be useful, we need to be able to compute degrees. Let  $f : S^n \rightarrow S^n$ . Assume that there exists  $y \in S^n$  such that  $f^{-1}(y) = \{x_1, \dots, x_m\}$  is finite. Pick neighbourhoods  $U_i \in \mathcal{U}_i$  and  $V \in \mathcal{V}$  homeomorphic to  $\mathbb{R}^n$  such that  $U_i \cap U_j = \emptyset$  if  $i \neq j$  and  $f|_{U_i} : U_i \rightarrow V \subseteq S^n$ .

**Definition.** The **local degree** is

$$\deg_{x_i} f = H_n(U_i, U_i \setminus \{x_i\}) \cong \mathbb{Z} \rightarrow H_n(V, V \setminus \{y\}) \cong \mathbb{Z}.$$

Note that by excision and the long exact sequence,

$$H_n(U_i, U_i \setminus \{x_i\}) \cong H_n(S^n, S^n \setminus \{x_i\}) \cong H_n(S^n) \cong \mathbb{Z}.$$

By fixing this,  $\deg_{x_i} f$  is well-defined.

**Lemma 3.11.** Under the assumption,

$$\deg f = \sum_{i=1}^m \deg_{x_i} f.$$

**Remark.** If  $f : S^n \rightarrow S^n$  is smooth,  $f^{-1}(y)$  is finite if  $y$  is a regular value for  $f$ , and by Sard's theorem, almost all values, in particular a dense set, are regular.

*Proof.*

$$\begin{array}{ccc}
 H_n(S^n) & \xrightarrow{\deg f} & H_n(S^n) \\
 \text{LES} \downarrow \sim & & \sim \downarrow \text{LES} \\
 H_n(S^n, S^n \setminus \{x_1, \dots, x_m\}) & \xrightarrow{f_*} & H_n(S^n, S^n \setminus \{y\}) \\
 \text{Excision} \uparrow \sim & & \sim \uparrow \text{Excision} \\
 H_n(\bigsqcup_i U_i, \bigsqcup_i U_i \setminus \{x_i\}) \cong \bigoplus_{i=1}^m H_n(U_i, U_i \setminus \{x_i\}) & \xrightarrow{(f|_{U_i})} & H_n(V, V \setminus \{y\})
 \end{array}
 \quad , \quad
 \begin{array}{ccc}
 1 & \xrightarrow{\quad} & \deg f \\
 \downarrow & & \downarrow \\
 (1, \dots, 1) & \rightarrow & \sum_i \deg_{x_i} f
 \end{array}$$

which implies the result.  $\square$

**Example.** Let  $p(z) = z^k + a_{k-1}z^{k-1} + \dots + a_0$  be a complex polynomial. Then  $p$  extends to a map  $\hat{p} : \mathbb{C} \cup \{\infty\} = S^2 \rightarrow S^2$  of degree  $\deg \hat{p} = k$ . As in lecture 1, show  $\hat{p} \simeq (q : z \mapsto z^k)$ . Now  $q^{-1}(1) = \{\zeta_1, \dots, \zeta_k\}$  and near each  $\zeta_i$ ,  $q$  is a local homeomorphism. And the different local homeomorphisms differ by rotation, so the local degrees at  $\zeta_i$  are all equal.

**Exercise.** The Klein bottle  $K$  has

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_2^{\text{cell}}(K) & \longrightarrow & C_1^{\text{cell}}(K) & \longrightarrow & C_0^{\text{cell}}(K) \longrightarrow 0 \\
 & & \downarrow \mathbb{R} & & \downarrow \mathbb{R} & & \downarrow \mathbb{R} \\
 & & \mathbb{Z} & \xrightarrow{1 \mapsto (0,2)} & \mathbb{Z} \oplus \mathbb{Z} & \xrightarrow{0} & \mathbb{Z}
 \end{array}$$

so  $H_1(K; \mathbb{Z}) \cong \mathbb{Z} \oplus \mathbb{Z}/2$  and  $H_2(K; \mathbb{Z}) = 0$ , again.

**Example.** The real projective space  $\mathbb{RP}^n = D^n \cup \mathbb{RP}^{n-1} = D^n \cup \dots \cup D^1 \cup \{\text{point}\}$  has one cell of each degree  $0 \leq i \leq n$ , so

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_n^{\text{cell}}(\mathbb{RP}^n) & \longrightarrow & \dots & \longrightarrow & C_0^{\text{cell}}(\mathbb{RP}^n) \longrightarrow 0 \\
 & & \downarrow \mathbb{R} & & & & \downarrow \mathbb{R} \\
 & & \mathbb{Z} & \xrightarrow{\quad} & \dots & \xrightarrow{\quad} & \mathbb{Z}
 \end{array}$$

Then

$$\eta : \partial D^k \xrightarrow{\phi} \mathbb{RP}^{k-1} \rightarrow \mathbb{RP}^{k-1}/\mathbb{RP}^{k-2} \cong S^{k-1}$$

is two-to-one and the local maps differ by the antipodal map, so  $\partial_k^{\text{cell}} = 1 + (-1)^k$ . Thus

$$C_{\bullet}^{\text{cell}}(\mathbb{RP}^n) = \begin{cases} \mathbb{Z} \xrightarrow{2} \mathbb{Z} \xrightarrow{0} \mathbb{Z} \rightarrow \dots \rightarrow \mathbb{Z} \xrightarrow{2} \mathbb{Z} \xrightarrow{0} \mathbb{Z} & n \text{ even} \\ \mathbb{Z} \xrightarrow{0} \mathbb{Z} \xrightarrow{2} \mathbb{Z} \rightarrow \dots \rightarrow \mathbb{Z} \xrightarrow{0} \mathbb{Z} \xrightarrow{2} \mathbb{Z} & n \text{ odd} \end{cases}$$

so

$$H_{\bullet}(\mathbb{RP}^n; \mathbb{Z}) = \begin{cases} \mathbb{Z} & \bullet = 0 \\ \mathbb{Z}/2 & 0 < \bullet < n, \bullet \text{ odd} \\ \mathbb{Z} & \bullet = n, n \text{ odd} \\ 0 & \text{otherwise} \end{cases}$$

**Exercise.** Let  $\Sigma_g$  be a  $4g$ -gon with edge identifications  $a_1 b_1 a_1^{-1} b_1^{-1} \dots a_g b_g a_g^{-1} b_g^{-1}$ . Then

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_2^{\text{cell}}(\Sigma_g) & \longrightarrow & C_1^{\text{cell}}(\Sigma_g) & \longrightarrow & C_0^{\text{cell}}(\Sigma_g) \longrightarrow 0 \\
 & & \downarrow \mathbb{R} & & \downarrow \mathbb{R} & & \downarrow \mathbb{R} \\
 & & \mathbb{Z} & \xrightarrow{\quad} & \mathbb{Z}^{2g} & \xrightarrow{\quad} & \mathbb{Z}
 \end{array}$$

which vanishes as for Klein and as  $H_0(\Sigma_g) = \mathbb{Z}$ .

### 3.5 Cellular cohomology

There is also **cellular cohomology**

$$C_{\text{cell}}^i(X) = H^i(X_i, X_{i-1}), \quad \partial_{\text{cell}}^* : H^i(X_i, X_{i-1}) \rightarrow H^{i+1}(X_{i+1}, X_i),$$

and  $H_{\text{cell}}^{\bullet}(X) \cong H_{\text{cell}}(X)$  by the exactly analogous argument as for homology.

Lecture 10  
Friday  
30/10/20

**Lemma 3.12.**  $C_{\text{cell}}^{\bullet}(X) = \text{Hom}(C_{\bullet}^{\text{cell}}(X), \mathbb{Z})$  and  $\partial_{\text{cell}}^*$  is the adjoint of  $\partial_{\text{cell}}$ .

See question sheet 2 for related results, in particular the existence of the  $h$ -maps.

*Proof.* If  $(\cdot)^{\vee}$  is the linear dual, then

$$\begin{array}{ccccc}
 H^i(X_i, X_{i-1}) & \xrightarrow{i^*} & H^i(X_i) & \xrightarrow{\partial^*} & H^{i+1}(X_{i+1}, X_i) \\
 \sim \downarrow h & & \downarrow h & & \sim \downarrow h \\
 \text{Hom}(H_i(X_i, X_{i-1}), \mathbb{Z}) & \xrightarrow{i_*^{\vee}} & \text{Hom}(H_i(X_i), \mathbb{Z}) & \xrightarrow{\partial^{\vee}} & \text{Hom}(H_{i+1}(X_{i+1}, X_i), \mathbb{Z})
 \end{array}$$

$\partial_{\text{cell}}^*$  (top curved arrow) and  $\partial_{\text{cell}}^{\vee}$  (bottom curved arrow)

A direct check shows the outer  $h$  maps are isomorphisms, so just need the diagram to commute. The left hand square commutes by naturality of  $h$  and the right hand square commutes by naturality of homology and cohomology long exact sequences.  $\square$

**Proposition 3.13.** Let  $X$  be a finite cell complex. Then

$$H^i(X; \mathbb{Z}) \cong (H_i(X; \mathbb{Z}) / \text{Tors}) \oplus \text{Tors } H_{i-1}(X; \mathbb{Z}).$$

Note that for an abelian group  $G$ ,

$$\text{Tors } G = \{\text{elements of finite order}\} = \{\text{all } \mathbb{Z}/k\text{-summands} \mid k \geq 2\} \leq G.$$

*Proof.* This is now just algebra. Let  $C_{\bullet}$  be a chain complex of free finitely generated abelian groups, and  $C^{\bullet} = \text{Hom}(C_{\bullet}, \mathbb{Z})$  the dual complex. Then

$$H^i(C^{\bullet}) = (H_i(C_{\bullet}) / \text{Tors}) \oplus \text{Tors } H_{i-1}(C_{\bullet}).$$

The key idea is to break  $C_{\bullet}$  into a collection of short exact sequences

$$0 \rightarrow Z_n \rightarrow C_n \rightarrow B_{n-1} \rightarrow 0, \quad 0 \rightarrow B_n \rightarrow Z_n \rightarrow H_n(C_{\bullet}) \rightarrow 0,$$

where  $Z_i = \ker(\partial_i : C_i \rightarrow C_{i-1})$  and  $B_i = \text{im}(\partial_{i+1} : C_{i+1} \rightarrow C_i)$ . In the former sequence, all the terms are free finitely generated abelian groups, so we can non-canonically split these and write  $C_n \cong Z_n \oplus B_{n-1}$  for all  $n$ , so there exists  $\alpha_n : B_{n-1} \rightarrow C_n$  such that  $\partial_n \circ \alpha_n = \text{id}$ . Then  $C_{\bullet}$  becomes

$$\cdots \rightarrow Z_{n+1} \oplus B_n \xrightarrow{B_n \rightarrow Z_n} Z_n \oplus B_{n-1} \xrightarrow{B_{n-1} \rightarrow Z_{n-1}} Z_{n-1} \oplus B_{n-2} \rightarrow \cdots, \quad (2)$$

that is  $C_{\bullet}$  breaks into a sum of complexes of the form

$$0 \rightarrow B_n \xrightarrow{A_n} Z_n \rightarrow 0.$$

By the Smith normal form, there exists a  $\mathbb{Z}$ -linear change of basis such that  $A_n$  has matrix

$$\begin{pmatrix} d_1 & & & & \\ & \ddots & & & \\ & & d_k & & \\ & & & 0 & \\ & & & & \ddots \\ & & & & & 0 \end{pmatrix}, \quad \begin{cases} d_1 \mid d_2 \\ \vdots \\ d_{k-1} \mid d_k \end{cases},$$

so (2) breaks into a direct sum of complexes of the form

$$0 \rightarrow \mathbb{Z} \xrightarrow{0} \mathbb{Z} \rightarrow 0, \quad 0 \rightarrow \mathbb{Z} \xrightarrow{d_i} \mathbb{Z} \rightarrow 0.$$

Now the relation  $H^i(C^{\bullet}) = (H_i(C_{\bullet}) / \text{Tors}) \oplus \text{Tors } H_{i-1}(C_{\bullet})$  is obvious, by just checking in these two examples.  $\square$

**Remark.** For abelian groups  $H$  and  $G$  set

$$\text{Ext}^1(H, G) = \{\text{short exact sequences } 0 \rightarrow G \rightarrow J \rightarrow H \rightarrow 0\} / \sim,$$

where two are equivalent if the obvious thing happens. That is, there exists  $\phi$  making the diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & G & \longrightarrow & J_1 & \longrightarrow & H \longrightarrow 0 \\ & & \parallel & & \downarrow \exists \phi & & \parallel \\ 0 & \longrightarrow & G & \longrightarrow & J_2 & \longrightarrow & H \longrightarrow 0 \end{array}$$

commute, so  $\phi$  is an isomorphism by the 5-lemma. The **universal coefficient theorem** says there are split exact sequences

$$0 \rightarrow \text{Ext}^1(H_{n-1}(X; \mathbb{Z}), G) \rightarrow H^n(X; G) \rightarrow \text{Hom}(H_n(X; \mathbb{Z}), G) \rightarrow 0.$$

We will not prove this.

### 3.6 Euler characteristic

Recall that if  $X$  is a finite cell complex, we saw  $H_i(X)$  is finitely generated for all  $i$ , and  $H_i(X) = 0$  if  $i > \dim X$ . So  $\bigoplus_i H_i(X; \mathbb{Q})$  is a finite-dimensional graded  $\mathbb{Q}$ -vector space.

**Definition.** The **Euler characteristic** of a finite cell complex  $X$  is

$$\chi(X) = \sum_{k \geq 0} (-1)^k \text{rk}_{\mathbb{Z}} H_k(X; \mathbb{Z}).$$

**Lemma 3.14.**

$$\chi(X) = \sum_{k \geq 0} (-1)^k N_k,$$

where  $N_k$  is the number of  $k$ -cells in  $X$ .

*Proof.*  $N_k$  is the rank of  $C_k^{\text{cell}}(X)$  and we just observed we have short exact sequences

$$0 \rightarrow B_k \rightarrow Z_k \rightarrow H_k(X) \rightarrow 0, \quad 0 \rightarrow Z_k \rightarrow C_k \rightarrow B_{k-1} \rightarrow 0.$$

Then

$$\text{rk } H_k(X) = \text{rk } Z_k - \text{rk } B_k = z_k - b_k, \quad \text{rk } C_k = \text{rk } Z_k + \text{rk } B_{k-1} = z_k + b_{k-1},$$

so

$$\begin{aligned} \sum_{k \geq 0} (-1)^k \text{rk}_{\mathbb{Z}} H_k(X; \mathbb{Z}) &= \sum_{k \geq 0} (-1)^k (z_k - b_k) = \sum_{k \geq 1} (-1)^k (z_k - (N_{k+1} - z_{k+1})) \\ &= \sum_{k \geq 0} (-1)^k N_{k+1} + z_0 = \sum_{k \geq 0} (-1)^k N_k. \end{aligned}$$

□

**Remark.** If  $\mathbb{F}$  is a field,

$$\chi(X) = \sum_{k \geq 0} (-1)^k \dim_{\mathbb{F}} H_k(X; \mathbb{F}).$$

**Example.**

- $S^4 \not\cong \mathbb{CP}^2$  as  $\chi(S^4) = 2$  and  $\chi(\mathbb{CP}^2) = 3$ , since  $S^4$  has one 0-cell and one 4-cell and  $\mathbb{CP}^2$  has one 4-cell, one 2-cell, and one 0-cell.
- $\chi(\Sigma_g) = 2 - 2g$ , since  $\Sigma_g$  has one 0-cell,  $2g$  distinct 1-cells, and one 2-cell.
- If  $A$  and  $B$  are finite cell complexes,  $A \times B$  has a product cell structure such that the open  $i$ -cells are of the form  $(j\text{-cell in } A) \times ((i-j)\text{-cell in } B)$ , so  $\chi(A \times B) = \chi(A) \chi(B)$ .
- If  $X = A \cup B$  is a union of two subcomplexes,  $\chi(X) = \chi(A) + \chi(B) - \chi(A \cap B)$ .