

Algebraic Geometry

Lectured by Prof Mark Gross
Typed by David Kurniadi Angdinata

Michaelmas 2020

Syllabus

Contents

0	Brief review of classical algebraic geometry and motivation for scheme theory	3
0.1	Classical algebraic geometry	3
0.2	Why schemes?	3
0.3	Categorical philosophy	4
0.4	Solutions over non-algebraically closed fields	4
0.5	Spectrum of a ring	5
1	Sheaves	6
1.1	Sheaves	6
1.2	Examples	7
1.3	Stalks	7
1.4	Sheafification	8
1.5	Kernels, cokernels, and images	9
1.6	Passing between spaces	10
2	Schemes	11
2.1	Localisation of a ring	11
2.2	Construction of the structure sheaf	11
2.3	Ringed spaces	12
2.4	Affine schemes	13
2.5	Projective schemes	16
2.6	Open and closed subschemes	18
2.7	Fibre products	19

0 Brief review of classical algebraic geometry and motivation for scheme theory

The following are the main references for the course.

- R Hartshorne, Algebraic geometry, 1977
- U Goertz and T Wedhorn, Algebraic geometry I, 2010
- R Vakil, The rising sea: foundations of algebraic geometry, 2017

Lecture 1
Friday
09/10/20

0.1 Classical algebraic geometry

Throughout this discussion, we take the base field k to be algebraically closed. An **affine variety** $V \subseteq \mathbb{A}^n(k)$, where, once one has chosen coordinates, $\mathbb{A}^n(k) = k^n$, is given by the vanishing of polynomials $f_1, \dots, f_r \in k[X_1, \dots, X_n]$. If $I = \langle f_1, \dots, f_r \rangle \triangleleft k[X_1, \dots, X_n]$ is any ideal, we set

$$\mathbb{V}(I) = \{z \in \mathbb{A}^n \mid \forall f \in I, f(z) = 0\}.$$

First set $\mathbb{P}^n(k) = (k^{n+1} \setminus \{0\})/k^*$ with **homogeneous coordinates** $(x_0 : \dots : x_n)$. A **projective variety** $V \subseteq \mathbb{P}^n$ is given by the vanishing of homogeneous polynomials $F_1, \dots, F_r \in k[X_0, \dots, X_n]$. If I is the ideal generated by the homogeneous ideals F_i , that is if $F \in I$ then so are all its homogeneous parts, we set

$$\mathbb{V}(I) = \{z \in \mathbb{P}^n \mid \forall F \in I \text{ homogeneous}, F(z) = 0\}.$$

If $V = \mathbb{V}(I) \subseteq \mathbb{A}^n$, set

$$\mathbb{I}(V) = \{f \in k[X_1, \dots, X_n] \mid \forall x \in V, f(x) = 0\}.$$

Observe that $\mathbb{V}(\mathbb{I}(V)) = V$, by tautology, and $\mathbb{I}(\mathbb{V}(I)) \supseteq \sqrt{I}$, which is obvious. Recall that the **radical** \sqrt{I} of the ideal I is defined by $f \in \sqrt{I}$ if and only if there exists $m > 0$ such that $f^m \in I$. **Hilbert's Nullstellensatz** states that, noting $k = \bar{k}$, $\mathbb{I}(\mathbb{V}(I)) = \sqrt{I}$. The **coordinate ring** is

$$k[V] = k[X_1, \dots, X_n] / \mathbb{I}(V).$$

This may be regarded as the ring of polynomial functions on V , and it is a finitely generated reduced k -algebra. Recall that a **k -algebra** is a commutative ring containing k as a subring. It is **finitely generated** if it is the quotient of a polynomial ring over k , and **reduced** if $a^m = 0$ implies that $a = 0$.

0.2 Why schemes?

A better question is what is wrong with varieties?

- With varieties, always work over algebraically closed fields. For example, let $I = \langle X^2 + Y^2 + 1 \rangle \subseteq \mathbb{R}[X, Y]$. Then $\mathbb{V}(I) = \emptyset$, but I is a prime ideal, hence radical, so $\mathbb{I}(\mathbb{V}(I)) = \mathbb{R}[X, Y] \neq I$.
- Number theory? Diophantine equations. If $I \subseteq \mathbb{Z}[X_1, \dots, X_n]$ is an ideal, have $\mathbb{V}(I) \subseteq \mathbb{Z}^n$. For example, $X^n + Y^n = Z^n$.
- Why should we only consider radical, or prime, ideals? For example, a natural situation is

$$X_1 = \mathbb{V}(X - Y^2) \subseteq \mathbb{A}^2, \quad X_2 = \mathbb{V}(X) \subseteq \mathbb{A}^2.$$

Then $X_1 \cap X_2 = \mathbb{V}(X - Y^2, X)$. Note $I = \langle X - Y^2, X \rangle = \langle X, Y^2 \rangle$ is not a radical ideal, because $Y \notin I$ and $Y^2 \in I$ so $Y \notin \sqrt{I}$. Recall the coordinate ring of X_i is $k[X_i] = k[X, Y]/I_i$. Then $k[X_1 \cap X_2] = k[X, Y]/\langle X, Y^2 \rangle \cong k[Y]/\langle Y^2 \rangle$. So thinking of the coordinate ring of $X_1 \cap X_2$ as functions on $X_1 \cap X_2$, we have a function Y whose square is zero, but is not itself zero.

0.3 Categorical philosophy

What is a point? In the category of sets, objects are sets, and if A and B are sets, then morphisms are $\text{Hom}(A, B)$, the set of maps $f : A \rightarrow B$. Let $*$ be a one-element set. Then the elements of any set X are in one-to-one correspondence with $\text{Hom}(*, X)$. In the category of affine varieties, objects are affine varieties and morphisms are $\text{Hom}(X, Y) = \text{Hom}_{k\text{-alg}}(k[Y], k[X])$. In this category, a point is a single point with coordinate ring k . Giving a morphism

$$\{\text{point}\} \rightarrow X = \mathbb{V}(I) \subseteq \mathbb{A}^n, \quad I \subseteq k[X_1, \dots, X_n],$$

for I a radical ideal, is the same as giving a homomorphism

$$\begin{aligned} \phi : k[X] = k[X_1, \dots, X_n]/I &\longrightarrow k \\ X_i &\longmapsto a_i \end{aligned}$$

Note that ϕ vanishes in I if and only if $f(a_1, \dots, a_n) = 0$ for all $f \in I$, which is if and only if $(a_1, \dots, a_n) \in \mathbb{V}(I) = X$. Note ϕ is surjective, and hence $\ker \phi$ is a maximal ideal. With k algebraically closed, the maximal ideals at $k[X]$ are all of the form $\langle X_1 - a_1, \dots, X_n - a_n \rangle$ for $(a_1, \dots, a_n) \in X$, a consequence of Hilbert's Nullstellensatz. That is, there exists one-to-one correspondences

$$\{\text{points of } X\} \longleftrightarrow \{k\text{-algebra homomorphisms } \phi : k[X] \rightarrow k\} \longleftrightarrow \{\text{maximal ideals of } k[X]\}.$$

0.4 Solutions over non-algebraically closed fields

What if k is not algebraically closed? We may want to consider solutions not just in $k^n = \mathbb{A}^n$ but $(k')^n$ for k' any field extension of k . That is, we may consider k -algebra homomorphisms

$$\begin{aligned} \phi : k[X] = k[X_1, \dots, X_r]/I &\longrightarrow k' \\ X_i &\longmapsto a_i \end{aligned}$$

This gives a tuple $(a_1, \dots, a_n) \in (k')^n$ with $f(a_1, \dots, a_n) = 0$ for all $f \in I$. Then ϕ need not be surjective, so can only say the image of ϕ is a subring of a field, hence an integral domain. Thus $\ker \phi$ is a prime ideal, and maximal if and only if $\text{im } \phi$ is a field.

Example. The \mathbb{R} -algebra homomorphism

$$\begin{aligned} \phi : \mathbb{R}[X, Y] / \langle X^2 + Y^2 + 1 \rangle &\longrightarrow \mathbb{C} \\ X &\longmapsto 0 \\ Y &\longmapsto i \end{aligned}$$

is surjective with kernel $\langle X, Y^2 + 1 \rangle$, since $\mathbb{R}[Y] / \langle Y^2 + 1 \rangle \cong \mathbb{C}$. This is a maximal ideal but is not of the form $\langle X - a, Y - b \rangle$ for $(a, b) \in \mathbb{R}^2$. If instead we considered the map

$$\begin{aligned} \mathbb{R}[X, Y] / \langle X^2 + Y^2 + 1 \rangle &\longrightarrow \mathbb{C} \\ X &\longmapsto 0 \\ Y &\longmapsto -i \end{aligned},$$

we get the same kernel. That is, $(0, i)$ and $(0, -i)$ are solutions to $X^2 + Y^2 + 1 = 0$, but they correspond to the same maximal ideal. In fact, this maximal ideal corresponds to a Galois orbit of $\text{Gal}(\mathbb{C}/\mathbb{R})$ of solutions.

There are more exotic points by taking even bigger fields.

Example. Let $k(X)$ be the field of fractions of $k[X] = \mathbb{R}[X, Y] / \langle X^2 + Y^2 + 1 \rangle$. There is an inclusion

$$\begin{aligned} k[X] &\longrightarrow k(X) \\ f &\longmapsto \frac{f}{1} \\ (X, Y) &\longmapsto (X, Y) \end{aligned}.$$

The kernel of this map is zero. This gives a solution to the equation $X^2 + Y^2 + 1 = 0$ with coordinates in the field $k(X)$. This solution is $(X, Y) \in \mathbb{A}^2(k(X))$.

The moral is that once we start looking at solutions to equation over any field, then we get maps $k[X] \rightarrow k'$ with kernel not necessarily maximal. What about solutions over rings?

Example. Let $A = \mathbb{Z}[X_1, \dots, X_n]/I$, and let R be any commutative ring. We define an R -valued point of $\text{Spec } A$ to be a ring homomorphism

$$\begin{array}{ccc} A & \longrightarrow & R \\ X_i & \longmapsto & r_i \end{array}.$$

Then $f(r_1, \dots, r_n) = 0$ for all $f \in I$. This gives a lot of flexibility. For example,

- $R = \mathbb{Z}$ gives diophantine equations,
- $R = \mathbb{F}_p$ gives solutions modulo p , and
- $R = \mathbb{Q}$ gives rational solutions.

Take this to its logical conclusion. Let A be a ring, where all rings are commutative in this course. Given A , we hope for some geometric object $\text{Spec } A$, the **spectrum** of A . For a ring R , the set of **R -valued points** of X is

$$X(R) = \text{Hom}_{\text{ring}}(A, R).$$

A morphism $X = \text{Spec } A \rightarrow Y = \text{Spec } B$ should be the same thing as giving a morphism $\phi: B \rightarrow A$. Define the category of **affine schemes** to be the opposite category to the category of rings. Define a **scheme** to be something which is locally isomorphic to an affine scheme. By analogy, a **manifold** is a topological space with an open cover $\{U_i\}$ with each U_i homeomorphic to an open subset of \mathbb{R}^n . To make sense of the definition of schemes, we need a lot of language.

0.5 Spectrum of a ring

Definition. Let A be a ring. Then

$$\text{Spec } A = \{\mathfrak{p} \subseteq A \mid \mathfrak{p} \text{ a prime ideal}\}.$$

For $I \subseteq A$ an ideal, define

$$\mathbb{V}(I) = \{\mathfrak{p} \subseteq A \mid \mathfrak{p} \text{ prime, } \mathfrak{p} \supseteq I\}.$$

Proposition 0.1. *The sets $\mathbb{V}(I)$ form the closed sets of a topology on $\text{Spec } A$, called the **Zariski topology**.*

Proof.

- $\mathbb{V}(A) = \emptyset$.
- $\mathbb{V}(0) = \text{Spec } A$.
- If $\{I_i\}_{i \in J}$ is a collection of ideals, then

$$\mathbb{V}\left(\sum_{i \in J} I_i\right) = \bigcap_{i \in J} \mathbb{V}(I_i).$$

- Claim that

$$\mathbb{V}(I_1 \cap I_2) = \mathbb{V}(I_1) \cup \mathbb{V}(I_2).$$

\supseteq Obvious.

\subseteq If $\mathfrak{p} \supseteq I_1 \cap I_2$ is prime, then $\mathfrak{p} \supseteq I_1$ or $\mathfrak{p} \supseteq I_2$. See Atiyah-Macdonald, Proposition 1.11.ii. ¹

□

Example. Let $A = k[X_1, \dots, X_n]$ with k algebraically closed and $I \subseteq A$ an ideal. Then the maximal ideals \mathfrak{m} of A containing I are in one-to-one correspondence with the zero set of I in $\mathbb{A}^n(k)$, so

$$\left\{ \langle X_1 - a_1, \dots, X_n - a_n \rangle \supseteq I, a_i \in k \right\} \quad \rightsquigarrow \quad \left\{ (a_1, \dots, a_n) \in \mathbb{V}(I) \subseteq \mathbb{A}^n(k) \right\}.$$

The new $\mathbb{V}(I)$ now extends this notion of zero set by including possible other prime ideals.

Example. If k is a field, $\text{Spec } k = \{0\}$, so the topological space cannot see the field.

We fix this by also thinking about what functions are on these spaces.

¹Exercise: try to prove without looking up

1 Sheaves

Fix a topological space X .

1.1 Sheaves

Definition. A **presheaf** \mathcal{F} on X consists of the following data.

- For every open set $U \subseteq X$ an abelian group $\mathcal{F}(U)$.
- Whenever given an inclusion $V \subseteq U \subseteq X$, a **restriction map** $\rho_{UV} : \mathcal{F}(U) \rightarrow \mathcal{F}(V)$, a homomorphism, such that
 - $\rho_{UU} = \text{id}_{\mathcal{F}(U)}$, and
 - if $W \subseteq V \subseteq U$, then $\rho_{UW} = \rho_{VW} \circ \rho_{UV}$.

Remark. Can think of a presheaf as a contravariant functor from the category of open sets of X , the category whose objects are open subsets of X and whose morphisms are inclusions of open sets, to the category of abelian groups. Can replace the category of abelian groups with any desired category, such as commutative rings.

Definition. A **morphism of presheaves** $f : \mathcal{F} \rightarrow \mathcal{G}$ is a collection of homomorphisms $f_U : \mathcal{F}(U) \rightarrow \mathcal{G}(U)$ such that for all $V \subseteq U$ the diagram

$$\begin{array}{ccc} \mathcal{F}(U) & \xrightarrow{f_U} & \mathcal{G}(U) \\ \rho_{UV} \downarrow & & \downarrow \rho_{UV} \\ \mathcal{F}(V) & \xrightarrow{f_V} & \mathcal{G}(V) \end{array}$$

is commutative.

Definition. A presheaf \mathcal{F} is a **sheaf** if it satisfies the following additional axioms.

- S1. If $U \subseteq X$ is covered by an open cover $\{U_i\}$ and $s \in \mathcal{F}(U)$ satisfies $s|_{U_i} = \rho_{UU_i}(s) = 0$ for all i , then $s = 0$.
- S2. If U and $\{U_i\}$ are as in S1 and $s_i \in \mathcal{F}(U_i)$ such that $s_i|_{U_i \cap U_j} = s_j|_{U_i \cap U_j}$ for all i and j , then there exists $s \in \mathcal{F}(U)$ with $s|_{U_i} = s_i$ for all i .

Remark.

- If \mathcal{F} is a sheaf, then $\emptyset \subseteq X$ is covered by the empty covering, and hence $\mathcal{F}(\emptyset) = 0$.
- S1 and S2 together can be described as saying, given U and $\{U_i\}_{i \in I}$,

$$0 \rightarrow \mathcal{F}(U) \xrightarrow{\alpha} \prod_{i \in I} \mathcal{F}(U_i) \xrightleftharpoons[\beta_2]{\beta_1} \prod_{i,j} \mathcal{F}(U_i \cap U_j)$$

is exact, where

$$\alpha(s) = (s|_{U_i})_{i \in I}, \quad \beta_1((s_i)_{i \in I}) = (s_i|_{U_i \cap U_j})_{i,j}, \quad \beta_2((s_i)_{i \in I}) = (s_j|_{U_i \cap U_j})_{i,j}.$$

Exactness means

- α is injective, which is S1,
- $\beta_1 \circ \alpha = \beta_2 \circ \alpha$, and
- for any $(s_i) \in \prod_{i \in I} \mathcal{F}(U_i)$, with $\beta_1((s_i)) = \beta_2((s_i))$, there exists $s \in \mathcal{F}(U)$ with $\alpha(s) = (s_i)$, which is S2.

1.2 Examples

Example.

- Let X be any topological space, and let

$$\mathcal{F}(U) = \{\text{continuous functions } U \rightarrow \mathbb{R}\}.$$

This is a sheaf, by

$$\begin{aligned} \rho_{UV} : \mathcal{F}(U) &\longrightarrow \mathcal{F}(V) \\ f &\longmapsto f|_V \end{aligned}.$$

S1. A continuous function is zero if it is zero on every open set of a cover.

S2. Continuous functions can be glued.

- Let $X = \mathbb{C}$ with the Euclidean topology, and let

$$\mathcal{F}(U) = \{f : U \rightarrow \mathbb{C} \mid f \text{ is a bounded analytic function}\}.$$

This is a presheaf. It satisfies S1, and does not satisfy S2. For example, consider the cover $\{U_i\}_{i \in \{1,2,\dots\}}$ of \mathbb{C} given by $U_i = \{z \in \mathbb{C} \mid |z| < i\}$ and

$$\begin{aligned} s_i : U_i &\longrightarrow \mathbb{C} \\ z &\longmapsto z \end{aligned}.$$

Note if $i < j$, then $U_i \cap U_j = U_i$ and $s_i|_{U_i \cap U_j} = s_j|_{U_i \cap U_j}$. But if we glue we get the function $z : \mathbb{C} \rightarrow \mathbb{C}$, which is not bounded. Note $\mathcal{F}(\mathbb{C}) = \mathbb{C}$.

- Take any group G and set $\mathcal{F}(U) = G$ for any open set U . This is called the **constant presheaf**. This is not a sheaf. Let $U = U_1 \sqcup U_2$. If we wanted a sheaf,

$$\begin{array}{ccc} \mathcal{F}(U_1) = G & & \mathcal{F}(U_2) = G \\ & \searrow \quad \swarrow & \\ & \mathcal{F}(U_1 \cap U_2) = \mathcal{F}(\emptyset) = 0 & \end{array},$$

so if S2 is satisfied, would want $s_1 \in \mathcal{F}(U_1)$ and $s_2 \in \mathcal{F}(U_2)$ to glue. We would then want to have $\mathcal{F}(U) = G \times G$. Now give G the discrete topology, and define instead

$$\mathcal{F}(U) = \{f : U \rightarrow G \text{ continuous}\},$$

that is f is locally constant. That is, if $x \in U$, there exists a neighbourhood $x \in V \subseteq U$ with $f|_V$ constant. This is called the **constant sheaf** and if U is non-empty and connected, then $\mathcal{F}(U) = G$.

- If X is an algebraic variety, and $U \subseteq X$ is a Zariski open subset, define

$$\mathcal{O}_X(U) = \{f : U \rightarrow k \mid f \text{ regular function}\}.$$

Roughly f is **regular** means that every point of U has an open neighbourhood on which f is expressed as a ratio of polynomials g/h with h non-vanishing on the neighbourhood. Then \mathcal{O}_X is a sheaf, called the **structure sheaf** of X .

1.3 Stalks

Definition. Let \mathcal{F} be a presheaf on X . Let $p \in X$. Then the **stalk** of \mathcal{F} at p is

$$\mathcal{F}_p = \{(U, s) \mid U \subseteq X \text{ is an open neighbourhood of } p, s \in \mathcal{F}(U)\} / \equiv,$$

where $(U, s) \equiv (V, s')$ if there exists $W \subseteq U \cap V$ also a neighbourhood of p such that $s|_W = s'|_W$. An equivalence class of a pair (U, s) is called a **germ**.

Remark. $\mathcal{F}_p = \varinjlim_{p \in U} \mathcal{F}(U)$.

Note that a morphism $f : \mathcal{F} \rightarrow \mathcal{G}$ of presheaves induces a morphism

$$f_p : \mathcal{F}_p \longrightarrow \mathcal{G}_p \\ (U, s) \longmapsto (U, f_U(s)) .$$

Proposition 1.1. *Let $f : \mathcal{F} \rightarrow \mathcal{G}$ be a morphism of sheaves. Then f is an isomorphism if and only if f_p is an isomorphism for all $p \in X$.*

Proof.

\implies Obvious.

\impliedby Assume f_p is an isomorphism for all $p \in X$. Need to show that $f_U : \mathcal{F}(U) \rightarrow \mathcal{G}(U)$ is an isomorphism for all $U \subseteq X$, as then we can define $(f^{-1})_U = (f_U)^{-1}$. Check that with this definition, $(f^{-1})_U$ is compatible with restriction maps, hence f^{-1} is a morphism of sheaves.²

- f_U is injective. Suppose $s \in \mathcal{F}(U)$, and $f_U(s) = 0$. Then for all $p \in U$, $f_p(U, s) = (U, f_U(s)) = (U, 0) = 0 \in \mathcal{G}_p$. Since f_p is injective, $(U, s) = 0$ in \mathcal{F}_p . That is, there exists an open neighbourhood V_p of p in U such that $s|_{V_p} = 0$. Since $\{V_p\}_{p \in U}$ cover U , we see by S1 that $s = 0$.
- f_U is surjective. Let $t \in \mathcal{G}(U)$ and write $t_p = (U, t) \in \mathcal{G}_p$. Since f_p is surjective, there exists $s_p \in \mathcal{F}_p$ with $f_p(s_p) = t_p$. That is, there exists $V_p \subseteq U$ an open neighbourhood of p , and a germ (V_p, s_p) such that $(V_p, f_{V_p}(s_p)) \equiv (U, t)$. By shrinking V_p if necessary, we can assume that $t|_{V_p} = f_{V_p}(s_p)$. Now on $V_p \cap V_q$,

$$f_{V_p \cap V_q}(s_p|_{V_p \cap V_q} - s_q|_{V_p \cap V_q}) = t|_{V_p \cap V_q} - t|_{V_p \cap V_q} = 0,$$

and hence by injectivity of $f_{V_p \cap V_q}$ already proved, we have $s_p|_{V_p \cap V_q} = s_q|_{V_p \cap V_q}$. By S2 the s_p 's glue to give an element $s \in \mathcal{F}(U)$ with $s|_{V_p} = s_p$, for all $p \in U$. Now

$$f_U(s)|_{V_p} = f_{V_p}(s|_{V_p}) = f_{V_p}(s_p) = t|_{V_p} .$$

By S1, applied to $f_U(s) - t$, we get $f_U(s) = t$. Thus f_U is surjective.

□

1.4 Sheafification

Theorem 1.2 (Sheafification). *Given a presheaf \mathcal{F} , there exists a sheaf \mathcal{F}^+ and a morphism $\theta : \mathcal{F} \rightarrow \mathcal{F}^+$ satisfying the following universal property. For any sheaf \mathcal{G} and morphism $\phi : \mathcal{F} \rightarrow \mathcal{G}$, there exists a unique morphism $\phi^+ : \mathcal{F}^+ \rightarrow \mathcal{G}$ such that $\phi^+ \circ \theta = \phi$, so*

$$\begin{array}{ccc} \mathcal{F} & \xrightarrow{\theta} & \mathcal{F}^+ \\ & \searrow \phi & \downarrow \phi^+ \\ & & \mathcal{G} \end{array}$$

The pair (\mathcal{F}^+, θ) is unique up to unique isomorphism, and is called the **sheafification** of \mathcal{F} .

Proof. See example sheet 1. The idea is to make \mathcal{F}^+ look like functions. Define

$$\mathcal{F}^+(U) = \left\{ s : U \rightarrow \bigsqcup_{p \in U} \mathcal{F}_p \mid \begin{array}{l} \forall p \in U, s(p) \in \mathcal{F}_p, \\ \forall p \in U, \exists p \in V \subseteq U, \exists t \in \mathcal{F}(V), \forall q \in V, s(q) = (V, t) \in \mathcal{F}_q \end{array} \right\} .$$

Then

$$\begin{array}{ccc} \theta_U : \mathcal{F}(U) & \longrightarrow & \mathcal{F}^+(U) \\ s & \longmapsto & (p \mapsto (U, s) \in \mathcal{F}_p) . \end{array}$$

□

Exercise. A recommendation is to do all exercises in chapter II.1 of Hartshorne.

²Exercise

1.5 Kernels, cokernels, and images

Lecture 4
Friday
16/10/20

Definition. Let $f : \mathcal{F} \rightarrow \mathcal{G}$ be a morphism of presheaves on a space X . We define the following.

- The **presheaf kernel** of f , $\ker f$, is the presheaf given by $(\ker f)(U) = \ker(f_U : \mathcal{F}(U) \rightarrow \mathcal{G}(U))$.
- The **presheaf cokernel** $\operatorname{coker} f$ is the presheaf given by $(\operatorname{coker} f)(U) = \operatorname{coker}(f_U) = \mathcal{G}(U) / \operatorname{im} f_U$.
- The **presheaf image** $\operatorname{im} f$ is the presheaf given by $(\operatorname{im} f)(U) = \operatorname{im} f_U$.

Exercise. Check that these are presheaves, that is restrictions work.

Remark 1.3. If $f : \mathcal{F} \rightarrow \mathcal{G}$ is a morphism of sheaves, then $\ker f$ is also a sheaf.

Proof. S1 is certainly satisfied. If $s \in (\ker f)(U) \subseteq \mathcal{F}(U)$ satisfies $s|_{U_i} = 0$ for all U_i in a cover of U so $s = 0$ by S1 for \mathcal{F} . Given $s_i \in (\ker f)(U_i)$ with $\{U_i\}$ an open cover of U , and with $s_i|_{U_i \cap U_j} = s_j|_{U_i \cap U_j}$, then there exists $s \in \mathcal{F}(U)$ with $s|_{U_i} = s_i$ by S2 for \mathcal{F} . But $f_U(s) = 0$ since $f_U(s)|_{U_i} = f_{U_i}(s|_{U_i}) = f_{U_i}(s_i) = 0$ so by S1, $f_U(s) = 0$. \square

Example. Let $X = \mathbb{P}^1$, or think of the Riemann sphere. Let $P, Q \in X$ be distinct points. Let \mathcal{G} be the sheaf of regular functions on X , or think of the sheaf of holomorphic functions. Let \mathcal{F} be the sheaf of regular functions on X which vanish at P and Q . Note $\mathcal{F}(U) = \mathcal{G}(U)$ if $U \cap \{P, Q\} = \emptyset$. Let $U = \mathbb{P}^1 \setminus \{P\}$ and $V = \mathbb{P}^1 \setminus \{Q\}$. Note $\mathcal{F}(\mathbb{P}^1) = 0$ and $\mathcal{G}(\mathbb{P}^1) = k$, because regular functions on \mathbb{P}^1 are constants. Let $f : \mathcal{F} \rightarrow \mathcal{G}$ be the obvious inclusion. Then

$$\begin{aligned} (\operatorname{coker} f)(\mathbb{P}^1) &= k, & (\operatorname{coker} f)(U) &= \mathcal{G}(U) / \mathcal{F}(U) = k[X] / \langle X \rangle = k, \\ (\operatorname{coker} f)(V) &= k, & (\operatorname{coker} f)(U \cap V) &= \mathcal{G}(U \cap V) / \mathcal{F}(U \cap V) = 0. \end{aligned}$$

If S2 holds, then we would need to have $(\operatorname{coker} f)(\mathbb{P}^1) = k \oplus k$. This is not a bug, but a feature.

Definition. Let $f : \mathcal{F} \rightarrow \mathcal{G}$ be a morphism of sheaves.

- The **sheaf kernel** $\ker f$ of f is just the presheaf kernel.
- The **sheaf cokernel** is the sheaf associated to the presheaf cokernel of f .
- The **sheaf image** is the sheaf associated to the presheaf image of f .

\mathcal{F} is a **subsheaf** of \mathcal{G} if we have inclusions $\mathcal{F}(U) \subseteq \mathcal{G}(U)$ for all U compatible with restrictions.

Exercise. The sheaf image $\operatorname{im} f$ is a subsheaf of \mathcal{G} .

We say f is **injective** if $\ker f = 0$. We say f is **surjective** if $\operatorname{im} f = \mathcal{G}$. We say a sequence of morphisms of sheaves

$$\dots \rightarrow \mathcal{F}^{i-1} \xrightarrow{f^i} \mathcal{F}^i \xrightarrow{f^{i+1}} \mathcal{F}^{i+1} \rightarrow \dots$$

is **exact** if $\ker f^{i+1} = \operatorname{im} f^i$ for all i . If $\mathcal{F}' \subseteq \mathcal{F}$ is a subsheaf, we write \mathcal{F}/\mathcal{F}' for the sheaf associated to the presheaf $U \mapsto \mathcal{F}(U) / \mathcal{F}'(U)$. That is, this is the cokernel of the inclusion $\mathcal{F}' \hookrightarrow \mathcal{F}$. A warning is if $f : \mathcal{F} \rightarrow \mathcal{G}$ is surjective, we do not necessarily have $\mathcal{F}(U) \rightarrow \mathcal{G}(U)$ surjective for all U .

Lemma 1.4. Let $f : \mathcal{F} \rightarrow \mathcal{G}$ be a morphism of sheaves. Then for all $p \in X$,

$$(\ker f)_p = \ker(f_p : \mathcal{F}_p \rightarrow \mathcal{G}_p), \quad (\operatorname{im} f)_p = \operatorname{im} f_p.$$

Proof. Have a map

$$\begin{array}{ccc} (\ker f)_p & \longrightarrow & \ker f_p \subseteq \mathcal{F}_p \\ (U, s) & \longmapsto & (U, s) \end{array}.$$

If $s \in (\ker f)(U) = \ker f_U$ represents a germ $(U, s) \in (\ker f)_p$, then $(U, s) \in \mathcal{F}_p$, and $f_p(U, s) = (U, f_U(s)) = (U, 0) = 0 \in \mathcal{G}_p$. So $(U, s) \in \ker f_p$.

- **Injective.** If $(U, s) = 0$ in \mathcal{F}_p , there exists a neighbourhood $V \subseteq U$ of p such that $s|_V = 0$. Then $(U, s) \sim (V, s|_V) = (V, 0) = 0$ in $(\ker f)_p$.
- **Surjective.** If $(U, s) \in \ker f_p$, then $(U, f_U(s)) = 0$ in \mathcal{G}_p . That is, there exists a neighbourhood $V \subseteq U$ of p such that $0 = f_U(s)|_V = f_V(s|_V)$. Thus $s|_V \in (\ker f)(V)$, and $(V, s|_V) \in (\ker f)_p$, and $(V, s|_V)$ maps to the same element in $\ker f_p$ represented by (U, s) .

Let $\text{im}' f$ be the presheaf image. An easy fact is if \mathcal{F} is a presheaf with associated sheaf \mathcal{F}^+ , then $\mathcal{F}_p \cong \mathcal{F}_p^+$ for all $p \in X$.³ Thus $(\text{im } f)_p = (\text{im}' f)_p$, so need to show $(\text{im}' f)_p \cong \text{im } f_p$. Define a map by

$$\begin{aligned} (\text{im}' f)_p &\longrightarrow \text{im } f_p \\ (U, s) &\longmapsto (U, s) \end{aligned}.$$

- **Injective.** If $(U, s) = 0$ in \mathcal{G}_p then there exists a neighbourhood $V \subseteq U$ of p such that $s|_V = 0$. Then $(U, s) \sim (V, 0)$ in $(\text{im}' f)_p$.
- **Surjective.** If $(U, s) \in \text{im } f_p$, then there exists $(V, t) \in \mathcal{F}_p$ with $(U, s) = f_p(V, t) = (V, f_V(t))$, so after shrinking U and V if necessary, then we can take $U = V$ and $f_U(t) = s$. Then $(U, s) \in (\text{im}' f)_p$.

□

Proposition 1.5. *Let $f : \mathcal{F} \rightarrow \mathcal{G}$ be a morphism of sheaves. Then*

1. *f is injective if and only if $f_p : \mathcal{F}_p \rightarrow \mathcal{G}_p$ is injective for all p , and*
2. *f is surjective if and only if $f_p : \mathcal{F}_p \rightarrow \mathcal{G}_p$ is surjective for all p .*

Proof.

1. f_p is injective for all p if and only if $\ker f_p = 0$ for all p , if and only if $(\ker f)_p = 0$ for all p , if and only if $\ker f = 0$,⁴ which is if and only if f is injective.
2. f_p is surjective for all p if and only if $\text{im } f_p = \mathcal{G}_p$ for all p , if and only if $(\text{im } f)_p = \mathcal{G}_p$ for all p , if and only if $\text{im } f = \mathcal{G}$,⁵ which is if and only if f is surjective.

□

Remark. Given $f : \mathcal{F} \rightarrow \mathcal{G}$, in fact $\mathcal{G}/\text{im } f \cong \text{coker } f$.⁶

1.6 Passing between spaces

Let $f : X \rightarrow Y$ be a continuous map between topological spaces, \mathcal{F} a sheaf on X , and \mathcal{G} a sheaf on Y . Define $f_*\mathcal{F}$ by, for $U \subseteq Y$

$$(f_*\mathcal{F})(U) = \mathcal{F}(f^{-1}(U)).$$

Exercise. Check $f_*\mathcal{F}$ is a sheaf on Y .

Define $f^{-1}\mathcal{G}$ to be the sheaf associated to the presheaf

$$U \subseteq X \mapsto \{(V, s) \mid V \supseteq f(U), V \text{ open}, s \in \mathcal{G}(V)\} / \sim,$$

where $(V, s) \sim (V', s')$ if there exists $W \subseteq V \cap V'$ such that $f(U) \subseteq W$, and $s|_W = s'|_W$.

Example. If $f : \{p\} \rightarrow X$ is an inclusion of a point, then $f^{-1}\mathcal{G} = \mathcal{G}_p$. This is a group but defines a sheaf on a one-point space. More generally, if $\iota : Z \hookrightarrow X$ is an inclusion of a subset with induced topology, we often write

$$\mathcal{F}|_Z = \iota^{-1}\mathcal{F}.$$

If Z is open in X , then this is easy, since if $U \subseteq Z$ then $\mathcal{F}|_Z(U) = \mathcal{F}(U)$.

Remark. If $s \in \mathcal{F}(U)$ we say s is a **section** of \mathcal{F} over U . We often write

$$\mathcal{F}(U) = \Gamma(U, \mathcal{F}),$$

thinking of $\Gamma(U, \cdot)$ as a functor from the category of sheaves on a space X to the category of abelian groups.

³Exercise: check

⁴Exercise: check by S1

⁵Exercise: check using $\text{im } f \subseteq \mathcal{G}$

⁶Exercise

2 Schemes

Want to construct a sheaf \mathcal{O} on $\text{Spec } A$, analogous to the sheaf of regular functions on a variety, and \mathcal{O} will be a sheaf of rings. That is, $\mathcal{O}(U)$ will be a ring for each open set U and restriction maps will be ring homomorphisms.

2.1 Localisation of a ring

Importantly recall the following. Let A be a ring, where all rings are commutative with unity, and $S \subseteq A$ be a multiplicatively closed subset, that is $1 \in S$ and if $s_1, s_2 \in S$ then $s_1 s_2 \in S$. We define a ring

$$S^{-1}A = \{(a, s) \mid a \in A, s \in S\} / \sim,$$

where $(a, s) \sim (a', s')$ if there exists $s'' \in S$ such that $s''(as' - a's) = 0$. Then $S^{-1}A$ is called the **localisation of A at S** . Note that we write a/s for the equivalence class of (a, s) . The usual equivalence relation on fractions is $a/s = a'/s'$ if and only if $as' = a's$. We need the extra possibility of killing $as' - a's$ with s'' if A is not an integral domain.

Example.

- Take $f \in A$ and $S = \{1, f, \dots\} \subseteq A$. Then we write $A_f = S^{-1}A$. These will correspond to open subsets.
- If $\mathfrak{p} \subseteq A$ is a prime ideal and $S = A \setminus \mathfrak{p}$, then
 - $1 \in S$, and
 - $a, b \in S$ and $ab \in \mathfrak{p}$ is a contradiction by definition of prime ideals, so $ab \in S$.

Then $A_{\mathfrak{p}} = S^{-1}A$ is the **localisation of A at \mathfrak{p}** . These will correspond to stalks.

2.2 Construction of the structure sheaf

Let

$$\mathcal{O}(U) = \left\{ s : U \rightarrow \bigsqcup_{\mathfrak{p} \in U} A_{\mathfrak{p}} \mid \begin{array}{l} \forall \mathfrak{p} \in U, s(\mathfrak{p}) \in A_{\mathfrak{p}} \\ \forall \mathfrak{p} \in U, \exists \mathfrak{p} \in V \subseteq U \text{ open}, \exists a, f \in A, \forall \mathfrak{q} \in V, f \notin \mathfrak{q}, s(\mathfrak{q}) = \frac{a}{f} \in A_{\mathfrak{q}} \end{array} \right\}.$$

Proposition 2.1. For any $\mathfrak{p} \in \text{Spec } A$, $\mathcal{O}_{\mathfrak{p}} = A_{\mathfrak{p}}$.

Proof. Have a map

$$\begin{array}{ccc} \mathcal{O}_{\mathfrak{p}} & \longrightarrow & A_{\mathfrak{p}} \\ (U, s) & \longmapsto & s(\mathfrak{p}) \end{array}.$$

- Surjective. Any element of $A_{\mathfrak{p}}$ can be written as a/f for some $a \in A$ and $f \notin \mathfrak{p}$. Then $\mathbb{D}(f) = \text{Spec } A \setminus \mathbb{V}(f) = \{\mathfrak{p} \in \text{Spec } A \mid f \notin \mathfrak{p}\}$, since $\mathbb{V}(f) = \{\mathfrak{p} \in \text{Spec } A \mid f \in \mathfrak{p}\}$. Now a/f defines an element of $\mathcal{O}(\mathbb{D}(f))$ given by

$$\begin{array}{ccc} s & : & \mathbb{D}(f) \longrightarrow A_{\mathfrak{q}} \\ & & \mathfrak{q} \longmapsto \frac{a}{f} \end{array},$$

and in particular, $s(\mathfrak{p}) = a/f \in A_{\mathfrak{p}}$.

- Injective. Let $\mathfrak{p} \in U \subseteq \text{Spec } A$ and $s \in \mathcal{O}(U)$ with $s(\mathfrak{p}) = 0$ in $A_{\mathfrak{p}}$. Want to show $(U, s) = 0$ in $\mathcal{O}_{\mathfrak{p}}$. By shrinking U if necessary, we can assume that s is given by $a, f \in A$ with $s(\mathfrak{q}) = a/f$ for all $\mathfrak{q} \in U$. In particular $f \notin \mathfrak{q}$ for all $\mathfrak{q} \in U$. Thus $a/f = 0/1$ in $A_{\mathfrak{p}}$ so there exists $h \in A \setminus \mathfrak{p}$ such that $0 = h \cdot (a \cdot 1 - f \cdot 0) = h \cdot a$ in A . Now let $V = U \cap \mathbb{D}(h)$. Then $(V, s|_V) = 0$, since for $\mathfrak{q} \in V$, $s|_V(\mathfrak{q}) = s(\mathfrak{q}) = a/f \in A_{\mathfrak{q}}$ and $h \cdot a = 0$, and $h \in A \setminus \mathfrak{q}$ so $h \cdot a = 0$ implies $a/f = 0/1$ in $A_{\mathfrak{q}}$. Thus $(U, s) = 0$ in $\mathcal{O}_{\mathfrak{p}}$.

□

Proposition 2.2. For any $f \in A$, $\mathcal{O}(\mathbb{D}(f)) = A_f$.

In particular, as $\text{Spec } A = \mathbb{D}(1)$, the **global sections** of \mathcal{O} is $\mathcal{O}(\text{Spec } A) = A_1 = A$.

Proof. Let

$$\begin{aligned} \psi : A_f &\longrightarrow \mathcal{O}(\mathbb{D}(f)) \\ \frac{a}{f^n} &\longmapsto \left(\mathfrak{p} \in \mathbb{D}(f) \mapsto \frac{a}{f^n} \in A_{\mathfrak{p}} \right), \end{aligned}$$

since $f \notin \mathfrak{p}$ implies that $f^n \notin \mathfrak{p}$ for all $n \geq 0$.

- **Injective.** If $\psi(a/f^n) = 0$, then for all $\mathfrak{p} \in \mathbb{D}(f)$, $a/f^n = 0$ in $A_{\mathfrak{p}}$, that is there exists $h \in A \setminus \mathfrak{p}$ such that $h \cdot a = 0$ in A . Let $I = \{g \in A \mid g \cdot a = 0\}$, the **annihilator** of a . So $h \in I$ and $h \notin \mathfrak{p}$, so $I \not\subseteq \mathfrak{p}$. This is true for all $\mathfrak{p} \in \mathbb{D}(f)$, so $\mathbb{V}(I) \cap \mathbb{D}(f) = \emptyset$. Thus $f \in \bigcap_{\mathfrak{p} \in \mathbb{V}(I)} \mathfrak{p} = \sqrt{I}$, the radical, so $f^m \in I$ for some $m > 0$. Thus $f^m \cdot a = 0$, so $a/f^n = 0$ in A_f . Thus ψ is injective.
- **Surjective.** Let $s \in \mathcal{O}(\mathbb{D}(f))$. Cover $\mathbb{D}(f)$ with open sets V_i on which s is represented as a_i/g_i with $a_i, g_i \in A$ such that $g_i \notin \mathfrak{p}$ whenever $\mathfrak{p} \in V_i$. Thus $V_i \subseteq \mathbb{D}(g_i)$. By question 1 on example sheet 1, the sets of the form $\mathbb{D}(h)$ form a base for the Zariski topology on $\text{Spec } A$. Thus we can assume $V_i = \mathbb{D}(h_i)$ for some $h_i \in A$. Since $\mathbb{D}(h_i) \subseteq \mathbb{D}(g_i)$, we have $\mathbb{V}(h_i) \supseteq \mathbb{V}(g_i)$, so $\sqrt{\langle h_i \rangle} \subseteq \sqrt{\langle g_i \rangle}$, so $h_i^n \in \langle g_i \rangle$ for some n , say $h_i^n = c_i g_i$, so $a_i/g_i = c_i a_i/h_i^n$. Now replace h_i by h_i^n , since this does not change open sets because in general $\mathbb{D}(h_i) = \mathbb{D}(h_i^n)$, and replace a_i by $c_i a_i$. The situation so far is that we can assume $\mathbb{D}(f)$ is covered by sets $\mathbb{D}(h_i)$ such that s is represented by a_i/h_i on $\mathbb{D}(h_i)$. Claim that $\mathbb{D}(f)$ can be covered by a finite number of the $\mathbb{D}(h_i)$, that is $\mathbb{D}(f)$ is quasi-compact. Since

$$\begin{aligned} \mathbb{D}(f) \subseteq \bigcup_i \mathbb{D}(h_i) &\iff \mathbb{V}(f) \supseteq \bigcap_i \mathbb{V}(h_i) = \mathbb{V}\left(\sum_i \langle h_i \rangle\right) &\iff f \in \bigcap_{\mathfrak{p} \in \mathbb{V}(\sum_i \langle h_i \rangle)} \mathfrak{p} \\ &\iff f \in \sqrt{\sum_i \langle h_i \rangle} &\iff \exists n, f^n \in \sum_i \langle h_i \rangle, \end{aligned}$$

we can write $f^n = \sum_{i \in I} b_i h_i$ for some finite index set I . Thus reversing this argument, $\mathbb{D}(f) \subseteq \bigcup_{i \in I} \mathbb{D}(h_i)$. We now pass to this finite subcover $\{\mathbb{D}(h_i)\}$. On $\mathbb{D}(h_i) \cap \mathbb{D}(h_j) = \mathbb{D}(h_i h_j)$, note a_i/h_i and a_j/h_j both represent s , so by injectivity shown in the last lecture, $a_i h_j / h_i h_j = a_i / h_i = a_j / h_j = a_j h_i / h_i h_j$ in $A_{h_i h_j}$. Thus for some n , $(h_i h_j)^n (h_j a_i - h_i a_j) = 0$ in A . We can pick an n sufficiently large to work for all pairs i and j . Rewriting, $h_j^{n+1} (h_i^n a_i) - h_i^{n+1} (h_j a_j) = 0$. Replace each h_i by h_i^{n+1} and a_i by $h_i^n a_i$, since $a_i/h_i = a_i h_i^n / h_i^{n+1}$. Thus we can assume that s is still represented on $\mathbb{D}(h_i)$ by a_i/h_i but also for each i and j have $h_i a_j = h_j a_i$. Note $f^n = \sum_i b_i h_i$ for $b_i \in A$, since $\{\mathbb{D}(h_i)\}$ cover $\mathbb{D}(f)$. Let $a = \sum_i b_i a_i$. Then for any j , $h_j a = \sum_i b_i a_i h_j = \sum_i b_i a_j h_i = f^n a_j$. Thus $a/f^n = a_j/h_j$ on $\mathbb{D}(h_j)$. Thus $\psi(a/f^n) = s$, so ψ is surjective. \square

We now have a topological space $\text{Spec } A$ equipped with a sheaf of rings \mathcal{O} .

2.3 Ringed spaces

Definition. A **ringed space** is a pair (X, \mathcal{O}_X) where

- X is a topological space, and
- \mathcal{O}_X is a sheaf of rings on X .

A **morphism of ringed spaces** $f : (X, \mathcal{O}_X) \rightarrow (Y, \mathcal{O}_Y)$ is the following data.

- $f : X \rightarrow Y$ a continuous map.
- $f^\# : \mathcal{O}_Y \rightarrow f_* \mathcal{O}_X$ a morphism of sheaves of rings, that is for each $U \subseteq Y$ open, we have a ring homomorphism $f_U^\# : \mathcal{O}_Y(U) \rightarrow (f_* \mathcal{O}_X)(U) = \mathcal{O}_X(f^{-1}(U))$.

Lecture 6
Wednesday
21/10/20

Example.

- Let X be a topological space, and let \mathcal{O}_X be the sheaf of continuous \mathbb{R} -valued functions. Then if (Y, \mathcal{O}_Y) is similarly defined, given $f : X \rightarrow Y$, we get $f^\# : \mathcal{O}_Y \rightarrow f_* \mathcal{O}_X$ defined by

$$\begin{aligned} f_U^\# : \mathcal{O}_Y(U) &\longrightarrow \mathcal{O}_X(f^{-1}(U)) \\ \phi &\longmapsto \phi \circ f \end{aligned} .$$

- Let X be a variety, and let \mathcal{O}_X be the sheaf of regular functions on X . A morphism of varieties $f : X \rightarrow Y$ is a continuous map inducing

$$\begin{aligned} f_U^\# : \mathcal{O}_Y(U) &\longrightarrow \mathcal{O}_X(f^{-1}(U)) \\ \phi &\longmapsto \phi \circ f \end{aligned} .$$

A ring is **local** if it has a unique maximal ideal.

Definition. A **locally ringed space** (X, \mathcal{O}_X) is a ringed space such that $\mathcal{O}_{X,p}$ is a local ring for all $p \in X$. A **morphism** $f : (X, \mathcal{O}_X) \rightarrow (Y, \mathcal{O}_Y)$ of **locally ringed spaces** is a morphism of ringed spaces such that the induced homomorphism $f_p^\# : \mathcal{O}_{Y,f(p)} \rightarrow \mathcal{O}_{X,p}$ is a **local homomorphism** for all $p \in X$.

- The map is defined by ⁷

$$\begin{aligned} f_p^\# : \mathcal{O}_{Y,f(p)} &\longrightarrow \mathcal{O}_{X,p} \\ (U, s) &\longmapsto (f^{-1}(U), f_U^\#(s)) \end{aligned} .$$

- A ring homomorphism $\phi : (A, \mathfrak{m}_A) \rightarrow (B, \mathfrak{m}_B)$ is **local** if $\phi^{-1}(\mathfrak{m}_B) = \mathfrak{m}_A$, where \mathfrak{m}_A is the maximal ideal of A . Note that $\phi(A \setminus \mathfrak{m}_A) = \phi(A^*) \subseteq B^* = B \setminus \mathfrak{m}_B$, where A^* is the set of invertible elements of A . Thus $\phi^{-1}(\mathfrak{m}_B) \subseteq \mathfrak{m}_A$ always.

Example. In the case of varieties, $\mathcal{O}_{X,p}$ has a unique maximal ideal

$$\{(U, f) \in \mathcal{O}_X(U) \mid f(p) = 0\} / \sim .$$

If $f(p) \neq 0$, f is nowhere vanishing on some neighbourhood of p , so after shrinking U , we can invert f . The local homomorphism condition just follows from the pull-back $\phi \circ f$ of a function ϕ vanishing at $f(p)$ vanishes at p .

2.4 Affine schemes

The key example $(\text{Spec } A, \mathcal{O})$ is a locally ringed space, which we call an affine scheme.

Theorem 2.3. *The category of affine schemes with locally ringed morphisms is equivalent to the opposite category of rings.*

Need to show that

- if $\phi : A \rightarrow B$ is a ring homomorphism, we obtain an induced morphism $(f, f^\#) : (\text{Spec } B, \mathcal{O}_{\text{Spec } B}) \rightarrow (\text{Spec } A, \mathcal{O}_{\text{Spec } A})$, and
- any morphism of affine schemes as locally ringed spaces arises in this way.

Proof.

- Given a ring homomorphism $\phi : A \rightarrow B$, define

$$\begin{aligned} f : \text{Spec } B &\longrightarrow \text{Spec } A \\ \mathfrak{p} &\longmapsto \phi^{-1}(\mathfrak{p}) \end{aligned} .$$

Note $\phi^{-1}(\mathfrak{p})$ is prime, since if $ab \in \phi^{-1}(\mathfrak{p})$, then $\phi(ab) = \phi(a)\phi(b) \in \mathfrak{p}$, thus either $\phi(a) \in \mathfrak{p}$ or $\phi(b) \in \mathfrak{p}$, and hence either $a \in \phi^{-1}(\mathfrak{p})$ or $b \in \phi^{-1}(\mathfrak{p})$. Then f is continuous, since

$$\begin{aligned} f^{-1}(\mathbb{V}(I)) &= f^{-1}(\{\mathfrak{p} \in \text{Spec } A \mid \mathfrak{p} \supseteq I\}) = \{\mathfrak{q} \in \text{Spec } B \mid f(\mathfrak{q}) \supseteq I\} \\ &= \{\mathfrak{q} \in \text{Spec } B \mid \phi^{-1}(\mathfrak{q}) \supseteq I\} = \{\mathfrak{q} \in \text{Spec } B \mid \mathfrak{q} \supseteq \phi(I)\} = \mathbb{V}(\phi(I)) . \end{aligned}$$

⁷Exercise: check well-defined

We need to construct $f^\# : \mathcal{O}_{\text{Spec } A} \rightarrow f_* \mathcal{O}_{\text{Spec } B}$. For $\mathfrak{p} \in \text{Spec } B$, we obtain a natural homomorphism

$$\begin{array}{ccc} \phi_{\mathfrak{p}} : A_{\phi^{-1}(\mathfrak{p})} & \longrightarrow & B_{\mathfrak{p}} \\ \frac{a}{s} & \longmapsto & \frac{\phi(a)}{\phi(s)} \end{array}.$$

Note $\phi_{\mathfrak{p}}$ is a local homomorphism, since the maximal ideal $\mathfrak{p}B_{\mathfrak{p}}$ of $B_{\mathfrak{p}}$ is generated by the image of \mathfrak{p} under the map

$$\begin{array}{ccc} B & \longrightarrow & B_{\mathfrak{p}} \\ b & \longmapsto & \frac{b}{1} \end{array},$$

and the maximal ideal $\phi^{-1}(\mathfrak{p})A_{\phi^{-1}(\mathfrak{p})}$ of $A_{\phi^{-1}(\mathfrak{p})}$ is generated by the image of $\phi^{-1}(\mathfrak{p})$ under the map

$$\begin{array}{ccc} A & \longrightarrow & A_{\phi^{-1}(\mathfrak{p})} \\ a & \longmapsto & \frac{a}{1} \end{array},$$

so have a commutative diagram

$$\begin{array}{ccccc} \phi^{-1}(\mathfrak{p}) & \subset & A & \xrightarrow{\phi} & B & \supset & \mathfrak{p} \\ & & \downarrow & & \downarrow & & \\ f(\mathfrak{p})A_{f(\mathfrak{p})} & \subset & A_{\phi^{-1}(\mathfrak{p})} & \xrightarrow{\phi_{\mathfrak{p}}} & B_{\mathfrak{p}} & \supset & \mathfrak{p}B_{\mathfrak{p}} \end{array},$$

thus $\phi_{\mathfrak{p}}^{-1}(\mathfrak{p}B_{\mathfrak{p}}) = \phi^{-1}(\mathfrak{p})A_{\phi^{-1}(\mathfrak{p})}$. Given $V \subseteq \text{Spec } A$ open, we may define

$$\begin{array}{ccc} f_V^\# : \mathcal{O}_{\text{Spec } A}(V) & \longrightarrow & \mathcal{O}_{\text{Spec } B}(f^{-1}(V)) \\ (\mathfrak{p} \in V \mapsto s(\mathfrak{p}) \in A_{\mathfrak{p}}) & \longmapsto & (\mathfrak{q} \in f^{-1}(V) \mapsto \phi_{\mathfrak{q}}(s(f(\mathfrak{q}))) \in B_{\mathfrak{q}}) \end{array}.$$

Note that we need to check the local coherence part of the definition of \mathcal{O} . That is, if s is locally given by a/h , then $f_V^\#(s)$ is locally given by $\phi(a)/\phi(h)$. This gives the desired map $f^\# : \mathcal{O}_{\text{Spec } A} \rightarrow f_* \mathcal{O}_{\text{Spec } B}$, and the induced map on stalks $f_{\mathfrak{p}}^\# : \mathcal{O}_{\text{Spec } A, f(\mathfrak{p})} \rightarrow \mathcal{O}_{\text{Spec } B, \mathfrak{p}}$ agrees with $\phi_{\mathfrak{p}} : A_{\phi^{-1}(\mathfrak{p})} \rightarrow B_{\mathfrak{p}}$, by construction. Hence $(f, f^\#)$ is a morphism of locally ringed spaces.

2. Now suppose given a morphism $(f, f^\#) : \text{Spec } B \rightarrow \text{Spec } A$ of locally ringed spaces. Take

$$\phi = f_{\text{Spec } A}^\# : \Gamma(\text{Spec } A, \mathcal{O}_{\text{Spec } A}) = A \rightarrow \Gamma(\text{Spec } B, \mathcal{O}_{\text{Spec } B}) = B.$$

We need to show ϕ gives rise to $(f, f^\#)$. We have $f_{\mathfrak{p}}^\# : \mathcal{O}_{\text{Spec } A, f(\mathfrak{p})} = A_{f(\mathfrak{p})} \rightarrow \mathcal{O}_{\text{Spec } B, \mathfrak{p}} = B_{\mathfrak{p}}$ a local homomorphism. This is compatible with the corresponding map on global sections, that is

$$\begin{array}{ccc} \Gamma(\text{Spec } A, \mathcal{O}_{\text{Spec } A}) & \xrightarrow{f_{\text{Spec } A}^\#} & \Gamma(\text{Spec } B, \mathcal{O}_{\text{Spec } B}) \\ \downarrow & & \downarrow \\ \mathcal{O}_{\text{Spec } A, f(\mathfrak{p})} & \xrightarrow{f_{\mathfrak{p}}^\#} & \mathcal{O}_{\text{Spec } B, \mathfrak{p}} \end{array}$$

is commutative. That is, we have a commutative diagram

$$\begin{array}{ccccc} f(\mathfrak{p}) & \subset & A & \xrightarrow{\phi} & B & \supset & \mathfrak{p} \\ & & \downarrow & & \downarrow & & \\ f(\mathfrak{p})A_{f(\mathfrak{p})} & \subset & A_{f(\mathfrak{p})} & \xrightarrow{f_{\mathfrak{p}}^\#} & B_{\mathfrak{p}} & \supset & \mathfrak{p}B_{\mathfrak{p}} \end{array}.$$

Then $(f_{\mathfrak{p}}^\#)^{-1}(\mathfrak{p}B_{\mathfrak{p}}) = f(\mathfrak{p})A_{f(\mathfrak{p})}$ since $f_{\mathfrak{p}}^\#$ is a local homomorphism, and by commutativity of the diagram, $f(\mathfrak{p}) = \phi^{-1}(\mathfrak{p})$. Thus f is induced by ϕ , and $f_{\mathfrak{p}}^\# = \phi_{\mathfrak{p}}$. So $f^\#$ is as constructed previously. \square

Remark. Demanding $(f, f^\#)$ was a morphism of locally ringed spaces was crucial to make the proof work.

Definition. An **affine scheme** is a locally ringed space isomorphic, in the category of locally ringed spaces, to $(\operatorname{Spec} A, \mathcal{O}_{\operatorname{Spec} A})$ for some ring A . A **scheme** is a locally ringed space (X, \mathcal{O}_X) with an open cover $\{(U_i, \mathcal{O}_X|_{U_i})\}$ with each $(U_i, \mathcal{O}_X|_{U_i})$ an affine scheme, where $\mathcal{O}_X|_{U_i}(V) = \mathcal{O}_X(V)$ for $V \subseteq U_i$ open. A **morphism of schemes** is a morphism of locally ringed spaces.

Example. Let k be a field. Then $\operatorname{Spec} k = (\{0\}, k)$.

- What does giving a morphism $f : \operatorname{Spec} k \rightarrow X$ to a scheme mean? First, this selects a point $x \in X$, the image of f . Second, we get a local ring homomorphism $f_x^\# : \mathcal{O}_{X,x} \rightarrow \mathcal{O}_{\operatorname{Spec} k,0} = k$, that is $(f_x^\#)^{-1}(0) = \mathfrak{m}_x \subseteq \mathcal{O}_{X,x}$, the maximal ideal of $\mathcal{O}_{X,x}$. Thus we get a factorisation $f_x^\# : \mathcal{O}_{X,x} \rightarrow \mathcal{O}_{X,x}/\mathfrak{m}_x \rightarrow k$, where $\mathcal{O}_{X,x}/\mathfrak{m}_x$ is a field, written as $\kappa(x)$, called the **residue field** of X at x . Thus f induces an inclusion $\kappa(x) \hookrightarrow k$. Conversely, given such an inclusion $\iota : \kappa(x) \hookrightarrow k$ of fields, we get a scheme morphism by defining $f(0) = x$, and

$$\begin{array}{ccc} f^\# & : & \mathcal{O}_X \longrightarrow f_*k \\ s & \longmapsto & \iota(s(x)) \end{array}, \quad s(x) \in \mathcal{O}_{X,x}.$$

The moral is that giving a morphism $f : \operatorname{Spec} k \rightarrow X$ is equivalent to giving a point $x \in X$ and an inclusion $\iota : \kappa(x) \rightarrow k$. Note that if $X = \operatorname{Spec} A$, giving $\operatorname{Spec} k \rightarrow \operatorname{Spec} A$ is equivalent to giving a homomorphism $A \rightarrow k$, which we viewed at the beginning of the course as a k -valued point on $\operatorname{Spec} A$.

- What does giving $X \rightarrow \operatorname{Spec} k$ mean? No information in the continuous map, but need also a map $f^\# : k \rightarrow f_*\mathcal{O}_X$, that is a map $k \rightarrow \Gamma(\operatorname{Spec} k, f_*\mathcal{O}_X) = \Gamma(X, \mathcal{O}_X)$. That is, $\Gamma(X, \mathcal{O}_X)$ carries a k -algebra structure. Note this induces k -algebra structures on $\mathcal{O}_X(U)$ for all U via the composition $k \rightarrow \mathcal{O}_X(X) \rightarrow \mathcal{O}_X(U)$ and similarly all stalks $\mathcal{O}_{X,p}$ are also k -algebras. We say X is a **scheme defined over k** . For example, in affine varieties, consider $A = k[X_1, \dots, X_n]/I$ with $I = \sqrt{I}$. Then $\operatorname{Spec} A$ is our replacement for $\mathbb{V}(I) \subseteq \mathbb{A}_k^n$, viewing $\operatorname{Spec} A$ as a scheme over k . If $k \subseteq k'$ is a field extension, a **k' -valued point** of X/k is a commutative diagram

$$\begin{array}{ccc} \operatorname{Spec} k' & \xrightarrow{\quad} & X \\ & \searrow & \swarrow \\ & \operatorname{Spec} k & \end{array}.$$

We write $X(k')$ for the set of such morphisms.

Remark. It is rare in algebraic geometry to work with schemes alone, but rather always working over a base scheme.

Fix a base scheme S . Define **Sch**/ S to be the category whose objects are morphisms $T \rightarrow S$ and morphisms are commutative diagrams

$$\begin{array}{ccc} T & \xrightarrow{\quad} & T' \\ & \searrow & \swarrow \\ & S & \end{array}.$$

We will frequently work with **Sch**/ $k = \mathbf{Sch}/\operatorname{Spec} k$. Given $T \rightarrow S$ and $X \rightarrow S$ objects in **Sch**/ S , a **T -valued point** of $X \rightarrow S$ is a morphism $T \rightarrow X$ over S , so

$$\begin{array}{ccc} T & \xrightarrow{\quad} & X \\ & \searrow & \swarrow \\ & S & \end{array},$$

and we write $X(T)$ for the set of T -valued points. The **Yoneda philosophy** is that $X(T)$ for all T determines X .

Lecture 8
Monday
26/10/20

Example. Fix a field k , and let $D = \operatorname{Spec} k[t] / \langle t^2 \rangle = (\{t\}, k[t] / \langle t^2 \rangle)$. Then t does not make sense as k -valued function anymore, as $t^2 = 0$. Let X be any scheme over k . What is $X(D)$? Given $f : D \rightarrow X$ a morphism of schemes over k , we get a point $x \in X$ as the image of f and a local homomorphism

$$\begin{aligned} f_x^\# : \mathcal{O}_{X,x} &\longrightarrow k[t] / \langle t^2 \rangle \\ \mathfrak{m}_x &\longmapsto \langle t \rangle \end{aligned}.$$

Note that \mathfrak{m}_x^2 maps to zero, hence we get a k -linear map $\mathfrak{m}_x / \mathfrak{m}_x^2 \rightarrow \langle t \rangle \cong k$ as a k -vector space. We also have a composed surjective k -algebra homomorphism $\mathcal{O}_{X,x} \rightarrow k[t] / \langle t \rangle \cong k$ with kernel \mathfrak{m}_x , and hence we have $\kappa(x) = \mathcal{O}_{X,x} / \mathfrak{m}_x \cong k$. So we get

- a k -valued point x with residue field k , and
- a k -vector space map $\mathfrak{m}_x / \mathfrak{m}_x^2 \rightarrow k$, that is an element of $(\mathfrak{m}_x / \mathfrak{m}_x^2)^*$, the dual vector space.

Then $(\mathfrak{m}_x / \mathfrak{m}_x^2)^*$ is called the **Zariski tangent space** to X at x . Think of D as a point plus an arrow.

Example. Glued schemes are a special case of a question on example sheet 1. Suppose given two schemes X_1 and X_2 and open subsets $U_i \subseteq X_i$. Recall U_i is also a locally ringed space $(U_i, \mathcal{O}_{X_i}|_{U_i})$, and in fact U_i is then a scheme. Given an isomorphism $f : U_1 \xrightarrow{\sim} U_2$, can glue X_1 and X_2 along U_1 and U_2 to get a scheme X with an open cover $\{X_1, X_2\}$, so $X = X_1 \sqcup X_2 / \sim$ such that $x_1 \in U_1 \sim x_2 \in U_2$ if $f(x_1) = x_2$, and need to define \mathcal{O}_X . Now take $\mathbb{A}_k^n = \operatorname{Spec} k[X_1, \dots, X_n]$, so $\mathbb{A}_k^1 = \operatorname{Spec} k[X]$. Take $X_1 = X_2 = \mathbb{A}_k^1$.

- Glue $U_1 = \mathbb{A}^1 \setminus \{0\} = \mathbb{D}(X) \subseteq X_1$ and $U_2 = \mathbb{A}^1 \setminus \{0\} = \mathbb{D}(X) \subseteq X_2$ via the identity map. This is the affine line with doubled origin.
- Could instead glue U_1 and U_2 via the map given by $X \mapsto X^{-1}$, where $U_1 = \operatorname{Spec} k[X]_X = U_2$ and

$$\begin{aligned} k[X]_X &\longrightarrow k[X]_X \\ X &\longmapsto X^{-1} \end{aligned}$$

induces an isomorphism $U_1 \rightarrow U_2$. When we glue, we get the projective line over k , \mathbb{P}_k^1 .

2.5 Projective schemes

Let S be a graded ring, that is

$$S = \bigoplus_{d \geq 0} S_d,$$

with S_d an abelian group, and product law satisfies $S_d \cdot S_{d'} \subseteq S_{d+d'}$.

Example. $S = k[X_0, \dots, X_n]$, and S_d is the space of polynomials which are homogeneous of degree d , that is spanned by monomials of degree d .

We write

$$S_+ = \bigoplus_{d \geq 1} S_d,$$

which we call the **irrelevant ideal**.

Definition. $I \subseteq S$ is a **homogeneous ideal** if I is generated by its homogeneous elements, that is elements in S_d for various d .

Definition. Let

$$\operatorname{Proj} S = \{\mathfrak{p} \in \operatorname{Spec} S \mid \mathfrak{p} \text{ is homogeneous, } \mathfrak{p} \not\supseteq S_+\}.$$

For $I \subseteq S$ a homogeneous ideal, set

$$\mathbb{V}(I) = \{\mathfrak{p} \in \operatorname{Proj} S \mid \mathfrak{p} \supseteq I\}.$$

Exercise. Check the $\mathbb{V}(I)$ form the closed sets of a topology on $\operatorname{Proj} S$.

Notation. For $\mathfrak{p} \in \text{Proj } S$, let

$$T = \{f \in S \setminus \mathfrak{p} \mid f \text{ is homogeneous}\}.$$

Then T is a multiplicatively closed subset of S , and let $S_{(\mathfrak{p})} \subseteq T^{-1}S$ be the subring of elements of degree zero, that is written in the form s/s' with $s \in S$ homogeneous and $s' \in T$ with $\deg s = \deg s'$. For $f \in S$ homogeneous, we write $S_{(f)} \subseteq S_f$ for the subset of elements of degree zero.

Can now define a sheaf \mathcal{O} on $\text{Proj } S$. For $U \subseteq \text{Proj } S$ open, set

$$\mathcal{O}(U) = \left\{ s : U \rightarrow \bigsqcup_{\mathfrak{p} \in U} S_{(\mathfrak{p})} \mid \begin{array}{l} \forall \mathfrak{p} \in U, s(\mathfrak{p}) \in S_{(\mathfrak{p})} \\ \forall \mathfrak{p} \in U, \exists V \subseteq U \text{ open, } \exists a, f \in S, \forall \mathfrak{q} \in V, f \notin \mathfrak{q}, s(\mathfrak{q}) = \frac{a}{f} \in S_{(\mathfrak{q})} \end{array} \right\},$$

where a and f are homogeneous of the same degree. As before, $\mathcal{O}_{\mathfrak{p}} = S_{(\mathfrak{p})}$.⁸ Is the locally ringed space $(\text{Proj } S, \mathcal{O})$ a scheme?

Notation. If $f \in S$ is homogeneous, then we write

$$\mathbb{D}_+(f) = \{\mathfrak{p} \in \text{Proj } S \mid f \notin \mathfrak{p}\},$$

which is an open set and $\mathbb{D}_+(f) = \text{Proj } S \setminus \mathbb{V}(f)$.

Proposition 2.4. $(\mathbb{D}_+(f), \mathcal{O}|_{\mathbb{D}_+(f)}) \cong \text{Spec } S_{(f)}$ as locally ringed spaces. Further, the open sets $\mathbb{D}_+(f)$ for $f \in S_+$ cover $\text{Proj } S$. Hence $(\text{Proj } S, \mathcal{O})$ is a scheme.

Proof. Will be on example sheet 2. □

Definition. If A is a ring, define

$$\mathbb{P}_A^n = \text{Proj } A[X_0, \dots, X_n].$$

Example. If k is an algebraically closed field, consider $\mathbb{P}_k^1 = \text{Proj } k[X_0, X_1]$. The closed points, that is points \mathfrak{p} such that $\{\mathfrak{p}\}$ is closed, correspond to maximal elements of $\text{Proj } S$.⁹ These maximal elements are ideals of the form $\langle aX_0 - bX_1 \rangle$. The only maximal homogeneous ideal of $k[X_0, X_1]$ is $\langle X_0, X_1 \rangle = S_+$, since any maximal ideal is of the form $\langle X_0 - a_0, X_1 - a_1 \rangle$. The other prime ideals of $k[X_0, X_1]$ are principal, that is of the form $\langle f \rangle$ with f irreducible or $f = 0$. For $\langle f \rangle$ to be homogeneous, f must be homogeneous. Any such polynomial splits into linear factors, all homogeneous, so in order for f to be irreducible it must be linear. Note we have a one-to-one correspondence between

$$\begin{aligned} \{\langle aX_0 - bX_1 \rangle \mid a, b \in K, a, b \text{ not both zero}\} &\longrightarrow (k^2 \setminus \{(0, 0)\})/k^* \\ \langle aX_0 - bX_1 \rangle &\longmapsto (b : a) \end{aligned},$$

where k^* acts by $(a, b) \mapsto (\lambda a, \lambda b)$ for $\lambda \in k^*$. The conclusion is that the closed points of \mathbb{P}_k^1 are in one-to-one correspondence with points of $(k^2 \setminus \{(0, 0)\})/k^*$. More generally, the closed points of \mathbb{P}_k^n are in one-to-one correspondence with points of $(k^{n+1} \setminus \{0\})/k^*$. Can see this by making use of the open cover $\{\mathbb{D}_+(X_i) \mid 0 \leq i \leq n\}$,¹⁰ which is an open cover since $\mathfrak{p} \notin \mathbb{D}_+(X_i)$ for any i implies that $X_i \in \mathfrak{p}$ for all i , so $S_+ \subseteq \mathfrak{p}$ and so $\mathfrak{p} \notin \text{Proj } S$.

Example. Let $S = k[X_0, \dots, X_n]$, but grade by $\deg X_i = w_i$, where w_0, \dots, w_n are positive integers. Define $\mathbb{WP}^n(w_0, \dots, w_n) = \text{Proj } S$, the **weighted projective space**. For example, $\mathbb{WP}^2(1, 1, 2)$ has an open cover $\{\mathbb{D}_+(X_i) \mid 0 \leq i \leq 2\}$. Consider $\mathbb{D}_+(X_2) = \text{Spec } S_{(X_2)}$. Note

$$S_{(X_2)} = k\left[\frac{X_0^2}{X_2}, \frac{X_0X_1}{X_2}, \frac{X_1^2}{X_2}\right] \cong k[U, V, W] / \langle UW - V^2 \rangle \subseteq S_{X_2},$$

so $\text{Spec } S_{(X_2)}$ is a quadric cone with a singular point. Similarly, $\mathbb{D}_+(X_0)$ and $\mathbb{D}_+(X_1)$ are both isomorphic to \mathbb{A}_k^2 .

⁸Exercise: check

⁹Exercise: check

¹⁰Exercise: good exercise

Example. Let $M = \mathbb{Z}^n$ and $M_{\mathbb{R}} = M \otimes_{\mathbb{Z}} \mathbb{R} = \mathbb{R}^n$. Let $\Delta \subseteq M_{\mathbb{R}}$ be a compact convex lattice polytope. That is, there exists a finite set $V \subseteq M$ such that Δ is the convex hull of V , that is the smallest convex set containing V . Let

$$C(\Delta) = \{(m, r) \in M_{\mathbb{R}} \oplus \mathbb{R} \mid m \in r\Delta, r \geq 0\} \subseteq M_{\mathbb{R}} \oplus \mathbb{R}.$$

Here $r\Delta = \{rm \mid m \in \Delta\}$. This is the **cone over Δ** . Let

$$S = k[C(\Delta) \cap (M \oplus \mathbb{Z})] = \bigoplus_{P \in C(\Delta) \cap (M \oplus \mathbb{Z})} kZ^P,$$

with multiplication given by $Z^P Z^{P'} = Z^{P+P'}$, since $C(\Delta) \cap (M \oplus \mathbb{Z})$ is a monoid, that is it is closed under addition and contains zero. This makes S into a ring, and it is graded by $\deg Z^{(m,r)} = r$. Define $\mathbb{P}_{\Delta} = \text{Proj } S$. This is called a **projective toric variety**.

- Let Δ be the convex hull of $\{0, e_1, \dots, e_n\}$ with e_1, \dots, e_n the standard basis of $M = \mathbb{Z}^n$. Check that $S = k[X_0, \dots, X_n]$ with standard grading $X_0 = Z^{(0,1)}$ and $X_i = Z^{(e_i,1)}$.¹¹ So $\mathbb{P}_{\Delta} = \mathbb{P}_k^n$.
- Let $n = 2$, and let Δ be the convex hull of $\{(0,0), (1,0), (0,1), (1,1)\}$. In S , the degree d monomials are $\{Z^{(a,b,d)} \mid 0 \leq a \leq d, 0 \leq b \leq d\}$. Any of these can be written as a product of monomials of degree one, that is the monomials $X = Z^{(0,0,1)}$, $Y = Z^{(1,0,1)}$, $W = Z^{(0,1,1)}$, and $T = Z^{(1,1,1)}$. Thus $S = k[X, Y, W, T] / \langle XT - YW \rangle$. So $\text{Proj } S$ can be thought of as a quadric surface in \mathbb{P}_k^3 .

2.6 Open and closed subschemes

Definition. An **open subscheme** of a scheme X is a scheme $(U, \mathcal{O}_X|_U)$ for $U \subseteq X$ an open subset. Note that this is a scheme because from question 1 and question 11 on the first example sheet, open affine subsets of X form a basis for the topology on X . An **open immersion** is a morphism $f : X \rightarrow Y$ which induces an isomorphism of X with an open subscheme of Y . A **closed immersion** $f : X \rightarrow Y$ is a morphism which is a homeomorphism onto a closed subset of Y , and the induced morphism $f^{\#} : \mathcal{O}_Y \rightarrow f_*\mathcal{O}_X$ is surjective. A **closed subscheme** of Y is an equivalence class of closed immersions, where

$$\begin{array}{ccc} X & \xrightarrow{\quad i \quad} & X' \\ & \searrow & \swarrow \\ & Y & \end{array}$$

are equivalent if there exists an isomorphism i making the diagram commute.

Example.

- Let $Y = \text{Spec } A$, let $I \subseteq A$ be an ideal, and let $X = \text{Spec } A/I$. Note the map of schemes induced by the quotient map $A \rightarrow A/I$ identifies $\text{Spec } A/I$ with $\mathbb{V}(I) \subseteq \text{Spec } A$. Thus $f : X \rightarrow Y$, induced by $A \rightarrow A/I$, satisfies the first condition of being a closed immersion. Note that $\mathcal{O}_Y \rightarrow f_*\mathcal{O}_X$ is surjective on stalks. For $\mathfrak{p} \in \mathbb{V}(I)$, $\mathcal{O}_{Y,\mathfrak{p}} = A_{\mathfrak{p}}$ and $(f_*\mathcal{O}_X)_{\mathfrak{p}} = \mathcal{O}_{X,\mathfrak{p}}$ since all open sets in X are of the form $U \cap X$ for U an open set of Y and $\mathcal{O}_{X,\mathfrak{p}} = (A/I)_{\mathfrak{p}/I}$. Certainly $A_{\mathfrak{p}} \rightarrow (A/I)_{\mathfrak{p}/I}$ is surjective.
- Let $\text{Spec } k[X, Y] / \langle X \rangle \rightarrow \text{Spec } k[X, Y] = \mathbb{A}^2$. This gives a closed subscheme structure to the set $\mathbb{V}(X)$. Note $\mathbb{V}(X^2, XY) = \mathbb{V}(X)$. This gives a closed immersion $\text{Spec } k[X, Y] / \langle X^2, XY \rangle \rightarrow \mathbb{A}^2$. This gives a different closed subscheme structure on $\mathbb{V}(X)$. Note these two subschemes are isomorphic away from the origin, which we can see by looking at $\mathbb{D}(Y) \subseteq \text{Spec } k[X, Y] / \langle X \rangle$, where

$$\mathbb{D}(Y) \cong \text{Spec } (k[X, Y] / \langle X \rangle)_Y = \text{Spec } k[Y]_Y.$$

Looking at $\mathbb{D}(Y) \subseteq \text{Spec } k[X, Y] / \langle X^2, XY \rangle$,

$$\mathbb{D}(Y) \cong \text{Spec } (k[X, Y] / \langle X^2, XY \rangle)_Y \cong \text{Spec } (k[X, Y]_Y / \langle X \rangle) \cong \text{Spec } k[Y]_Y.$$

¹¹Exercise

2.7 Fibre products

Let \mathcal{C} be a category and

$$\begin{array}{ccc} & Y & \\ & \downarrow g & \\ X & \xrightarrow{f} & Z \end{array}$$

be a diagram in \mathcal{C} . Then the **fibre product**, if it exists, is an object W equipped with morphisms $p : W \rightarrow X$ and $q : W \rightarrow Y$ such that $f \circ p = g \circ q$ satisfying the following universal property. For any W' equipped with maps $p' : W' \rightarrow X$ and $q' : W' \rightarrow Y$ such that $f \circ p' = g \circ q'$, there exists a unique morphism $h : W' \rightarrow W$ making the diagram

$$\begin{array}{ccccc} W' & & \xrightarrow{q'} & & Y \\ & \searrow \exists! h & & \searrow q & \\ & & W & \xrightarrow{q} & Y \\ & \swarrow p' & \downarrow p & & \downarrow g \\ & & X & \xrightarrow{f} & Z \end{array}$$

commute, that is $p \circ h = p'$ and $q \circ h = q'$. Note that if the fibre product exists, it is unique up to unique isomorphism.

Example. Let \mathcal{C} be the category of sets. Then

$$X \times_Z Y = \{(x, y) \in X \times Y \mid f(x) = g(y)\}.$$

It will be helpful to think about the fibre product, and more generally other universal properties, via the Yoneda lemma.

Definition. Let \mathcal{C} be a category. Write h_X for the contravariant functor

$$\begin{array}{rcl} h_X : & \mathcal{C} & \longrightarrow \mathbf{Set} \\ & Y & \longmapsto \mathrm{Hom}(Y, X) \\ f : Y \rightarrow Z & \longmapsto & (\phi \in \mathrm{Hom}(Z, X) \mapsto \phi \circ f \in \mathrm{Hom}(Y, X)) \end{array}.$$

Recall that a **natural transformation** between contravariant functors $F, G : \mathcal{C} \rightarrow \mathcal{D}$, written as $T : \mathcal{C} \rightarrow \mathcal{D}$, consists of the data $T(X) : F(X) \rightarrow G(X)$ for all $X \in \mathrm{Ob} \mathcal{C}$ such that for all $f : X \rightarrow Y$ in \mathcal{C}

$$\begin{array}{ccc} F(X) & \xleftarrow{F(f)} & F(Y) \\ T(X) \downarrow & & \downarrow T(Y) \\ G(X) & \xleftarrow{G(f)} & G(Y) \end{array}$$

is commutative.

Lemma 2.5 (Yoneda's lemma). *The set of natural transformations between $h_X : \mathcal{C} \rightarrow \mathbf{Set}$ and $G : \mathcal{C} \rightarrow \mathbf{Set}$ is $G(X)$.*

Proof. Given $\eta \in G(X)$, we need to define a map

$$\begin{array}{ccc} h_X(Y) = \mathrm{Hom}(Y, X) & \longrightarrow & G(Y) \\ f & \longmapsto & G(f)(\eta) \end{array},$$

for all objects $Y \in \mathcal{C}$. Check that this defines a natural transformation $h_X \rightarrow G$.¹² Conversely, given $T : h_X \rightarrow G$ a natural transformation, take $\eta = T(X)(\mathrm{id}_X)$. Check that these two maps are inverse to each other.¹³ \square

Corollary 2.6. *The set of natural transformations $h_X \rightarrow h_Y$ is $h_Y(X) = \mathrm{Hom}(X, Y)$.*

¹²Exercise

¹³Exercise

Definition. A contravariant functor $F : \mathcal{C} \rightarrow \mathbf{Set}$ is said to be **representable** if $F \cong h_X$ for some $X \in \text{Ob } \mathcal{C}$.

Lots of questions in algebraic geometry are about representability of functors. Redefining, the fibre product in a category \mathcal{C} is an object which represents the functor

$$T \mapsto \text{Hom}(T, X) \times_{\text{Hom}(T, Z)} \text{Hom}(T, Y),$$

since an element of the set $\text{Hom}(T, X) \times_{\text{Hom}(T, Z)} \text{Hom}(T, Y)$ is a commutative diagram

$$\begin{array}{ccccc} T & & & & \\ & \searrow q & & & \\ & & W & \dashrightarrow & Y \\ & & \downarrow & & \downarrow g \\ & & X & \xrightarrow{f} & Z \\ & \nearrow p & & & \end{array}$$

The advantage of using Yoneda is that we can check identities using fibre products using identities of the products of sets.

Example. In **Set**,

$$\begin{aligned} (A \times_B B) \times_C D &\longleftrightarrow A \times_B D \\ ((a, c), d) &\longmapsto (a, d) \\ ((a, f(d)), d) &\longleftarrow (a, d) \end{aligned}, \quad f : D \rightarrow C.$$

Then we have two functors

$$\begin{array}{ccc} T & \longrightarrow & h_A(T) \times_{h_B(T)} h_C(T) \times_{h_C(T)} h_D(T) \\ & \searrow & \downarrow \sim \\ & & h_A(T) \times_{h_B(T)} h_D(T) \end{array},$$

and natural transformations showing those functors are isomorphic, and hence represent isomorphic objects.