

I3 WITH RCOS QUESTIONS

- 1 (i) Express $\cos \theta + (\sqrt{3}) \sin \theta$ in the form $R \cos(\theta - \alpha)$, where $R > 0$ and $0 < \alpha < \frac{1}{2}\pi$, giving the exact values of R and α . [3]

(ii) Hence show that
$$\int_0^{\frac{1}{2}\pi} \frac{1}{(\cos \theta + (\sqrt{3}) \sin \theta)^2} d\theta = \frac{1}{\sqrt{3}}.$$
 [4]

9709/03/M/J/07

- 2 (i) Express $4 \cos \theta + 3 \sin \theta$ in the form $R \cos(\theta - \alpha)$, where $R > 0$ and $0 < \alpha < \frac{1}{2}\pi$. Give the value of α correct to 4 decimal places. [3]

(ii) Hence

(a) solve the equation $4 \cos \theta + 3 \sin \theta = 2$ for $0 < \theta < 2\pi$, [4]

(b) find
$$\int \frac{50}{(4 \cos \theta + 3 \sin \theta)^2} d\theta.$$
 [3]

9709/31/M/J/13

- 3 (i) Express $(\sqrt{3}) \cos x + \sin x$ in the form $R \cos(x - \alpha)$, where $R > 0$ and $0 < \alpha < \frac{1}{2}\pi$, giving the exact values of R and α . [3]

(ii) Hence show that

$$\int_{\frac{1}{6}\pi}^{\frac{1}{2}\pi} \frac{1}{((\sqrt{3}) \cos x + \sin x)^2} dx = \frac{1}{4}\sqrt{3}. \quad [4]$$

9709/33/M/J/13

4 **Throughout this question the use of a calculator is not permitted.**

- (i) Express $\cos \theta + 2 \sin \theta$ in the form $R \cos(\theta - \alpha)$, where $R > 0$ and $0 < \alpha < \frac{1}{2}\pi$. Give the exact values of R and $\tan \alpha$. [3]

(ii) Hence, showing all necessary working, show that
$$\int_0^{\frac{1}{4}\pi} \frac{15}{(\cos \theta + 2 \sin \theta)^2} d\theta = 5.$$
 [5]

9709/33/M/J/18