

I4 WITH TRIG IDENTITY QUESTIONS

- 1 (i) Prove the identity

$$\sin^2 \theta \cos^2 \theta \equiv \frac{1}{8}(1 - \cos 4\theta). \quad [3]$$

- (ii) Hence find the exact value of

$$\int_0^{\frac{1}{3}\pi} \sin^2 \theta \cos^2 \theta \, d\theta. \quad [3]$$

9709/03/M/J/04

- 2 (i) Using the expansions of $\cos(3x - x)$ and $\cos(3x + x)$, prove that

$$\frac{1}{2}(\cos 2x - \cos 4x) \equiv \sin 3x \sin x. \quad [3]$$

- (ii) Hence show that

$$\int_{\frac{1}{6}\pi}^{\frac{1}{3}\pi} \sin 3x \sin x \, dx = \frac{1}{8}\sqrt{3}. \quad [3]$$

9709/31/M/J/10

- 3 (i) Prove the identity $\cos 4\theta + 4 \cos 2\theta \equiv 8 \cos^4 \theta - 3$.

[4]

- (ii) Hence

(a) solve the equation $\cos 4\theta + 4 \cos 2\theta = 1$ for $-\frac{1}{2}\pi \leq \theta \leq \frac{1}{2}\pi$, [3]

(b) find the exact value of $\int_0^{\frac{1}{4}\pi} \cos^4 \theta \, d\theta$. [3]

9709/31/M/J/11

- 4 (i) By differentiating $\frac{1}{\cos x}$, show that if $y = \sec x$ then $\frac{dy}{dx} = \sec x \tan x$. [2]

(ii) Show that $\frac{1}{\sec x - \tan x} \equiv \sec x + \tan x$. [1]

(iii) Deduce that $\frac{1}{(\sec x - \tan x)^2} \equiv 2 \sec^2 x - 1 + 2 \sec x \tan x$. [2]

(iv) Hence show that $\int_0^{\frac{1}{4}\pi} \frac{1}{(\sec x - \tan x)^2} \, dx = \frac{1}{4}(8\sqrt{2} - \pi)$. [3]

9709/31/O/N/12

5 (i) Prove that $\cot \theta + \tan \theta \equiv 2 \operatorname{cosec} 2\theta$. [3]

(ii) Hence show that $\int_{\frac{1}{6}\pi}^{\frac{1}{3}\pi} \operatorname{cosec} 2\theta \, d\theta = \frac{1}{2} \ln 3$. [4]

9709/31/O/N/13

6 (i) Prove the identity $\tan 2\theta - \tan \theta \equiv \tan \theta \sec 2\theta$. [4]

(ii) Hence show that $\int_0^{\frac{1}{6}\pi} \tan \theta \sec 2\theta \, d\theta = \frac{1}{2} \ln \frac{3}{2}$. [4]

9709/31/O/N/16

7 (i) Prove that if $y = \frac{1}{\cos \theta}$ then $\frac{dy}{d\theta} = \sec \theta \tan \theta$. [2]

(ii) Prove the identity $\frac{1 + \sin \theta}{1 - \sin \theta} \equiv 2 \sec^2 \theta + 2 \sec \theta \tan \theta - 1$. [3]

(iii) Hence find the exact value of $\int_0^{\frac{1}{4}\pi} \frac{1 + \sin \theta}{1 - \sin \theta} \, d\theta$. [4]

9709/32/M/J/17

8 (i) Show that $\frac{2 \sin x - \sin 2x}{1 - \cos 2x} \equiv \frac{\sin x}{1 + \cos x}$. [4]


(ii) Hence, showing all necessary working, find $\int_{\frac{1}{3}\pi}^{\frac{1}{2}\pi} \frac{2 \sin x - \sin 2x}{1 - \cos 2x} \, dx$, giving your answer in the form $\ln k$. [4]

9709/32/M/J/18

9 (i) By first expanding $\sin(2x + x)$, show that $\sin 3x \equiv 3 \sin x - 4 \sin^3 x$. [4]

(ii) Hence, showing all necessary working, find the exact value of $\int_0^{\frac{1}{3}\pi} \sin^3 x \, dx$. [4]

9709/31/M/J/19



The diagram shows the curve $y = \sin 3x \cos x$ for $0 \leq x \leq \frac{1}{2}\pi$ and its minimum point M . The shaded region R is bounded by the curve and the x -axis.

- (i) By expanding $\sin(3x + x)$ and $\sin(3x - x)$ show that

$$\sin 3x \cos x = \frac{1}{2}(\sin 4x + \sin 2x). \quad [3]$$

- (ii) Using the result of part (i) and showing all necessary working, find the exact area of the region R . [4]

- (iii) Using the result of part (i), express $\frac{dy}{dx}$ in terms of $\cos 2x$ and hence find the x -coordinate of M , giving your answer correct to 2 decimal places. [5]

9709/32/M/J/19

11 Let $f(\theta) = \frac{1 - \cos 2\theta + \sin 2\theta}{1 + \cos 2\theta + \sin 2\theta}$.

- (i) Show that $f(\theta) = \tan \theta$. [3]

- (ii) Hence show that $\int_{\frac{1}{6}\pi}^{\frac{1}{4}\pi} f(\theta) d\theta = \frac{1}{2} \ln \frac{3}{2}$. [4]

9709/33/M/J/19

- 12 (i) By first expanding $\cos(2x + x)$, show that $\cos 3x \equiv 4 \cos^3 x - 3 \cos x$. [4]

- (ii) Hence solve the equation $\cos 3x + 3 \cos x + 1 = 0$, for $0 \leq x \leq \pi$. [2]

- (iii) Find the exact value of $\int_{\frac{1}{6}\pi}^{\frac{1}{3}\pi} \cos^3 x dx$. [4]

9709/31/O/N/19