

No. of Continuation Sheets attached _____

Sheet No. _____



PUNJAB COLLEGES

Name _____

Roll No. _____

Class _____

Section (if any) _____

Subject _____

Date _____

Student's Signature: _____

Invigilator's Name: _____

Signature: _____

(To be filled by the Examiner)

AWARD LIST

Q. No.	01 (MCQs)	02	03	04	05	06	07	08	09	10	Total Marks
Marks Obtained											

Examiner's Name: _____

Signature: _____

Please start writing from here

* Fourier Series of $f(x)$ over $(-P, P)$ is

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \left(a_n \cos \frac{n\pi x}{P} + b_n \sin \frac{n\pi x}{P} \right)$$

where $a_0 = \frac{1}{P} \int_{-P}^P f(x) dx$

$$a_n = \frac{1}{P} \int_{-P}^P f(x) \cos \frac{n\pi x}{P} dx$$

$$b_n = \frac{1}{P} \int_{-P}^P f(x) \sin \frac{n\pi x}{P} dx$$

* Ex. Expand $f(x) = \begin{cases} 0 & -\pi < x < 0 \\ \pi - x & 0 \leq x < \pi \end{cases}$

in a Fourier Series.

$$\begin{aligned} \text{Sol: } a_0 &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) dx = \frac{1}{\pi} \int_{-\pi}^0 0 dx + \int_0^{\pi} (\pi - x) dx \\ &= \frac{1}{\pi} \left(\pi x - \frac{x^2}{2} \right) \Big|_0^{\pi} = \pi/2 \end{aligned}$$

$$a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cdot \cos nx \, dx$$

$$= \frac{1}{\pi} \int_{-\pi}^0 0 \, dx + \frac{1}{\pi} \int_0^{\pi} (\pi - x) \cos nx \, dx$$

$$= \frac{1}{\pi} \left[(\pi - x) \cdot \frac{\sin nx}{n} \Big|_0^{\pi} - \frac{1}{n} \int_0^{\pi} \sin nx \, dx \right]$$

$$= -\frac{1}{n\pi} \frac{\cos nx}{n} \Big|_0^{\pi} = -\frac{1}{n\pi} \left[\frac{\cos n\pi}{n} - \frac{\cos 0}{n} \right]$$

$$= -\frac{1}{n\pi} \left[\frac{(-1)^n}{n} - \frac{1}{n} \right]$$

$$= +\frac{1}{n^2\pi} (1 - (-1)^n) = \frac{1 - (-1)^n}{n^2\pi}$$

$$b_n = \frac{1}{\pi} \int_{-\pi}^{\pi} (\pi - x) \sin nx \, dx = \frac{1}{\pi} \int_{-\pi}^0 0 \, dx + \int_0^{\pi} (\pi - x) \sin nx \, dx$$

$$= \frac{1}{\pi} \left[(\pi - x) \left(-\frac{\cos nx}{n} \right) \Big|_0^{\pi} + \int_0^{\pi} \frac{\cos nx}{n} (-1) \, dx \right]$$

$$= \frac{1}{\pi} \left[(0 + \frac{\pi}{n}) + \frac{1}{n} \int_0^{\pi} \cos nx \, dx \right]$$

$$= \frac{1}{\pi} - \frac{1}{n} \left| \frac{\sin nx}{n} \right|_0^{\pi} = \frac{1}{n} - 0 = \frac{1}{n}$$

$$\text{Hence } f(x) = \frac{\pi}{4} + \sum_{n=1}^{\infty} \left\{ \frac{1 - (-1)^n}{n^2\pi} \cos nx + \frac{1}{n} \sin nx \right\}$$

* Fourier Cosine Series

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \frac{a_n \cos \frac{n\pi}{p} x}{p}$$

$$a_0 = \frac{2}{p} \int_0^p f(x) dx$$

$$a_n = \frac{2}{p} \int_0^p f(x) \cos \frac{n\pi}{p} x dx$$

* Fourier Sine Series

$$f(x) = \sum_{n=1}^{\infty} \frac{b_n \sin \frac{n\pi}{p} x}{p}$$

$$b_n = \frac{2}{p} \int_0^p f(x) \cdot \sin \frac{n\pi}{p} x dx$$

Ex. $f(x) = x$; $-2 < x < 2$

Expand the above odd function in a Fourier Series.

$$b_n = \int_0^2 x \cdot \sin \frac{n\pi}{2} x dx = \frac{4(-1)^{n+1}}{n\pi}$$

$$\text{Hence } f(x) = \frac{4}{\pi} \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n} \sin \frac{n\pi}{2} x$$

Ex. $f(x) = x^2$; $0 < x < L$

Expand in a Cosine Series.

$$a_0 = \frac{2}{L} \int_0^L x^2 dx = \frac{2}{L} \left[\frac{x^3}{3} \right]_0^L$$

$$= \frac{2}{L} \left(\frac{L^3}{3} \right) = \frac{2L^2}{3}$$

$$a_n = \frac{2}{L} \int_0^L x^2 \cos \frac{n\pi x}{L} dx = \frac{4L^2 (-1)^n}{n^2 \pi^2}$$

Hence $f(x) = \frac{L^2}{3} + \frac{4L^2}{\pi^2} \sum_{n=1}^{\infty} \frac{(-1)^n}{n^2} \cos \frac{n\pi x}{L}$