

Ex. 1 Find a Parameterization of the cone

$$z = \sqrt{x^2 + y^2} ; 0 \leq z \leq 1$$

Let $x = r \cos \theta$ $y = r \sin \theta$ and $z = \sqrt{x^2 + y^2} = r$

where $0 \leq r \leq 1$ & $0 \leq \theta \leq 2\pi$

we have the parameterization

$$\vec{r}(r, \theta) = (r \cos \theta) \underline{i} + (r \sin \theta) \underline{j} + r \underline{k}$$

with $0 \leq r \leq 1$; $0 \leq \theta \leq 2\pi$

$$\vec{r}_r = \frac{\partial \vec{r}}{\partial r} = \cos \theta \underline{i} + \sin \theta \underline{j} + \underline{k}$$

$$\vec{r}_\theta = \frac{\partial \vec{r}}{\partial \theta} = -r \sin \theta \underline{i} + r \cos \theta \underline{j} + \underline{k}$$

$$\vec{r}_r \times \vec{r}_\theta = \begin{vmatrix} \underline{i} & \underline{j} & \underline{k} \\ \cos \theta & \sin \theta & 1 \\ -r \sin \theta & r \cos \theta & 0 \end{vmatrix}$$

$$\vec{r}_r \times \vec{r}_\theta = -r \cos \theta \underline{i} - r \sin \theta \underline{j} + r \underline{k}$$

$$|\vec{r}_r \times \vec{r}_\theta| = \sqrt{r^2 \cos^2 \theta + r^2 \sin^2 \theta + r^2} = \sqrt{2r^2} = \sqrt{2} r$$

$$\text{Surface Area} = \int_0^{2\pi} \int_0^1 \sqrt{2} r \, dr \, d\theta = \int_0^{2\pi} \frac{\sqrt{2}}{2} d\theta = \frac{\sqrt{2}}{2} \theta \Big|_0^{2\pi}$$

$$= \sqrt{2} \pi \text{ (unit)}^2$$

Ex. 2 Find a Parameterization and Surface
area of the sphere $x^2 + y^2 + z^2 = a^2$

We consider

$$x = a \sin \phi \cos \theta \quad y = a \sin \phi \sin \theta \quad z = a \cos \phi$$

$$0 \leq \phi \leq \pi \quad ; \quad 0 \leq \theta \leq 2\pi$$

$$\vec{r}(\phi, \theta) = (a \sin \phi \cos \theta) \underline{i} + (a \sin \phi \sin \theta) \underline{j} + a \cos \phi \underline{k}$$

$$\text{with } 0 \leq \phi \leq \pi \quad ; \quad 0 \leq \theta \leq 2\pi$$

$$\vec{r}_\phi \times \vec{r}_\theta = \begin{vmatrix} \underline{i} & \underline{j} & \underline{k} \\ a \cos \phi \cos \theta & a \cos \phi \sin \theta & -a \sin \phi \\ -a \sin \phi \sin \theta & a \sin \phi \cos \theta & 0 \end{vmatrix}$$

$$\vec{r}_\phi \times \vec{r}_\theta = (a^2 \sin^2 \phi \cos \theta) \underline{i} + (a^2 \sin^2 \phi \sin \theta) \underline{j} + a^2 \sin \phi \cos \phi \underline{k}$$

$$|\vec{r}_\phi \times \vec{r}_\theta| = a^2 \sin \phi$$

$$A = \int_0^{2\pi} \int_0^\pi a^2 \sin \phi \, d\phi \, d\theta$$

$$= \int_0^{2\pi} [-a^2 \cos \phi]_0^\pi \, d\theta = \int_0^{2\pi} 2a^2 \, d\theta = 4\pi a^2 \text{ (unit)}^2$$

Ex-3 Find S.A from the bottom of Paraboloid
 $x^2 + y^2 - z = 0$ by the Plane $z=4$

Here we can choose

$$x^2 + y^2 \leq 4 \quad \& \quad \vec{p} = \underline{k}$$

$$\& \quad F(x, y, z) = x^2 + y^2 - z$$

$$\nabla F = 2x\underline{i} + 2y\underline{j} - \underline{k}$$

$$|\nabla F| = \sqrt{4x^2 + 4y^2 + 1}$$

$$|\nabla F \cdot \underline{p}| = |1 - 1| = \sqrt{1} = 1$$

$$S.A = \int_0^{2\pi} \int_0^2 (4r^2 + 1)^{1/2} r dr d\theta$$

$$= \int_0^{2\pi} \left[\frac{1}{12} (4r^2 + 1)^{3/2} \right]_0^2 d\theta$$

$$= \frac{\pi}{6} (17\sqrt{17} - 1) (\text{unit})^2$$

Ex. 4 $F(x,y,z) = x^2$ over the cone $z = \sqrt{x^2 + y^2}$;

$$0 \leq z \leq 1$$

$$\iint_S x^2 d\sigma = \int_0^{2\pi} \int_0^1 (h^2 \cos^2 \theta) \sqrt{2} h \, dh \, d\theta$$

from Ex 1.

Because

$$(1) \quad \iint_S F(x,y,z) d\sigma = \iint_R F(f(u,v), g(u,v), h(u,v)) |\vec{r}_u \times \vec{r}_v| \, du \, dv$$

$$= \sqrt{2} \int_0^{2\pi} \int_0^1 h^3 \cos^2 \theta \, dh \, d\theta$$

$$= \frac{\sqrt{2}}{4} \int_0^{2\pi} \cos^2 \theta \, d\theta = \frac{\sqrt{2}}{4} \left[\frac{\theta}{2} + \frac{1}{4} \sin 2\theta \right]_0^{2\pi} = \frac{\pi \sqrt{2}}{4}$$