

8.4 Trigonometric Substitutions

①


Date: / /

→ Trigonometric Substitutions occur when we replace the variable of integration by a trigonometric function.

→ The most common substitutions are:


① $x = a \tan \theta$

$$\sqrt{a^2 + x^2} = a |\sec \theta|$$




② $x = a \sin \theta$

$$\sqrt{a^2 - x^2} = a |\cos \theta|$$



③ $x = a \sec \theta$

$$\sqrt{x^2 - a^2} = a |\tan \theta|$$



→ These substitutions are effective in transforming integrals involving $\sqrt{a^2 + x^2}$, $\sqrt{a^2 - x^2}$, $\sqrt{x^2 - a^2}$ into integrals we can evaluate directly

Trigonometric
function

$$1 + \tan^2 \theta = \sec^2 \theta$$

$$1 - \sin^2 \theta = \cos^2 \theta$$

$$\sec^2 \theta - 1 = \tan^2 \theta$$

(2)

Date: / /

* Read details from book (Pg 467)

$$x = a \tan \theta \Rightarrow \theta = \tan^{-1}\left(\frac{x}{a}\right) \text{ with } -\frac{\pi}{2} < \theta < \frac{\pi}{2}$$

$$x = a \sin \theta \Rightarrow \theta = \sin^{-1}\left(\frac{x}{a}\right) \text{ with } -\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$$

$$x = a \sec \theta \Rightarrow \theta = \sec^{-1}\left(\frac{x}{a}\right) \text{ with } \begin{cases} 0 \leq \theta < \frac{\pi}{2} & \text{if } \frac{x}{a} \geq 1 \\ \frac{\pi}{2} < \theta \leq \pi & \text{if } \frac{x}{a} \leq -1 \end{cases}$$


Procedure for a Trigonometric Substitution

- ① Write down the substitution for x , calculate dx and specify the selected values of θ for substitution.
- ② Substitute the trigonometric expression and the calculated differential into the integrand and then simplify the results algebraically.
- ③ Integrate the trigonometric integral, keeping in mind the restrictions on the angle θ for reversibility.
- ④ Draw an appropriate reference triangle to reverse the substitution in the integration result and convert it back to the original variable x .

Date: / /

* Example 1: $\int \frac{dx}{\sqrt{4+x^2}} = \int \frac{dx}{\sqrt{2^2+x^2}}$

Sol: Let $x = 2 \tan \theta$
 $x = 2 \tan \theta$



$$dx = 2 \sec^2 \theta d\theta, \quad -\frac{\pi}{2} < \theta < \frac{\pi}{2}$$

$$4+x^2 = 4 + 4 \tan^2 \theta$$

$$= 4(1 + \tan^2 \theta) = 4 \sec^2 \theta$$

$$\int \frac{dx}{\sqrt{4+x^2}} = \int \frac{dx}{\sqrt{2^2+x^2}} = \int \frac{2 \sec^2 \theta d\theta}{\sqrt{4 \sec^2 \theta}}$$

$$= \int \frac{2 \sec^2 \theta d\theta}{2 \sec \theta}$$

$$= \int \sec \theta d\theta$$

$$= \ln |\sec \theta + \tan \theta| + C$$

$$= \ln \left| \frac{\sqrt{4+x^2}}{2} + \frac{x}{2} \right| + C$$

R.W

$$\bullet x = 2 \tan \theta$$

$$\tan \theta = \frac{x}{2}$$

$$\bullet 4+x^2 = 4 \sec^2 \theta$$

$$\sec^2 \theta = \frac{4+x^2}{4}$$


$$\sec \theta = \frac{\sqrt{4+x^2}}{2}$$

* Example 3: Evaluate $\int \frac{x^2}{\sqrt{9-x^2}} dx = \int \frac{x^2}{\sqrt{3^2-x^2}} dx$

Sol: Let $x = 3 \sin \theta$

$$\theta = \sin^{-1} \left(\frac{x}{3} \right) \Leftrightarrow x = 3 \sin \theta$$

$$\frac{x}{3} = \sin \theta \Leftrightarrow dx = 3 \cos \theta d\theta, \quad -\frac{\pi}{2} < \theta < \frac{\pi}{2}$$



$$9-x^2 = 9 - (3 \sin \theta)^2$$

$$= 9 - 9 \sin^2 \theta$$

$$9-x^2 = 9 \cos^2 \theta$$

$$\cos^2 \theta = \frac{9-x^2}{9}$$

$$\cos \theta = \frac{\sqrt{9-x^2}}{3}$$

$$= 9(1 - \sin^2 \theta) = 9 \cos^2 \theta$$

(4)

Date: / /

$$\int \frac{x^2}{\sqrt{9-x^2}} dx = \int \frac{9 \sin^2 \theta \cdot 3 \cos \theta d\theta}{\sqrt{9 \cos^2 \theta}}$$

$$= 9 \int \frac{3 \sin^2 \theta \cos \theta d\theta}{3 \cos \theta}$$

$$= 9 \int \sin^2 \theta d\theta \quad \left(\because \sin^2 \theta = \frac{1 - \cos 2\theta}{2} \right)$$

$$= 9 \int \left(\frac{1 - \cos 2\theta}{2} \right) d\theta$$

$$= \frac{9}{2} \left[\theta - \frac{\sin 2\theta}{2} \right] + C$$

$$= \frac{9}{2} \left(\theta - \frac{2 \cos \theta \sin \theta}{2} \right) + C \quad \left(\because \sin 2\theta = 2 \cos \theta \sin \theta \right)$$

$$= \frac{9}{2} (\theta - \sin \theta \cos \theta) + C$$

$$= \frac{9}{2} \left(\frac{\sin^{-1} x}{3} - \frac{x}{3} \cdot \frac{\sqrt{9-x^2}}{3} \right) + C$$

$$= \frac{9}{2} \sin^{-1} \frac{x}{3} - \frac{3}{2} x \sqrt{9-x^2} + C$$

$\xleftrightarrow{\hspace{1.5cm}}$

(5)

Date: / /

* Example 4: $\int \frac{dx}{\sqrt{25x^2 - 4}}$ $x > \frac{2}{5}$

Sol: $\sqrt{25x^2 - 4} = \sqrt{25\left(x^2 - \frac{4}{25}\right)}$

$$= \sqrt{25\left(x^2 - \left(\frac{2}{5}\right)^2\right)}$$

$$= \sqrt{25} \sqrt{x^2 - \left(\frac{2}{5}\right)^2} = 5 \sqrt{x^2 - \left(\frac{2}{5}\right)^2}$$

$$x = a \sec \theta \Rightarrow x = \frac{2}{5} \sec \theta$$

$$0 < \theta < \frac{\pi}{2}$$

$$dx = \frac{2}{5} \sec \theta \tan \theta d\theta$$

$$x^2 - \left(\frac{2}{5}\right)^2 = \left(\frac{2}{5} \sec \theta\right)^2 - \left(\frac{2}{5}\right)^2$$

$$= \frac{4}{25} \sec^2 \theta - \frac{4}{25} = \frac{4}{25} (\sec^2 \theta - 1)$$

$$= \frac{4}{25} \tan^2 \theta$$

$$\int \frac{dx}{\sqrt{25x^2 - 4}} = \int \frac{dx}{5 \sqrt{x^2 - \left(\frac{2}{5}\right)^2}} = \int \frac{\frac{2}{5} \sec \theta \tan \theta d\theta}{5 \cdot \sqrt{\frac{4}{25} \tan^2 \theta}}$$

$$= \int \frac{\frac{2}{5} \sec \theta \tan \theta d\theta}{5 \cdot \frac{2}{5} \tan \theta}$$

$$= \frac{1}{5} \int \sec \theta d\theta = \frac{1}{5} \ln |\sec \theta + \tan \theta| + C$$

$$= \frac{1}{5} \ln \left| \frac{5x}{2} + \frac{\sqrt{25x^2 - 4}}{2} \right| + C$$

← →