



# Lecture 27

---

## Linear Regression







Slides created by John DeNero ([denero@berkeley.edu](mailto:denero@berkeley.edu)) and Ani Adhikari ([adhikari@berkeley.edu](mailto:adhikari@berkeley.edu))  
Contributions by Fahad Kamran ([fhdkmrn@berkeley.edu](mailto:fhdkmrn@berkeley.edu)) and Vinitra Swamy ([vinitra@berkeley.edu](mailto:vinitra@berkeley.edu))

# **Announcements**

# Correlation (Review)

# The Correlation Coefficient $r$

- Measures **linear** association
- Based on standard units
- $-1 \leq r \leq 1$ 
  - $r = 1$ : scatter is perfect straight line sloping up
  - $r = -1$ : scatter is perfect straight line sloping down
- $r = 0$ : No linear association; *uncorrelated*



# Definition of $r$

---

**Correlation Coefficient ( $r$ ) =**

average of	product of	x in standard units	and	y in standard units
---------------	------------	---------------------------	-----	---------------------------


Measures how clustered the scatter is around a straight line

---

**Prediction**


# Galton's Heights

---



# Galton's Heights


---






# Galton's Heights

---



# Where is the prediction line?

---




$$r = 0.99$$

---

# Where is the prediction line?

---




$$r = 0.0$$

---

# Where is the prediction line?

---




$r = 0.5$

---

# Where is the prediction line?

---



$r = 0.2$

---

# Nearest Neighbor Regression

---

A method for prediction:

- Group each  $x$  with a representative  $x$  value (rounding)
- Average the corresponding  $y$  values for each group

For each representative  $x$  value, the corresponding prediction is the average of the  $y$  values in the group.

Graph these predictions.

If the association between  $x$  and  $y$  is linear, then points in the graph of averages tend to fall on the regression line.

---

# Linear Regression

(Demo)

# Regression to the Mean

---

A statement about  $x$  and  $y$  pairs

- Measured in *standard units*
- Describing the deviation of  $x$  from 0 (the average of  $x$ 's)
- And the deviation of  $y$  from 0 (the average of  $y$ 's)

*On average*,  $y$  deviates from 0 less than  $x$  deviates from 0

Regression  
Line

$$y_{(\text{su})} = r \times x_{(\text{su})}$$

Correlation

Not true for all points — a statement about averages

---



# Slope & Intercept

# Regression Line Equation

---

In original units, the regression line has this equation:

$$\frac{\text{estimate of } y - \text{average of } y}{\text{SD of } y} = r \times \frac{\text{the given } x - \text{average of } x}{\text{SD of } x}$$

estimated y in standard units

x in standard units


Lines can be expressed by *slope* & *intercept*

$$y = \text{slope} \times x + \text{intercept}$$


---

# Regression Line

## Standard Units



## Original Units



# Slope and Intercept

---

estimate of  $y$  = slope \*  $x$  + intercept

$$\text{slope of the regression line} = r \cdot \frac{\text{SD of } y}{\text{SD of } x}$$

$$\text{intercept of the regression line} = \text{average of } y - \text{slope} \cdot \text{average of } x$$

(Demo)

---

# Scenario Question

---

You use a regression line to predict height based on weight and get a slope of .52 inches per pound.

I eat a lot of ice cream and gain 1 pound.

**True or False:**

My regression line predicts I will gain 1 inch on my height

---

# Scenario Question

---

## False

The regression line is a **statement about averages**. Given two groups of people with 1 lb difference, we expect the average height of the heavier group to be .52 inches greater than the average height of the lighter group

The regression line is based on a snapshot of time and looking at holistic trends. We are **not** following 1 person and intending to make a prediction about them

---