

Modeling and Inference for Rotavirus Dynamics in Niger

(joint work with Jaewoo Park, Josh Goldstein, Matt Ferrari)

ISBA 2018 Meeting, Edinburgh, Scotland.
June 2018.

Murali Haran

Department of Statistics, Penn State University

Outline

Rotavirus

Rotavirus

- ▶ Rotavirus infection causes significant morbidity, mortality worldwide
- ▶ In sub-Saharan Africa, cause of diarrheal disease often unknown due to lack of infrastructure
- ▶ Uncertainty about the underlying transmission dynamics
- ▶ Our Goal:
 - ▶ Estimate burden of rotavirus in the Maradi region of Niger
 - ▶ Predict impact of vaccination on rotavirus burden
- ▶ Methods:
 - ▶ Compare several compartmental SIR type models.
 - ▶ Account for model uncertainty via Bayesian model averaging (BMA).

Two Sources of Rotavirus Data

- ▶ Surveillance data collected over two years (12/2009 - 4/2011) from hospitals and health centers
 - ▶ 9,600 children ≤ 5 years presenting with severe diarrhea
 - ▶ For each child: Dates of consultation, results of a rapid rotavirus diagnostic test (30% test positive)
 - ▶ A subset (378) of cases are also genotyped
- ▶ Cluster survey of households with children ≤ 5
 - ▶ Hospital surveillance data can miss a majority of cases
 - ▶ Use cluster survey to estimate reporting rates (42.9% sought care at a hospital or health center).

Vaccination


- ▶ Two dose strategy: 2 months, 4 months of age
- ▶ Assume vaccination confers same level of protection as an infection.
- ▶ Both short-term (5 years) and long-term (20 years) effects are investigated.

SIR Compartmental Models


- ▶ We consider 5 different SIR models (Pitzer et al., 2012)
- ▶ Notation:
 - ▶ (M) the number of individuals in maternal immunity
 - ▶ (S) the number of susceptible individuals
 - ▶ (E) the number of individuals in incubation period
 - ▶ (I) the number of infected individuals
 - ▶ (R) the number of recovered individuals

Model Examples

B



C




Model Examples

- ▶ Models differ by characteristics such as:
 - ▶ Whether they track severe and mild cases separately
 - ▶ Whether they allow for successive infections
 - ▶ Whether they allow for incubation period
 - ▶ How immunity is granted


Inference and Model Details

- ▶ Bayesian inference via Markov chain Monte Carlo
 - ▶ Periodic solution for the disease dynamics is required for likelihood evaluation
- ▶ Dynamics Accounting for Vaccination
 - ▶ Given estimated model parameters, dynamic models integrated forward to steady state solution.
 - ▶ Then, the dynamics are modified to account for vaccination.
- ▶ BMA (Bates and Granger, 1969; Hoeting et al., 1999)
 - ▶ Posterior model probability (PMP) measures how well each model is supported.
 - ▶ Combining information across models

Distribution of Cases Across Age Groups



Model-averaged predicted vaccine impact



- Over the short term (Left), models predict an overall decline in total burden, but an increase in the magnitude of peak incidence.
- For a fixed (70%) level of coverage, we predict 38.9% reduction of severe rotavirus (99%CI : (37.1%; 42.6%))

over the long-term (Middle)


Pre-vaccination

Model Comparison Summary

Model	Probability	R_0	Burden
A	0	30.7 (25.8,34.3)	9.2 (8.1,10.1)
B	0.01	13.9 (12.7,15.4)	3.5 (3.1,3.8)
C	0.92	13.4 (11.7,15.3)	3.5 (3.1,3.9)
D	0.03	11.2 (9.4,12.7)	3.6 (3.2,4.1)
E	0.04	10.3 (9.5,12.6)	3.2 (2.9,3.5)
BMA		13.4 (10.3,15.4)	3.5 (2.9,4.2)

- ▶ Model C performs the best, Model A performs significantly worse.
- ▶ Estimates of burden and basic reproductive ratio (R_0) are significantly larger for Model A.

Model-averaged (BMA) burden estimates



- Models successfully capture the observed dynamics (dots). Seasonal effects explain observed double incidence peak.
- Models predict a steep decline in cases in children under 1y of age following the epidemic peak.

Conclusions

- ▶ We develop flexible models that allow for a variety of different transmission dynamics for rotavirus
- ▶ BMA approach can account for uncertainty in the underlying dynamics.
- ▶ We can predict short-term/long-term reduction in rotavirus due to vaccination.
- ▶ Evidence that human activity is the primary vector of transmission.
 - ▶ The peak transmission estimated with a maximum in early March.
 - ▶ Urban population density is at its maximum due to seasonal rural-urban migration (Bharti et al., 2011)

Acknowledgement

The authors are grateful to Epicentre for providing the data sets for this research project. MF is funded by a grant from the Ecology and Evolution of Infectious Disease program of the NSF/NIH (award number 1 R01 GM105247-01).

None of the authors has conflicts of interest.