

# Sequential Monte Carlo Estimation of High Dimensional Latent Variable Models

Application to Stochastic Volatility Models

---

Han Xiao

Nov. 2018

Pennsylvania State University

# Motivation

- **Estimate** high-dimensional latent variable models

Stochastic volatility models: Widely used in practice and option pricing. Application Given asset returns, study the underlying volatility.

# Motivation

- **Estimate** high-dimensional latent variable models

Stochastic volatility models: Widely used in practice and option pricing. Application Given asset returns, study the underlying volatility.

$$y_t = \exp\left(\frac{h_t}{2}\right) \varepsilon_t \quad (1)$$

$$h_{t+1} = \mu + \phi(h_t - \mu) + \sigma\eta_t \quad (2)$$

where  $\varepsilon$  and  $\eta$  are iid standard normal distribution.

- $y_t$ : asset return in reality  $\rightarrow$  Observable
- $h_t$ : risk underlying the asset  $\rightarrow$  Latent variable
- Parameter:  $\Theta = \{\mu, \phi, \sigma^2\}$

# Motivation

- **Estimate** high-dimensional latent variable models

Stochastic volatility models: Widely used in practice and option pricing. Application Given asset returns, study the underlying volatility.

$$y_t = \exp\left(\frac{h_t}{2}\right) \varepsilon_t \quad (1)$$

$$h_{t+1} = \mu + \phi(h_t - \mu) + \sigma\eta_t \quad (2)$$

where  $\varepsilon$  and  $\eta$  are iid standard normal distribution.

- $y_t$  : asset return in reality  $\rightarrow$  Observable
- $h_t$ : risk underlying the asset  $\rightarrow$  Latent variable
- Parameter:  $\Theta = \{\mu, \phi, \sigma^2\}$
- **Goal:** learn parameters and latent variables recursively
  - use real time data and estimate efficiently

# Computational challenge

The difficult part occurs in the latent variable (Equation (2)), i.e., likelihood function is not easily available:

$$f(h_t|y_t, \Theta) \propto \underbrace{f(y_t|h_t, \Theta)}_{\text{observed return}} \underbrace{f(h_t|y_{t-1}, \Theta)}_{\text{latent variable: volatility}} \quad (3)$$

# Computational challenge

The difficult part occurs in the latent variable (Equation (2)), i.e., likelihood function is not easily available:

$$f(h_t|y_t, \Theta) \propto \underbrace{f(y_t|h_t, \Theta)}_{\text{observed return}} \underbrace{f(h_t|y_{t-1}, \Theta)}_{\text{latent variable: volatility}} \quad (3)$$

$$f(h_t|y_{t-1}, \Theta) = \int \underbrace{f(h_t|h_{t-1}, \Theta)}_{\text{proposed AR(1)}} \underbrace{f(h_{t-1}|y_{t-1}, \Theta)}_{\text{hard closed form}} dh_{t-1} \quad (4)$$

$\Rightarrow$  *Difficult to evaluate the objective function*

# Computational challenge

The difficult part occurs in the latent variable (Equation (2)), i.e., likelihood function is not easily available:

$$f(h_t|y_t, \Theta) \propto \underbrace{f(y_t|h_t, \Theta)}_{\text{observed return}} \underbrace{f(h_t|y_{t-1}, \Theta)}_{\text{latent variable: volatility}} \quad (3)$$

$$f(h_t|y_{t-1}, \Theta) = \int \underbrace{f(h_t|h_{t-1}, \Theta)}_{\text{proposed AR(1)}} \underbrace{f(h_{t-1}|y_{t-1}, \Theta)}_{\text{hard closed form}} dh_{t-1} \quad (4)$$

$\Rightarrow$  *Difficult to evaluate the objective function*

**Challenge 1:** Cannot directly draw **sample** from  $f(h_t|y_{t-1}, \Theta)$

# Computational challenge

The difficult part occurs in the latent variable (Equation (2)), i.e., likelihood function is not easily available:

$$f(h_t|y_t, \Theta) \propto \underbrace{f(y_t|h_t, \Theta)}_{\text{observed return}} \underbrace{f(h_t|y_{t-1}, \Theta)}_{\text{latent variable: volatility}} \quad (3)$$

$$f(h_t|y_{t-1}, \Theta) = \int \underbrace{f(h_t|h_{t-1}, \Theta)}_{\text{proposed AR(1)}} \underbrace{f(h_{t-1}|y_{t-1}, \Theta)}_{\text{hard closed form}} dh_{t-1} \quad (4)$$


$\Rightarrow$  *Difficult to evaluate the objective function*

**Challenge 1:** Cannot directly draw **sample** from  $f(h_t|y_{t-1}, \Theta)$

**Challenge 2:** Computational **complexity** of simulation increases with  
the number of time points in data



# Intuition for sequential Monte Carlo Method



## Sequential Monte Carlo: particle filter algorithm

- Main Idea: draw particles  $\{h_{t-1}^j\}$  from filtered distribution  $f(h_{t-1}|y_{t-1})$ , and derive  $f(h_t|y_t)$  using Equation (2). Then sequentially draw particles  $\{h_t^j\}$

# Sequential Monte Carlo: particle filter algorithm

- Main Idea: draw particles  $\{h_{t-1}^j\}$  from filtered distribution  $f(h_{t-1}|y_{t-1})$ , and derive  $f(h_t|y_t)$  using Equation (2). Then sequentially draw particles  $\{h_t^j\}$
- Sequentially draw  $M$  particles at each time point  $t$   
 $\{h_{t-1}\} = \{h_{t-1}^1, \dots, h_{t-1}^M\}$  from the filtered distributions:  $f(h_{t-1}|y_{t-1})$ , where  $M$  is the number of particles, then get

$$h_t = \mu + \phi(h_{t-1}^j - \mu)$$

# Sequential Monte Carlo: particle filter algorithm

- Main Idea: draw particles  $\{h_{t-1}^j\}$  from filtered distribution  $f(h_{t-1}|y_{t-1})$ , and derive  $f(h_t|y_t)$  using Equation (2). Then sequentially draw particles  $\{h_t^j\}$
- Sequentially draw  $M$  particles at each time point  $t$   
 $\{h_{t-1}\} = \{h_{t-1}^1, \dots, h_{t-1}^M\}$  from the filtered distributions:  $f(h_{t-1}|y_{t-1})$ , where  $M$  is the number of particles, then get

$$h_t = \mu + \phi(h_{t-1}^j - \mu)$$

This gives Monte Carlo approximations

$$f(h_t|y_{t-1}, \Theta) \approx \frac{1}{M} \sum_{j=1}^M f(h_t|h_{t-1}^j, \Theta) \quad (5)$$

$$f(h_t|y_t, \Theta) \propto f(y_t|h_t, \Theta) \frac{1}{M} \sum_{j=1}^M f(h_t|h_{t-1}^j, \Theta) \quad (6)$$

# Sequential Monte Carlo: particle filter algorithm

- Main Idea: draw particles  $\{h_{t-1}^j\}$  from filtered distribution  $f(h_{t-1}|y_{t-1})$ , and derive  $f(h_t|y_t)$  using Equation (2). Then sequentially draw particles  $\{h_t^j\}$
- Sequentially draw  $M$  particles at each time point  $t$   
 $\{h_{t-1}\} = \{h_{t-1}^1, \dots, h_{t-1}^M\}$  from the filtered distributions:  $f(h_{t-1}|y_{t-1})$ , where  $M$  is the number of particles, then get

$$h_t = \mu + \phi(h_{t-1}^j - \mu)$$

This gives Monte Carlo approximations

$$f(h_t|y_{t-1}, \Theta) \approx \frac{1}{M} \sum_{j=1}^M f(h_t|h_{t-1}^j, \Theta) \quad (5)$$

$$f(h_t|y_t, \Theta) \propto f(y_t|h_t, \Theta) \frac{1}{M} \sum_{j=1}^M f(h_t|h_{t-1}^j, \Theta) \quad (6)$$

- I explore different sampling methods: importance-sampling, bootstrap, auxiliary variable

## MCMC Algorithm

1. Initialize  $h_1$  and  $\Theta^{(1)}$
2. Sample  $h_t$  from  $h_t^{(k+1)} | h_{t-1}^{(k)}, y, \Theta^{(k)}$  for  $t = 1, \dots, n$
3. Sample  $\sigma^{2(k+1)} | y, h^{(k+1)}, \phi^{(k)}, \mu^{(k)}$
4. Sample  $\phi^{(k+1)} | h^{(k+1)}, \mu^{(k)}, \sigma^{2(k+1)}$
5. Sample  $\mu^{(k+1)} | h^{(k+1)}, \phi^{(k+1)}, \sigma^{2(k+1)}$
6. Goto 2

## MCMC Algorithm

1. Initialize  $h_1$  and  $\Theta^{(1)}$
2. Sample  $h_t$  from  $h_t^{(k+1)} | h_{t-1}^{(k)}, y, \Theta^{(k)}$  for  $t = 1, \dots, n$
3. Sample  $\sigma^{2(k+1)} | y, h^{(k+1)}, \phi^{(k)}, \mu^{(k)}$
4. Sample  $\phi^{(k+1)} | h^{(k+1)}, \mu^{(k)}, \sigma^{2(k+1)}$
5. Sample  $\mu^{(k+1)} | h^{(k+1)}, \phi^{(k+1)}, \sigma^{2(k+1)}$
6. Goto 2

## Computing is more expensive

- Need to sample from  $h_t^{(k+1)} | h_{t-1}^{(k)}, y, \Theta^{(k)}$  at each time point
- $\{h_t\}, \Theta$  are highly correlated, resulting in slow convergence

ACF

# Comparison: Efficiency and Accuracy





- Simulated data:  $\mu_0 = 0.5$ ,  $\phi_0 = 0.985$ ,  $\sigma_0^2 = 0.04$
- Real data: Daily S&P 500 returns, 1950 – 2018

	Simulated Data				Real Data			
	MCMC	SISR	Boot	APF	MCMC	SISR	Boot	APF
Computational Efficiency								
ESS	5302	<b>724</b>	902	964	1107	<b>576</b>	756	826
time	113.106	<b>0.117</b>	0.228	0.311	193.733	<b>2.048</b>	3.766	4.274
MSE for last latent variable $h_N$								
mean	<b>0.013</b>	0.098	0.100	0.097	1.174	0.134	0.134	<b>0.132</b>
std	0.315	<b>0.041</b>	0.045	0.043	<b>0.047</b>	2.096	2.082	2.003
MSE for parameters								
$\mu$	0.200	0.010	<b>0.000</b>	0.090	—	—	—	—
$\phi$	0.006	<b>0.000</b>	0.000	0.000	—	—	—	—
$\sigma^2$	0.004	<b>0.001</b>	0.005	0.003	—	—	—	—



# Comparison: Latent Variable Estimates

Posterior:  $h_t, y_t$



# Conclusion

- SMC Strength
  - Efficient: large sample and iteration, more parameters
  - The algorithm can be easily parallelized
  - The computational complexity does not increase with increase in time
  - As the number of particles  $M$  increases, the accuracy increases.

# Conclusion

- SMC Strength
  - Efficient: large sample and iteration, more parameters
  - The algorithm can be easily parallelized
  - The computational complexity does not increase with increase in time
  - As the number of particles  $M$  increases, the accuracy increases.
- SMC Weakness
  - Sensitive to outliers
  - Poor approximation at tails
  - Inference is dominated by a few particles with high weight: Potential solution is auxiliary particle filter

# Conclusion

- SMC Strength
  - Efficient: large sample and iteration, more parameters
  - The algorithm can be easily parallelized
  - The computational complexity does not increase with increase in time
  - As the number of particles  $M$  increases, the accuracy increases.
- SMC Weakness
  - Sensitive to outliers
  - Poor approximation at tails
  - Inference is dominated by a few particles with high weight: Potential solution is auxiliary particle filter
- Auxiliary particle filter may be better at handling outliers and heavy tails
  - I found some tentative evidence from simulation

**Thank you**

# Appendix

- Application
- Auxiliary particle filter algorithm
- MCMC simulation autocorrelation
- Sequence density

# Application: Stochastic Volatility Models

1. Option price: Black-Scholes, Heston and Hull-White model
2. Long run risk model
3. Industry

[Back](#)

# Methodology: Sequential Monte Carlo

## SMC Algorithm: auxiliary particle filter [Back](#)

1. Given  $\{h_{t-1}^1, \dots, h_{t-1}^M\}$  from  $f(h_{t-1}|y_{t-1}, \Theta)$  calculate

$$\begin{aligned}\widehat{h}_t^{*j} &= \mu + \phi(h_{t-1}^j - \mu) \\ w_j &= f_N(y_t | \exp(\widehat{h}_t^{*j})), \quad j = 1, \dots, M\end{aligned}$$

and sample  $R$  times with probability  $\{w_j\}$ . Let the sampled index be  $k_1, \dots, k_R$ , and associate these with  $\widehat{h}_t^{*k_1}, \dots, \widehat{h}_t^{*k_R}$

2. For each value of  $k_j$  from Step 1 simulate

$$h_t^{*j} \sim \mathcal{N}(\mu + \phi(h_{t-1}^{*k_j} - \mu), \sigma^2), \quad j = 1, \dots, R$$




3. Resample  $\{h_t^{*1}, \dots, h_t^{*R}\}$  with probability

$$\frac{\mathcal{N}(\mu + \phi(h_{t-1}^{*j} - \mu), \sigma^2)}{\mathcal{N}(\mu + \phi(h_{t-1}^{*k_j} - \mu), \sigma^2)}$$



to produce the filtered sample  $\{h_t^1, \dots, h_t^M\}$  from  $f(h_t|y_t, \Theta)$



# Slow convergence in MCMC: Autocorrelation



# Density analysis: simulated and real data



Back

# Reference

- Gordon, N. J., Salmond, D. J., and Smith, A. F. M. (1993), Novel approach to nonlinear/non-Gaussian Bayesian state estimation, IEEE Proceedings Part F: Communications, Radar and Signal Processing, 140, 107–113.
- Kim, S., Shephard, N., and Chib, S. (1998). Stochastic volatility: likelihood inference and comparison with ARCH models. The review of economic studies, 65(3), 361-393.
- Pitt, M. K. and Shephard, N. (1999), Filtering via simulation: auxiliary particle filters, Journal of the American Statistical Association, 94, 590–599.
- Creal, D. (2012). A survey of sequential Monte Carlo methods for economics and finance. Econometric reviews, 31(3), 245-296.