Multi-Dimensional Array related problems (Total 15 questions)

SL	Problem statement	Difficulty
		levels

Sample input	Sample output
987654321	9 8 7
987034321	654
	321
111222333	111
111222333	222
	333
nd column-wise.	ers into a <i>m by n</i> array (2D) and print them both row-wise
Sample input (m,n)	Sample output
23	Row-wise: 1 2 3 6 5 4
123	Column-wise: 1 6 2 5 3 4
6 5 4 3 3	Pow wise: 1 1 1 2 2 2 2 2
111	Row-wise: 1 1 1 2 2 2 3 3 3 Column-wise: 1 2 3 1 2 3 1 2 3
222	Column-wise: 123123123
333	
VAP that will take inputs of a 3 his matrix. http://www.mathsisfu	by 3 matrix into a 2D array. Now find the determinant of un.com/algebra/matrix-determinant.html
VAP that will take inputs of a 3 his matrix. http://www.mathsisfu Sample input	un.com/algebra/matrix-determinant.html Sample output
VAP that will take inputs of a 3 his matrix. http://www.mathsisfusample input 123	un.com/algebra/matrix-determinant.html
VAP that will take inputs of a 3 his matrix. http://www.mathsisfu Sample input	un.com/algebra/matrix-determinant.html Sample output

elements of its two diagonals.		
Sample input	Sample output	
5	Major diagonal: 1 4 2 9 4	
12345	Minor diagonal: 5 2 2 7 1	
54321		
22222		
67890		
19374		
		*
	WAP that will take the size of an identity matrix from the user and generate the identity matrix into a 2D array. Finally display it. Reference: http://en.wikipedia.org/wiki/Identity_matrix	
Sample input	Sample output	
5	10000	
	01000	
	00100	
	00010	
•	0 0 0 0 1 yo m x n sized matrix into two 2D array, suppose A and B. ay all the elements from matrix / 2D array C.	*
Now do C = A + B. Finally displa	yo m x n sized matrix into two 2D array, suppose A and B. ay all the elements from matrix / 2D array C.	*
Now do C = A + B. Finally displa Sample input	yo m x n sized matrix into two 2D array, suppose A and B. ay all the elements from matrix / 2D array C. Sample output	*
Now do C = A + B. Finally displa Sample input 2 3	yo m x n sized matrix into two 2D array, suppose A and B. ay all the elements from matrix / 2D array C. Sample output 2 3 4	*
Now do C = A + B. Finally displa Sample input 2 3 1 2 3	yo m x n sized matrix into two 2D array, suppose A and B. ay all the elements from matrix / 2D array C. Sample output	*
Sample input 2 3 1 2 3 2 3 4	yo m x n sized matrix into two 2D array, suppose A and B. ay all the elements from matrix / 2D array C. Sample output 2 3 4	*
Sample input 2 3 1 2 3 2 3 4 1 1 1	yo m x n sized matrix into two 2D array, suppose A and B. ay all the elements from matrix / 2D array C. Sample output 2 3 4	*
Now do C = A + B. Finally displa Sample input 2 3 1 2 3 2 3 4	yo m x n sized matrix into two 2D array, suppose A and B. ay all the elements from matrix / 2D array C. Sample output 2 3 4	*
Sample input 2 3 1 2 3 2 3 4 1 1 1 2 2 2	yo m x n sized matrix into two 2D array, suppose A and B. ay all the elements from matrix / 2D array C. Sample output 2 3 4	*
Sample input 2 3 1 2 3 2 3 4 1 1 1 2 2 2 WAP that will take inputs of tw do C = A * B (multiplication). Fi	Sample output 2 3 4 4 5 6 20 3 x 3 sized matrix into two 2D array, suppose A and B. Now nally display all the elements from matrix / 2D array C.	
Sample input 2 3 1 2 3 2 3 4 1 1 1 2 2 2 WAP that will take inputs of two do C = A * B (multiplication). Fi	yo m x n sized matrix into two 2D array, suppose A and B. ay all the elements from matrix / 2D array C. Sample output 2 3 4 4 5 6 yo 3 x 3 sized matrix into two 2D array, suppose A and B. Now nally display all the elements from matrix / 2D array C. Sample output	
Sample input 2 3 1 2 3 2 3 4 1 1 1 2 2 2 WAP that will take inputs of tw do C = A * B (multiplication). Fi Sample input 1 2 3	yo m x n sized matrix into two 2D array, suppose A and B. ay all the elements from matrix / 2D array C. Sample output	
Sample input 2 3 1 2 3 2 3 4 1 1 1 2 2 2 WAP that will take inputs of tw do C = A * B (multiplication). Fi Sample input 1 2 3 4 5 6	Sample output 2 3 4 4 5 6 20 3 x 3 sized matrix into two 2D array, suppose A and B. Now nally display all the elements from matrix / 2D array C. Sample output 2 3 4 4 5 6 Sample output 2 3 4 4 5 6	
Sample input 2 3 1 2 3 2 3 4 1 1 1 2 2 2 WAP that will take inputs of tw do C = A * B (multiplication). Fi Sample input 1 2 3 4 5 6 7 8 9	yo m x n sized matrix into two 2D array, suppose A and B. ay all the elements from matrix / 2D array C. Sample output	
Sample input 2 3 1 2 3 2 3 4 1 1 1 2 2 2 WAP that will take inputs of tw do C = A * B (multiplication). Fi Sample input 1 2 3 4 5 6	Sample output 2 3 4 4 5 6 20 3 x 3 sized matrix into two 2D array, suppose A and B. Now nally display all the elements from matrix / 2D array C. Sample output 2 3 4 4 5 6 Sample output 2 3 4 4 5 6	
Sample input 2 3 1 2 3 2 3 4 1 1 1 2 2 2 WAP that will take inputs of tw do C = A * B (multiplication). Fi Sample input 1 2 3 4 5 6 7 8 9	Sample output 2 3 4 4 5 6 20 3 x 3 sized matrix into two 2D array, suppose A and B. Now nally display all the elements from matrix / 2D array C. Sample output 2 3 4 4 5 6 Sample output 2 3 4 4 5 6	

Sample input	Sample output	
33	Max: 9	
123	Location: [2][1]	
456		
292		
2 3	Max: 9	
987	Location: [0][0]	
3 4 5		
be an odd number). Then cald	eger inputs into a square matrix of dimension n (where n must culate sum of the integers at first row, last row and two ease see the sample input-output.	*
Sample input	Sample output	
5	52	
4		
12345		
23416		
2 3 4 1 6 3 4 9 6 7		
2 3 4 1 6 3 4 9 6 7 4 2 6 7 8		
2 3 4 1 6 3 4 9 6 7		
2 3 4 1 6 3 4 9 6 7 4 2 6 7 8	23	
2 3 4 1 6 3 4 9 6 7 4 2 6 7 8 5 4 3 2 1	23	
2 3 4 1 6 3 4 9 6 7 4 2 6 7 8 5 4 3 2 1 7 1 1 1 1 1 1 1 1 1 1 1 1 1	23	
2 3 4 1 6 3 4 9 6 7 4 2 6 7 8 5 4 3 2 1 7 1 1 1 1 1 1 1 1 1 1 1 1 1	23	
2 3 4 1 6 3 4 9 6 7 4 2 6 7 8 5 4 3 2 1 7 1 1 1 1 1 1 1 1 1 1 1 1 1	23	
2 3 4 1 6 3 4 9 6 7 4 2 6 7 8 5 4 3 2 1 7 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	23	
2 3 4 1 6 3 4 9 6 7 4 2 6 7 8 5 4 3 2 1	23	

10. WAP that will take (n x n) integer inputs into a square matrix of dimension n (where n must be an odd number). Then calculate sum of the integers based on following position pattern (consider only the boxed position during the sum). Please see the input-output.

**

Sample input	Sample output
5 1 2 3 4 5 2 3 4 1 6 3 4 9 6 7 4 2 6 7 8 5 4 3 2 1	71
7 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	25

11. WAP that will take (n x n) integer inputs into a square matrix of dimension n (where n must be an odd number). Then calculate sum of the integers based on following position pattern (consider only the boxed position during the sum). Please see the input-output.

**

Sample input	Sample output	
5 1 2 3 4 5 2 3 4 1 6 3 4 9 6 7 4 2 6 7 8 5 4 3 2 1	65	
7 1 1 1 1 1 1 1 1 1 1 1 1 1	33	

12. WAP that will take (m x n) integer inputs into a matrix of dimension m x n. Now reverse that matrix within itself and display it. Reversal means swap 1st column with the nth column, swap 2nd column with the (n-1)th column and so on... Sample input Sample output 3 3 321 123 654 456 292 292 26 654321 123456 456789 987654 **13**. WAP that will take (n x n) integer inputs into a square matrix of dimension n. Now determine whether the matrix is symmetric or not. Reference: http://en.wikipedia.org/wiki/Symmetric matrix Sample input Sample output 3 Yes 1 7 3 7 4 5 3 5 6 2 No 1 3 4 2 14. WAP that will take $(m \times n)$ positive integer inputs into a matrix of dimension $m \times n$. Now replace all the duplicate integers by -1 in that matrix. Finally display it. Sample input Sample output 3 3 1 7 3 1 7 3 -1 4 5 7 4 5 -1 -1 6 3 5 6 26 2 -1 -1 -1 -1 2 2 2 2 2 2 6 5 4 3 - 1 1 6 5 4 3 2 1

Sample input	Sample output	
3 3	41	
1 7 3		
7 4 5		
3 5 6		
2 6	33	
2 2 2 2 2 2		
6 5 4 3 2 1		