

Selected Problems Chapter 5

Linear Algebra Done Right, Sheldon Axler, 3rd Edition

Mustaf Ahmed

August 3, 2021

Problem Example 5.8. Suppose $T \in \mathcal{L}(F^2)$ is defined by $T(w, z) = (-z, w)$. Find the eigenvectors and eigenvalues of T if $F = \mathbb{R}$. Find the eigenvectors and eigenvalues of T if $F = \mathbb{C}$

Proof. **Part(a).**

Assume T has eigenvectors and eigenvalues with $F = \mathbb{R}$. The equation $\lambda(w, z) = (-z, w)$ holds and leads to the following system of equations:

$$\lambda w = -z \tag{1}$$

$$\lambda z = w. \tag{2}$$

$$\tag{3}$$

Solving for λ , we have $\lambda^2 = -1$, which only has solutions in \mathbb{C} . This contradiction means T has no eigenvectors and eigenvalues.

Part(b).

In part(a), we showed that the eigenvalues of T must be in the complex numbers. The equation from part(a) $\lambda^2 = -1$ has the solutions $\lambda = i$ and $\lambda = -i$. The eigenvectors corresponding to $\lambda = i$ are of the form $(w, -iw)$ for any $w \in \mathbb{C}$; the eigenvectors corresponding to $\lambda = -i$ are of the form (w, iw) .

□