

Selected Problems Chapter 2

Linear Algebra Done Right, Sheldon Axler, 3rd Edition

Mustaf Ahmed

December 28, 2020

Problem 2.A.11. Suppose v_1, \dots, v_m is linearly independent in V and $w \in V$. Show that v_1, \dots, v_m, w is linearly independent if and only if $w \notin \text{span}(v_1, \dots, v_m)$.

Proof. For the forward direction, assume for a contradiction that the list v_1, \dots, v_m, w is linearly independent and $w \in \text{span}(v_1, \dots, v_m)$. We can then choose $a_1, \dots, a_m \in F$ such that

$$w = \sum_{i=1}^m a_i v_i,$$

so we have that

$$\left(\sum_{i=1}^m a_i v_i\right) - w = 0.$$

Since not all of the coefficients are equal to zero, the list v_1, \dots, v_m, w is not linearly independent, which is a contradiction.

For the backwards direction, assume for a contradiction that $w \notin \text{span}(v_1, \dots, v_m)$ and v_1, \dots, v_m, w is linearly dependent. We can choose $a_1, \dots, a_m, a_{m+1} \in F$, where not all of the coefficients are zero, such that

$$\left(\sum_{i=1}^m a_i v_i\right) + a_{m+1} w = 0.$$

Since v_1, \dots, v_m are linearly independent, a_{m+1} can't be equal to zero, otherwise we'd reach a contradiction. Thus, a_{m+1} is a non-zero coefficient, so we have that

$$w = \sum_{i=1}^m \frac{-a_i}{a_{m+1}} v_i$$

, by subtracting $a_{m+1}w$ and dividing out by $-a_{m+1}$. Thus, we conclude that $w \in \text{span}(v_1, \dots, v_m)$, which is a contradiction.

□

Problem 2.29 Basis Criterion. A list v_1, \dots, v_n of vectors in V is a basis of V if and only if every $v \in V$ can be written uniquely in the form

$$v = a_1v_1 + \dots + a_nv_n,$$

where $a_1, \dots, a_n \in F$.

Proof. For the forward direction, assume v_1, \dots, v_n are vectors in V that form a basis for V . Given $v \in V$, we want v to be written uniquely in the form

$$v = a_1v_1 + \dots + a_nv_n,$$

where $a_1, \dots, a_n \in F$. Since the list forms a basis for V , we can choose $a_1, \dots, a_n \in F$ such that

$$v = a_1v_1 + \dots + a_nv_n.$$

Suppose there exists $b_1, \dots, b_n \in F$ such that

$$v = b_1v_1 + \dots + b_nv_n.$$

. Then,

$$0 = (a_1 - b_1)v_1 + \dots + (a_n - b_n)v_n,$$

. so by linear independence, each coefficient must be equal, meaning that v is uniquely determined.

Next, we must show the backwards direction. Assume that every $v \in V$ can be written uniquely as a linear combination of v_1, \dots, v_n . By definition, v_1, \dots, v_n spans V . We must now show the list is linearly independent. The zero vector is in V , so $0 \in V$ can be written as a linear combination of v_1, \dots, v_n , namely

$$0 = 0v_1 + \dots + 0v_n,$$

. which is unique by assumption. Thus, the list satisfies the conditions of linear independence. \square

Problem 2.B.5. Prove or disprove : there exists a basis p_0, p_1, p_2, p_3 of $P_3(F)$ such that none of the polynomials p_0, p_1, p_2, p_3 has degree 2.

Proof. This is a true statement. Let $p_0 = 1, p_1 = x, p_2 = x^2 + x^3, p_3 = x^3$. We must show that this list of vectors spans $P_3(F)$ and is linearly independent.

Given $g = a + bx + cx^2 + dx^3 \in P_3(F)$, we must show the existence of coefficients in F such that g is in the span of the list of vectors defined earlier. Choosing $c_0 = a, c_1 = b, c_2 = c, c_3 = d - c$, we have that

$$\begin{aligned}
\left(\sum_{i=0}^3 c_i p_i\right) &= a + bx + c(x^2 + x^3) + (d - c)x^3 \\
&= a + bx + cx^2 + c(x^3 - x^3) + dx^3 \\
&= a + bx + cx^2 + dx^3 \\
&= g.
\end{aligned}$$

Next, we must show that our list of vectors in $P_3(F)$ is linearly independent. Given $g \in P_3(F)$, suppose there exists $a_0, a_1, a_2, a_3 \in F$ and $b_0, b_1, b_2, b_3 \in F$ such that

$$g = \left(\sum_{i=0}^3 a_i p_i\right) = \left(\sum_{i=0}^3 b_i p_i\right).$$

We want a unique representation g . Subtracting the two representations, we have that

$$\begin{aligned}
0 &= \left(\sum_{i=0}^3 a_i p_i\right) - \left(\sum_{i=0}^3 b_i p_i\right) = \left(\sum_{i=0}^3 (a_i - b_i) p_i\right) \\
&= (a_0 - b_0) + (a_1 - b_1)x + (a_2 - b_2)(x^2 + x^3) + (a_3 - b_3)x^3 \\
&= (a_0 - b_0) + (a_1 - b_1)x + (a_2 - b_2)x^2 + (a_2 - b_2)x^3 + (a_3 - b_3)x^3
\end{aligned}$$

This implies that $a_0 = b_0, a_1 = b_1, a_2 = b_2$. Since $a_2 = b_2$, the fourth term disappears and the final term must also have that $a_3 = b_3$. Thus, all vectors in $P_3(F)$ can be represented uniquely with our list of vectors.

□

Problem 2.B.7. Prove or give a counterexample: If v_1, v_2, v_3, v_4 is a basis for V and U is a subspace of V such that $v_1, v_2 \in U$ and $v_3 \notin U$ and $v_4 \notin U$, then v_1, v_2 is a basis for U .

Proof. This is a false statement. Let $V = \mathbb{R}^4$, with the basis being the standard basis. Consider the span following collection of vectors:

$$\begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 1 \\ 1 \end{pmatrix}$$

□

Problem 2.C.1. Suppose V is finite-dimensional and U is a subspace of V such that $\dim U = \dim V$. Prove that $U = V$.

Proof. Let $m = \dim U$. We can choose some basis for U to be $u_1, \dots, u_m \in U$. Since U is a subset of V , we can extend this basis to be a basis of V . However, since all basis for V have the same length, our basis for U can not be extended further, and thus, is already a basis for V . □

Problem 2.C.6(a). Let $U = \{p \in P_4(F) : p(2) = p(5)\}$. Find a basis for U .

Proof. First, we will propose a basis, and then prove that our list is a basis.

The constraints on the set U lead to the following equation:

$$a + 2b + 4c + 8d + 16e = a + 5b + 25c + 125d + 625e.$$

Solving for b , we get

$$b = -7c - 39d - 203e.$$

Thus, U is spanned by the following vectors:

$$a \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix} + c \begin{pmatrix} 0 \\ -7 \\ 1 \\ 0 \\ 0 \end{pmatrix} + d \begin{pmatrix} 1 \\ -39 \\ 0 \\ 1 \\ 0 \end{pmatrix} + e \begin{pmatrix} 1 \\ -203 \\ 0 \\ 0 \\ 1 \end{pmatrix}$$

. The vectors represent the polynomials $1, x^2 - 7x, x^3 - 39x, x^4 - 203x$, which are our proposed basis.

Since the list spans U , it is sufficient to show that the list is linearly independent. Each polynomial is not in the span of the previous polynomials in the list, so by the linear dependence lemma, the list is linearly independent. □

Problem 2.C.6(b). Extend the basis in part (a) to a basis of $P_4(F)$.

Proof. The standard basis of $P_4(F)$ is $1, x, x^2, x^3, x^4$; this means that the dimension of $P_4(F)$ is 5. We know that if a linearly independent list has the same length as a basis, the linear independent list is a basis itself. Thus, it is sufficient to extend our basis for U by a single polynomial.

Adding the polynomial x to our list, we get a new list: $1, x, x^2 - 7x, x^3 - 39x, x^4 - 203x$. Each polynomial in this new list is not in the span of polynomials before it because each polynomial has a unique highest degree. By the linear dependence lemma, this new list is linearly independent. □

Problem 2.C.6(c). Find a subspace W of $P_4(F)$ such that $P_4(F) = U \oplus W$.

Proof. We will first define the second vector space by

$$W = \{ax \mid a \in F\}.$$

Clearly, this is a subspace of $P_4(F)$.

We want that $U \cap W = \{0\}$ because this will imply that U and W form a direct sum. Suppose $e \in U \cap W$. Then we have the following equation

$$e = ax = b + c(x^2 - 7x) + d(x^3 - 39x) + e(x^4 - 203x)$$

, where all coefficients are in F . Since e is at most degree 1, c, d and e must be zero. Thus, we have that $e = ax = b$, which means that a is zero and e is the zero vector.

Next, we must show that $P_4(F) = U \oplus W$. A basis $U \oplus W$ can clearly be the basis we defined for $P_4(F)$, so the two sets are equal. □

Problem 2.C.11. Suppose that U and W are subspaces of \mathbb{R}^8 such that $\dim U = 3$, $\dim W = 5$ and $U + W = \mathbb{R}^8$. Prove that $\mathbb{R}^8 = U \oplus W$

Proof. We will prove this by contradiction. Suppose there exists a non-zero vector $v \in U \cap W$. Let $B = \{u_1, u_2, u_3, w_1, w_2, w_3, w_4, w_5\}$, where u_i form a basis for U and w_j form a basis for W . Since $U + W = \mathbb{R}^8$, B spans \mathbb{R}^8 . Also, $|B| = 8$ and $\dim \mathbb{R}^8 = 8$, so B is a basis for \mathbb{R}^8 .

We want a contradiction with the fact that B is a linearly independent list. Choose $a_1, a_2, a_3 \in F$ and $b_1, b_2, b_3, b_4, b_5 \in F$ such that

$$v = a_1u_1 + a_2u_2 + a_3u_3,$$

and

$$v = b_1w_1 + b_2w_2 + b_3w_3 + b_4w_4 + b_5w_5.$$

Then, we have that

$$0 = a_1u_1 + a_2u_2 + a_3u_3 - (b_1w_1 + b_2w_2 + b_3w_3 + b_4w_4 + b_5w_5),$$

where not all the coefficients are zero. This means that B is not a linearly independent list, which is a contradiction. □

Problem 2.C.12. Suppose U and W are both five-dimensional subspaces of \mathbb{R}^9 . Prove that $U \cap W \neq \{0\}$.

Proof. Using theorem 2.43, we have that

$$\begin{aligned} \dim(U + W) &= \dim U + \dim W - \dim(U \cap W) \\ &= 10 - \dim(U \cap W). \end{aligned}$$

Since $U + W$ is a subspace of \mathbb{R}^9 , $\dim(U + W) \leq \dim(\mathbb{R}^9)$. Thus, we have

$$\dim(\mathbb{R}^9) = 9 \geq 10 - \dim(U \cap W)$$

By rearranging terms, $\dim(U \cap W) \geq 1$, so $U \cap W$ is not the trivial subspace. □