

# Selected Problems Chapter 5

## Linear Algebra Done Right, Sheldon Axler, 3rd Edition

Mustaf Ahmed

August 4, 2021

**Problem Example 5.8.** Suppose  $T \in \mathcal{L}(F^2)$  is defined by  $T(w, z) = (-z, w)$ . Find the eigenvectors and eigenvalues of  $T$  if  $F = \mathbb{R}$ . Find the eigenvectors and eigenvalues of  $T$  if  $F = \mathbb{C}$

*Proof.* **Part(a).**

Assume  $T$  has eigenvectors and eigenvalues with  $F = \mathbb{R}$ . The equation  $\lambda(w, z) = (-z, w)$  holds and leads to the following system of equations:

$$\lambda w = -z \tag{1}$$

$$\lambda z = w. \tag{2}$$

$$\tag{3}$$

Solving for  $\lambda$ , we have  $\lambda^2 = -1$ , which only has solutions in  $\mathbb{C}$ . This contradiction means  $T$  has no eigenvectors and eigenvalues.

**Part(b).**

In part(a), we showed that the eigenvalues of  $T$  must be in the complex numbers. The equation from part(a)  $\lambda^2 = -1$  has the solutions  $\lambda = i$  and  $\lambda = -i$ . The eigenvectors corresponding to  $\lambda = i$  are of the form  $(w, -iw)$  for any  $w \in \mathbb{C}$ ; the eigenvectors corresponding to  $\lambda = -i$  are of the form  $(w, iw)$ .

□

**Problem Theorem 5.10 Linearly Independent Eigenvectors.** Let  $T \in \mathcal{L}(V)$ . Suppose  $(\lambda_1, \dots, \lambda_n)$  are distinct eigenvalues of  $T$ , and  $(v_1, \dots, v_n)$  are corresponding eigenvectors. Then  $(v_1, \dots, v_n)$  is a linearly independent list.

*Proof.* For a contradiction, suppose  $(v_1, \dots, v_n)$  is a linearly dependent list. Then choose  $a_1, \dots, a_n \in F$  where not all are zero such that  $0 = a_1 v_1 + \dots + a_n v_n$ . By the linear dependence lemma, choose the smallest  $j$  such that  $v_j = \frac{a_1}{a_j} v_1 + \dots + \frac{a_{j-1}}{a_j} v_{j-1}$ . Applying  $T$ , we have

$$\lambda_j v_j = \frac{a_1 \lambda_1}{a_j} v_1 + \dots + \frac{a_{j-1} \lambda_{j-1}}{a_j} v_{j-1}. \quad (4)$$

Subtracting the left side, we have

$$0 = \frac{a_1(\lambda_1 - \lambda_j)}{a_j} v_1 + \dots + \frac{a_{j-1}(\lambda_{j-1} - \lambda_j)}{a_j} v_{j-1}. \quad (5)$$

Each  $(\lambda_k - \lambda_j) > 0$  because the eigenvalues are distinct. Since we chose  $j$  to be the smallest such that the  $v_j$  is in the span of the preceding vectors,  $a_k = 0$  for  $k = 1, \dots, j-1$ . Thus,  $v_j = 0$ , but that is a contradiction because  $v_j$  is an eigenvector. □