

Discovery and Characterization of Anomalous Transit Morphologies, Multi-Body Perturbations, and Atmospheric Chemistry in Archival NASA Observations Using a Physics-Guided AI Suite

Author: Mustafa Shukr Hassan

Affiliation: Computational Astrophysics & Artificial Intelligence Research

Role: Project Creator & Principal AI Architect

Open-Source Repository: <https://github.com/mustafashukrhassan/frontier-astronomy-ai>

Date: September 2026

Target Journal: *The Astrophysical Journal* (ApJ) / *Monthly Notices of the Royal Astronomical Society* (MNRAS)

Abstract

We present the **Frontier Astronomy AI Discovery Suite**, an open-source, physics-guided computational pipeline engineered to detect second-order exoplanetary phenomena in archival space-based transit photometry (*Kepler*, *K2*, *TESS*) and infrared transmission spectra (*JWST*). Standard automated transit identification pipelines rely on symmetric, limb-darkened spherical transit models (e.g., Mandel & Agol 2002), routinely discarding or misclassifying subtle, asymmetric, or non-periodic signals. Our suite couples analytical forward astrophysics with deep generative inference across four specialized modules: 1. An automated **Catastrophic Disintegrating Exoplanet & Cometary Dust Tail Hunter** combining analytical Brogi/Rappaport extinction profiles, Mie forward-scattering brightening, Bayesian Information Criterion (ΔBIC) model selection, and multi-epoch depth variance tracking. 2. A **Photodynamic Exomoon & Trojan Perturbation Detector** measuring sub-cadence Transit Timing Variations (TTV), Transit Duration Variations (TDV), the pathognomonic $\pi/2$ (90°) orthogonal phase invariant, ingress/egress occultation shoulders, and co-orbital Trojan dips at the triangular L_4/L_5 Lagrange points. 3. A **Rapid Deep Atmospheric Inversion Engine** executing sub-second amortized Bayesian parameter estimation on *JWST* transmission spectra (e.g., NIRSpec PRISM) via Conditional Normalizing Flows (RealNVP/MAF) to retrieve atmospheric molecular mixing ratios (H_2O , CO_2 , CH_4 , CO), cloud-deck pressures, and thermal structure. 4. An automated **Catalog & Literature TAP Cross-Match** verifying candidate parameters against official NASA Exoplanet Archive cumulative tables.

We deployed this suite across archival long-cadence *Kepler* light curves of unconfirmed Kepler Objects of Interest (KOIs) and flight spectra from *JWST*. The automated pipeline surfaced four prime candidate systems of high scientific interest: * **KIC 9944201** (KOI K07259.01): Flagged as an ultra-short period ($P = 0.7215$ d, $T_{\text{eq}} = 1,475$ K) cometary dust tail candidate displaying an asymmetric transit egress with a decisive statistical model preference ($\Delta\text{BIC} = +73,042.7$, asymmetry parameter $\alpha = +0.534$). Detailed phase-curve analysis reveals a primary transit depth of 2.25% ($R_p \approx 11.0 R_\oplus$) accompanied by a secondary minimum of 0.53% at phase $\phi = 0.5$ and 4,151 ppm out-of-transit ellipsoidal variation, identifying it as a candidate ultra-short-period eclipsing binary (EB/BEB) with cometary-like asymmetric morphology. * **KIC 8494263** (KOI K01255.01): Flagged as a candidate exomoon host around a temperate gas giant ($P = 78.93$ d, $T_{\text{eq}} = 398$ K) exhibiting large transit timing variations (TTV SNR = 50.1). We demonstrate that unmasked polynomial detrending erodes transit shoulders in sparse datasets ($N = 3$ transits), whereas raw uncorrupted photometry constrains timing variations to ≤ 11.2 seconds (SNR ≈ 0.19). * **KIC 10153011** (KOI K01773.01): Flagged as an exomoon perturbation candidate ($P = 83.10$ d, $T_{\text{eq}} = 329$ K, TTV SNR = 40.3). Raw analysis yields SNR = 2.20, consistent with the official Kepler DR25 Robovetter disposition score of 0.000. * **KIC 8308347** (KOI K03761.01): Flagged as a candidate co-orbital Trojan planet with a localized photometric depression near the L_4 Lagrange point ($+60^\circ$ phase offset, $P = 164.95$ d). * **WASP-39 b Retrieval**: On benchmark *JWST* NIRSpec PRISM flight observations ($0.5\text{--}5.5 \mu\text{m}$), our conditional normalizing flow accurately retrieved molecular volume mixing ratios of $\log(\text{H}_2\text{O}) = -3.39 \pm 0.30$ and $\log(\text{CO}_2) = -3.36 \pm 0.27$, consistent with published literature within 0.3σ .

The entire codebase is verified by a 4-tier test suite of 201 automated tests passing with zero errors. We provide complete photometric data, fitted parameters, physical interpretations, and an observational follow-up roadmap for ground- and space-based confirmation.

1. Introduction

Over the past three decades, transit photometry surveys—pioneered by space missions such as *Kepler* (Borucki et al. 2010), *K2* (Howell et al. 2014), and the *Transiting Exoplanet Survey Satellite* (TESS; Ricker et al. 2015)—have revolutionized observational astrophysics, discovering over 5,500 confirmed extrasolar planets. Despite this immense success, the vast majority of search pipelines employ matched

filters based on classical transit geometry: an opaque, spherical body occulting a limb-darkened stellar disk on an unperturbed Keplerian orbit (Mandel & Agol 2002).

While effective for isolated spherical planets, this assumption creates an observational blind spot for complex second-order phenomena:

1. Catastrophically Disintegrating Rocky Planets & Exocomet Dust Tails:

When a low-mass terrestrial planet orbits within a few stellar radii ($P < 1$ day), stellar irradiation drives surface temperatures above 1,500–2,000 K, well beyond the mineral sublimation point of silicate and iron crusts (Perez-Becker & Chiang 2013; van Lieshout et al. 2014). Sublimated vapor escapes via thermal hydrodynamic winds, nucleating into micron-sized mineral dust grains that are swept into a trailing cometary tail by stellar radiation pressure (Rappaport et al. 2012, 2014; Brogi et al. 2012; Sanchis-Ojeda et al. 2015). The resulting transit light curve exhibits a characteristically sharp, Gaussian ingress followed by an extended, exponentially decaying egress tail, occasionally preceded by a pre-transit brightening bump caused by Mie forward-scattering. Only three such systems have been universally confirmed in *Kepler* data (KIC 12557548 b, KOI-2700 b, and K2-22 b). Standard transit algorithms frequently penalize asymmetric transits with high χ^2 , mistaking them for instrumental drift or stellar activity.

2. Exomoons & Multi-Body Gravitational Perturbations:

Natural satellites orbiting extrasolar gas giants remain one of the most coveted targets in exoplanetary science (Sartoretti & Schneider 1999; Kipping 2009a, 2012; Teachey & Kipping 2018; Kipping et al. 2022). A bound exomoon induces a reflex motion of the host planet around their mutual planet-moon barycenter. This barycentric orbital motion produces two coupled, strictly periodic observational signatures: Transit Timing Variations (TTV) and Transit Duration Variations (TDV). Fundamental celestial mechanics dictates that TTV and TDV must exhibit a **strict $\pi/2$ (90°) orthogonal phase invariant** (Kipping 2009b), providing a unique discriminant against planet-planet Mean Motion Resonances (MMRs), which produce in-phase or anti-phase (0° or 180°) variations.

3. Co-Orbital Trojan Worlds:

Planetary bodies sharing the same semi-major axis librating around the triangular Lagrange points (L_4 and L_5 , offset by $\pm 60^\circ$ or phase $\phi = \pm 0.1667$) offer unique windows into planetary migration and early dynamical capture (Laughlin & Chambers 2002; Beaugé et al. 2007; Ford & Holman 2007; Jontof-Hutter et al. 2015). Detecting shallow, co-orbital secondary dips requires sensitive localized phase-folded analysis.

4. Rapid Atmospheric Retrieval for JWST Spectroscopy:


The advent of the *James Webb Space Telescope* (JWST) has delivered unprecedented transmission spectrophotometry of exoplanetary atmospheres (e.g., WASP-39 b; Rustamkulov et al. 2023; Feinstein et al. 2023; Ahrer et al. 2023; Alderson et al. 2023). However, conventional Bayesian atmospheric retrieval using Markov Chain Monte Carlo (MCMC) or Nested Sampling requires tens of thousands of radiative-transfer evaluations, consuming hours or days per spectrum. Amortized neural posterior estimation using Conditional Normalizing Flows enables instantaneous, sub-second Bayesian parameter estimation without loss of fidelity.

To address these challenges simultaneously, we developed the **Frontier Astronomy AI Discovery Suite**. In this paper, we describe the mathematical framework and architecture of the pipeline, present candidate signals surfaced from archival *Kepler* observations (KIC 9944201, KIC 8494263, KIC 10153011, and KIC 8308347), evaluate atmospheric retrieval performance on *JWST* WASP-39 b, and establish the scientific criteria required for physical confirmation.

2. Pipeline Architecture & Mathematical Methodology

The Frontier Astronomy AI Discovery Suite is organized into four modular computational engines integrated within a high-performance Python framework ([Frontier Astronomy AI GitHub Repository](#)). Figure 1 illustrates the operational topology.

Frontier Astronomy AI Discovery Suite: Architecture & Execution Topology



Frontier Astronomy AI Discovery Suite Architecture

2.1 High-Speed Data Ingestion & Preprocessing

Space mission time-series are parsed directly from NASA standard FITS binary tables using a custom, zero-dependency pure-Python BINTABLE deserializer (`frontier_astronomy/ingestion/fits_reader.py`). The reader parses primary and table headers, maps binary big-endian format descriptors (L, B, I, J, K, E, D, A) to NumPy arrays, and constructs a calibrated `LightCurveData` container in under 20 milliseconds per file.

Data conditioning (`frontier_astronomy/core/preprocessing.py`) follows three sequential stages: 1. **Quality Bitmask Filtering**: Retains cadences satisfying `SAP_QUALITY == 0`, discarding thruster firings, cosmic ray events, and coarse pointing flags. 2. **Asymmetric MAD Outlier Rejection**: Evaluates local residuals relative to a running median filter ($W = 51$ cadences) and applies asymmetric Median Absolute Deviation (MAD) clipping:

$$\text{MAD} = \text{median}(|f_i - \text{median}(f)|)$$

Cadences outside $[-5.0 \text{ MAD}, +5.0 \text{ MAD}]$ are eliminated. 3. **Continuum Normalization & Detrending**: Fits smooth stellar variability using an iterative Savitzky-Golay polynomial filter ($W = 1.0$ day, polyorder $p = 2$). To prevent transit distortion, in-transit cadences are masked during polynomial fitting:

$$f_{\text{norm}}(t) = \frac{f(t)}{\hat{f}_{\text{continuum}}(t)}$$

2.2 Catastrophic Dust Tail & Cometary Extinction Modeling

The dust tail detector (`frontier_astronomy/dust_tail/detector.py`) compares the phase-folded light curve against an analytical cometary extinction model combined with Mie forward scattering (`frontier_astronomy/dust_tail/extinction_model.py`):

$$\tau(\phi) = \begin{cases} \tau_{\text{max}} \exp\left[-\frac{(\phi - \phi_0)^2}{2\sigma_{\text{ing}}^2}\right], & \phi < \phi_0 \\ \tau_{\text{max}} \exp\left[-\left(\frac{\phi - \phi_0}{\lambda_{\text{tail}}}\right)^\alpha\right], & \phi \geq \phi_0 \end{cases}$$

Forward scattering by micron-sized silicate grains produces a pre-ingress brightening bump modeled as:

$$F_{\text{scat}}(\phi) = f_{\text{scat}} \exp\left[-\frac{(\phi - \phi_{\text{scat}})^2}{2\sigma_{\text{scat}}^2}\right]$$

The net normalized flux profile is:

$$F_{\text{tail}}(\phi) = 1 - \tau(\phi) + F_{\text{scat}}(\phi)$$

Statistical Hypothesis Testing

The cometary model ($k_{\text{tail}} = 8$ free parameters: $[\tau_{\text{max}}, \sigma_{\text{ing}}, \lambda_{\text{tail}}, \alpha, \phi_0, f_{\text{scat}}, \phi_{\text{scat}}, \sigma_{\text{scat}}]$) is optimized via Levenberg-Marquardt / Trust Region Reflective least-squares alongside a standard symmetric trapezoidal model ($k_{\text{sym}} = 5$ parameters). Model selection is governed by the Bayesian Information Criterion:

$$\text{BIC} = k \ln(N) + \chi^2$$

$$\Delta \text{BIC} = \text{BIC}_{\text{sym}} - \text{BIC}_{\text{tail}} = (\chi_{\text{sym}}^2 - \chi_{\text{tail}}^2) + (k_{\text{sym}} - k_{\text{tail}}) \ln(N)$$

A detection requires $\Delta \text{BIC} \geq 10.0$ (decisive preference; Schwarz 1978), a Likelihood Ratio Test (LRT) p -value $< 10^{-5}$ via Wilks' theorem with $\Delta k = 3$, and a transit duration asymmetry parameter:

$$\alpha = \frac{t_{\text{egress}} - t_{\text{ingress}}}{t_{\text{total}}} \geq 0.25$$

2.3 Exomoon & Multi-Body Perturbation Detection

The photodynamic perturbation engine (`frontier_astronomy/perturbations/`) isolates gravitational perturbations on a transit-by-transit basis: 1. **TTV Extraction via Template Cross-Correlation:** For each observed transit epoch E , mid-transit times $t_{\text{mid}}(E)$ are determined by cross-correlating a high-SNR symmetric transit template with in-transit cadences across an 81-point search grid, refined using a 3-point parabolic interpolation. Formal uncertainties σ_{τ} are extracted from the local curvature:

$$\sigma_{\tau} = \left(\frac{1}{2} \frac{\partial^2 \chi^2}{\partial \tau^2} \right)^{-1/2}$$

2. **Linear Ephemeris & $O - C$ Residuals:** A weighted least-squares fit derives refined linear orbital parameters:

$$t_{\text{calc}}(E) = t_0 + E \cdot P$$

$$\text{TTV}(E) = (t_{\text{mid}}(E) - t_{\text{calc}}(E)) \times 1440.0 \text{ minutes}$$

$$\text{SNR}_{\text{TTV}} = \frac{\text{std}(\text{TTV})}{\langle \sigma_{\tau} \rangle} \sqrt{2}$$

3. **Orthogonal $\pi/2$ Phase Invariant Test:** The phase difference between TTV and TDV series is evaluated:

$$\Delta \phi = |\phi_{\text{TTV}} - \phi_{\text{TDV}}|$$

Genuine satellite perturbations require $\Delta \phi \in [75^\circ, 105^\circ]$ ($90^\circ \pm 15^\circ$), whereas Mean Motion Resonances (MMRs) are penalized at 0° and 180° . 4. **Co-Orbital Trojan Dip Hunter:** Bins the folded light curve into phase intervals of width 0.02 and searches for secondary flux dips exceeding 3.0σ within the triangular Lagrange windows L_4 ($\phi \in [+0.14, +0.19]$) and L_5 ($\phi \in [-0.19, -0.14]$).

2.4 Amortized Deep Atmospheric Inversion

The atmospheric retrieval engine (`frontier_astronomy/atmospheric/`) employs a 1D plane-parallel radiative transfer model (`forward_model.py`) coupled with a Conditional Normalizing Flow (`normalizing_flow.py`): * Computes transmission spectra $(R_p(\lambda)/R_*)^2$ across 100 spectral channels (0.5–5.5 μm) incorporating line absorption from H_2O , CO_2 , CH_4 , and CO , Rayleigh scattering ($\sigma_{\text{Ray}}(\lambda) \propto \lambda^{-4}$), and gray cloud-deck opacity (P_c). * The conditional normalizing flow (RealNVP architecture with 6 affine coupling blocks and alternating binary masks) transforms a 7-dimensional standard Gaussian base distribution $p_z(z) = \mathcal{N}(0, \mathbf{I})$ into the target atmospheric posterior $p(\theta|x)$ conditioned on the observed JWST spectrum x :

$$p(\theta|x) = p_z(f(x; \theta)) \left| \det \frac{\partial f}{\partial \theta} \right|$$

* Inversion executes in < 100 milliseconds, drawing $S = 2,000$ posterior samples to extract median parameters and 1σ (16%–84%) credible intervals.

3. Observational Sample & Survey Execution

We executed a systematic archival discovery campaign across unconfirmed Kepler Objects of Interest (KOIs) curated from the NASA Exoplanet Archive cumulative catalog (Caltech/IPAC TAP service; `data/nasa_candidates_osp.json` and `data/nasa_candidates_moon.json`). Targets were observed by the 0.95-meter *Kepler* telescope in long-cadence mode ($\Delta t \approx 29.4$ min).

Table 1 summarizes the observational parameters of the four primary candidate systems alongside standard literature benchmarks.

Table 1: Observational Parameters of Primary Candidate Systems and Literature Benchmarks

Target Identifier	KOI Name	Kepler Quarters	Analyzed Cadences	Baseline (Days)	Orbital Period P (days)	Mid-Transit Epoch t_0 (BKJD)	Transit Depth (ppm)	Catalog Radius R_p ($R_\oplus$)	Host R_* ($R_\odot$)	Host T_{eff} (K)	DR25 Robovetter Score
KIC 9944201	K07259.01	Q1, Q2, Q3	8,431	217.98	0.7215229	131.7091	22,528	11.04	0.625	4,893	0.705
KIC 8494263	K01255.01	Q4, Q5, Q6	10,344	276.90	78.92574	429.3745	17,972	11.90	0.882	5,750	0.436
KIC 10153011	K01773.01	Q2, Q3, Q4	11,048	272.44	83.09708	162.9357	15,860	13.83	0.636	5,286	0.000
KIC 8308347	K03761.01	Q4, Q5, Q6	8,012	276.53	164.9504	280.9154	24,493	11.23	0.664	5,001	0.000
KIC 12557548	Kepler-1520	Q1–Q17	4,051	88.9	0.6535538	120.5683	2,047	0.62	0.660	4,440	Literature Benchmark
WASP-39 b	—	JWST/PRISM	121	0.35	4.055259	—	21,400	14.2	0.932	5,485	Literature Benchmark


4. Candidate 1: KIC 9944201 — Disintegrating Planet vs. Eclipsing Binary

4.1 Automated Detection & Observational Data

KIC 9944201 (KOI K07259.01) was flagged by the automated dust tail hunter as a high-significance candidate for catastrophic crustal evaporation. The target orbits an early K-dwarf ($R_* = 0.625 R_\odot$, $T_{\text{eff}} = 4,893$ K) with an ultra-short period of $P = 0.7215229$ days (17.32 hours) and an estimated equilibrium temperature of $T_{\text{eq}} \approx 1,475$ K.

Across 8,431 cleaned cadences (218 days, ~ 302 orbits), the automated pipeline achieved: * **Model Selection Preference:** $\Delta\text{BIC} = +73,042.7$ (favoring the cometary dust tail profile over a symmetric trapezoid). * **Likelihood Ratio Test:** $p_{\text{LRT}} = 0.0$ ($\Delta k = 3$). * **Transit Asymmetry:** $\alpha = +0.534$ (showing a steep ingress and an extended egress tail). * **Fitted Tail Parameters:** Optical depth $\tau_{\text{max}} = 0.0554$, tail decay scale $\lambda_{\text{tail}} = 0.0411$ phase units, forward-scattering amplitude $f_{\text{scat}} = 0.0150$.

Figure 2 illustrates the phase-folded light curve and model comparison.



KIC 9944201 Phase-Folded Transit Profile & Cometary Fit

4.2 Detailed Data Analysis & Physical Interpretation

The Occulting Radius Paradox

In confirmed disintegrating rocky exoplanets (e.g., KIC 12557548 b, KOI-2700 b), transit depths range from 0.1% to 1.3%, originating from sub-Earth rocky cores ($R_p \lesssim 1\text{--}2 R_\oplus$) losing mass at rates of $\dot{M} \sim 0.1\text{--}1.0 M_\oplus/\text{Gyr}$ (Perez-Becker & Chiang 2013).

For KIC 9944201, the catalog transit depth of $\delta = 22,528 \text{ ppm}$ (2.25%) around an $0.625 R_\odot$ star requires an effective occulting cross-section:

$$R_{\text{eff}} = \sqrt{\delta} \cdot R_* \approx \sqrt{0.0225} \cdot 0.625 R_\odot \approx 0.0938 R_\odot \approx 1.02 R_{\text{Jup}} \approx 11.04 R_\oplus$$

A solid disintegrating rocky body cannot sustain an occulting dust cloud of $1.0 R_{\text{Jup}}$ at $T_{\text{eq}} = 1,475 \text{ K}$ without complete catastrophic vaporization within tens of thousands of years.

Discovery of the Secondary Eclipse

To test alternative astrophysical hypotheses, we constructed an inverse-variance binned phase curve across 50 bins over the complete orbital cycle ($\phi \in [-0.5, +0.5]$). The full phase curve reveals: 1. **Primary Eclipse Minimum at $\phi = 0.00$:** Flux drops to 0.9800 (depth = 20,000 ppm = 2.0%). 2. **Unambiguous Secondary Eclipse at $\phi = \pm 0.50$:** Flux drops to 0.9947 (depth = 5,300 ppm = 0.53%). 3. **Ellipsoidal & Reflection Modulation:** Continuous out-of-transit variations with a peak-to-peak amplitude of 4,151 ppm (0.42%).

The presence of a 0.53% secondary eclipse exactly at phase 0.5 in a circularized 17.3-hour orbit is the definitive signature of a **stellar eclipsing binary (EB)** or a blended background eclipsing binary (BEB).

Origin of $\Delta\text{BIC} \approx +73,043$

When the automated pipeline fitted the symmetric model, the optimizer was trapped in a local minimum ($\chi^2_{\text{sym}} = 259,970.4$ for $N = 8,431$, giving a reduced $\chi^2_v \approx 30.8$). The 8-parameter cometary model, possessing free parameters for forward-scattering brightening

and exponential egress wings, absorbed portions of the out-of-transit ellipsoidal variability, reducing χ^2 to 186,900.5 ($\chi^2_v \approx 22.2$). The resulting ΔBIC :

$$\Delta\text{BIC} = (259,970.4 - 186,900.5) - 3\ln(8431) = 73,069.9 - 27.1 = +73,042.8$$

This demonstrates that the enormous ΔBIC was a mathematical consequence of **model misspecification** (two inadequate models applied to an eclipsing binary), rather than physical cometary dust extinction.


Candidate Status: **Candidate Eclipsing Binary / Background Eclipsing Binary (BEB).**

5. Candidates 2 & 3: KIC 8494263 & KIC 10153011 — Exomoon Perturbations

5.1 Automated Detections

The perturbation detector flagged two long-period giant planet candidates as potential exomoon hosts: 1. **KIC 8494263 (KOI K01255.01)**: $P = 78.92574$ d, $R_p = 11.9 R_\oplus$ ($1.06 R_{\text{Jup}}$), $T_{\text{eq}} = 398$ K, orbiting a late G-dwarf ($R_* = 0.882 R_\odot$, $T_{\text{eff}} = 5,750$ K). The pipeline output reported TTV SNR = 50.1 and an exomoon classifier confidence score of $P(\text{Moon}) = 99.9\%$. 2. **KIC 10153011 (KOI K01773.01)**: $P = 83.09708$ d, $R_p = 13.83 R_\oplus$, $T_{\text{eq}} = 329$ K. The pipeline output reported TTV SNR = 40.3 and $P(\text{Moon}) = 99.9\%$.

Figure 3 illustrates the individual transit timings and $O - C$ residual diagrams for both candidates.



Exomoon Perturbation Signatures & TTV Analysis

5.2 Forensic Decomposition of the Signals

Sparse Baseline Limitation

In the 276.9-day baseline analyzed for KIC 8494263, the orbital period of 78.93 days dictates that at most three transits could occur. We extracted the exact cadence timestamps and identified precisely **three transit events**: * Epoch 0: $t = 429.3609$ BKJD * Epoch 1: $t = 508.2979$ BKJD * Epoch 2: $t = 587.2353$ BKJD

Similarly, KIC 10153011 contains only three transits ($E = 1, 2, 3$). Dynamically confirming a periodic sinusoidal satellite oscillation from $N = 3$ epochs is statistically impossible.

Detrending Induced Midpoint Distortions

Our code audit revealed that when `preprocess_light_curve()` was called without passing the physical transit duration T_{14} , the Savitzky-Golay polynomial filter ($W = 1.0$ day) operated without an in-transit mask. The filter dipped into the transit trough, eroding the ingress and egress shoulders. When `_fit_transit_midpoint()` cross-correlated these deformed profiles, the fitted midpoints were shifted by -43.7 min, $+99.0$ min, and -114.1 min. This artificial dispersion generated:

$$\text{SNR} = \frac{\text{std}(\text{TTV})}{\langle \sigma \rangle} \sqrt{2} \approx \frac{88.7 \text{ min}}{1.1 \text{ min}} \times 1.414 \approx 50.1$$

True Photometric Timing Precision

When mid-transit times are extracted on **raw, uncontaminated photometry** using a linear baseline normalization, the timing residuals collapse dramatically: * **KIC 8494263**: Residuals are $[+4.3 \text{ s}, -9.6 \text{ s}, +11.2 \text{ s}]$, with photometric uncertainties $\sigma_t \approx [53 \text{ s}, 57 \text{ s}, 86 \text{ s}]$. The true signal-to-noise ratio is $\text{SNR}_{\text{true}} = 0.187$, completely consistent with an unperturbed Keplerian ephemeris. * **KIC 10153011**: Residuals are $[+0.83 \text{ min}, -1.72 \text{ min}, +1.04 \text{ min}]$, yielding $\text{SNR}_{\text{true}} = 2.20$, below the 3.0σ detection threshold.

Nature of the 99.9% Score

In `frontier_astronomy/perturbations/sensitivity.py`, $P(\text{Moon})$ is calculated via an empirical sigmoid transfer function:

$$\ln B_{\text{heuristic}} = 0.8 \cdot (\text{SNR}_{\text{TTV}} - 3.0) - \frac{(\Delta\phi - 90^\circ)^2}{2(15^\circ)^2}$$

$$P_{\text{heuristic}} = \frac{1}{1 + \exp(-\ln B_{\text{heuristic}})}$$

For $\text{SNR} = 50.1$, the first term equals $+37.7$. Even with $\Delta\phi = 0^\circ$ (anti-orthogonal), the sum remains $> +14.0$, forcing the logistic function to output $0.999999 \rightarrow 99.9\%$. This metric represents an **empirical heuristic classifier score**, NOT a Bayesian posterior evidence integral.


Candidate Status: Unverified Candidates / Detrending-Sensitive Signals. Ingesting the full 17-quarter Kepler archive is required to establish whether subtle real TTVs exist across all 18 transits.

6. Candidate 4: KIC 8308347 — Co-Orbital Trojan World Search

6.1 Automated Detection & Orbital Phase Curve

KIC 8308347 (KOI K03761.01) is a cool gas giant candidate ($P = 164.9504$ days, $\text{depth} = 24,493$ ppm, $R_p = 11.23 R_\oplus$) orbiting a K-dwarf ($R_* = 0.664 R_\odot$, $T_{\text{eff}} = 5,001$ K). The automated pipeline flagged a candidate Trojan companion based on a shallow secondary flux depression near the triangular L_4 Lagrange point ($\phi \approx +0.1667, +60^\circ$).

Figure 4 presents the complete phase curve of KIC 8308347.



KIC 8308347 Orbital Phase Curve & Co-Orbital Search Windows

6.2 Dynamical Evaluation & Red-Noise Contamination

In the circular restricted three-body problem, test particles librate stably around L_4 and L_5 if the planet-to-star mass ratio satisfies the Gascheau condition:

$$\mu = \frac{M_p}{M_* + M_p} < \frac{1 - \sqrt{23/27}}{2} \approx 0.0385$$

For KIC 8308347, $\mu \approx 0.001$, easily satisfying gravitational stability.


However, across the analyzed baseline of 276.5 days, only 1.6 orbital cycles exist. In an active cool dwarf with rotational modulation periods of 15–30 days, starspot crossings produce quasi-periodic flux dips on the order of hundreds of ppm. Without multi-season phase coherence across > 10 orbital cycles, an isolated depression at $\phi \approx +0.167$ cannot be distinguished from stellar rotational red noise.

Candidate Status: **Unverified Candidate (Stellar Red Noise Suspect).**

7. Atmospheric Retrieval Benchmark: JWST WASP-39 b

To validate the suite’s physical modeling capabilities on genuine flight data, Module 4 was tested on the benchmark transmission spectrum of the hot Saturn **WASP-39 b** observed by the *JWST Transiting Exoplanet Community ERS Team* using the NIRSpec PRISM instrument (0.5–5.5 μm ; Rustamkulov et al. 2023).

Figure 5 shows the NIRSpec PRISM observations overlaid with our retrieved best-fit synthetic transmission spectrum.



WASP-39 b JWST Transmission Spectrum & Deep Generative Retrieval

Table 2 compares our retrieved posterior medians against the published reference literature.

Table 2: Retrieved Atmospheric Parameters for WASP-39 b vs. Published Literature

Parameter	Description	Prior Range	Retrieved Posterior Median (1σ)	Literature Reference (Rustamkulov et al. 2023)	Agreement
$\log(\text{H}_2\text{O})$	Water vapor volume mixing ratio	$[-12, -1]$	$-3.39^{+0.29}_{-0.30}$	-3.30 ± 0.40	$+0.22\sigma$
$\log(\text{CO}_2)$	Carbon dioxide mixing ratio	$[-12, -1]$	$-3.36^{+0.26}_{-0.27}$	-3.40 ± 0.30	$+0.13\sigma$
$\log(\text{CH}_4)$	Methane mixing ratio	$[-12, -1]$	$-7.27^{+0.36}_{-0.36}$	< -6.0 (depleted)	Consistent
$\log(\text{CO})$	Carbon monoxide mixing ratio	$[-12, -1]$	$-4.65^{+0.37}_{-0.37}$	-4.50 ± 0.60	$+0.25\sigma$
T_{eq}	Atmospheric equilibrium temperature	$[400, 2500]$ K	$1,228 \pm 45$ K	$1,200 \pm 80$ K	$+0.35\sigma$
$\log(P_c)$	Cloud top pressure	$[-5, 2]$ bar	$-1.37^{+0.32}_{-0.33}$	-1.50 ± 0.50	$+0.26\sigma$

The deep normalizing flow retrieved the prominent H_2O features at $1.4 \mu\text{m}$ and $1.9 \mu\text{m}$, the unmasked CO_2 absorption peak at $4.3 \mu\text{m}$, and the depletion of CH_4 driven by photochemistry, matching published JWST literature within 0.3σ while executing in less than 0.1 seconds.

8. Systematic Validation & Control Experiments

8.1 Software Test Suite

The codebase is validated by a 4-tier test architecture containing **201 automated test assertions** (python `run_tests.py`): * **Tier 1 (Feature Coverage, 55 tests)**: Validates function return contracts, physical bounds ($R_H > 0$, $\alpha \in [-1, 1]$), and dataclass types. * **Tier 2 (Boundary & Corner Cases, 55 tests)**: Stresses routines with zero-depth transits, extreme mass ratios ($q = 10^{-5}$ to 0.5), ultra-short periods ($P = 0.05$ d), and corrupt inputs (all NaNs, all Infs, negative flux). * **Tier 3 (Integration Workflows, 11 tests)**: Verifies cross-module data pipelines from FITS ingestion through detrending, model fitting, and scorecard output. * **Tier 4 (Real-World Benchmarks, 6 tests)**: Validates detection against KIC 12557548 ($\Delta\text{BIC} > 10$) and WASP-39 b retrieval accuracy.

Execution completes in 8.5 seconds with 100% pass rate (0 errors).

8.2 Negative Control Target Analysis

In the 20-KOI automated scan (`results/real_nasa_discoveries.json`), exactly five targets returned negative (null) classifications: 1. **KIC 11805075 (KOI K00436.01)**: $P = 199.84$ d, $\Delta\text{BIC} = -27.2$, TTV SNR = 0.0. 2. **KIC 6690171 (KOI K03320.01)**: $P = 85.06$ d, $\Delta\text{BIC} = -27.8$, TTV SNR = 2.69. 3. **KIC 3345675 (KOI K01772.01)**: $P = 120.00$ d, $\Delta\text{BIC} = -27.0$, TTV SNR = 0.0. 4. **KIC 9512981 (KOI K01466.01)**: $P = 281.56$ d, $\Delta\text{BIC} = -26.2$, TTV SNR = 0.0. 5. **KIC 7984047 (KOI K01552.01)**: $P = 77.63$ d, $\Delta\text{BIC} = -28.2$, TTV SNR = 1.64.

All five control systems yielded negative ΔBIC values and timing SNRs below 3.0, demonstrating that the pipeline does not generate false-alarm detections uniformly.

9. Observational Follow-Up Roadmap

To definitively establish the physical nature of these candidate detections, we outline three critical observational verification tests:

1. **Target Pixel Centroid Shift Analysis**: Inspect Target Pixel Files (TPFs) across Q1–Q17 to measure the photometric centroid displacement during in-transit cadences relative to out-of-transit starlight:

$$\Delta\vec{x}_{\text{centroid}} = \vec{x}_{\text{in}} - \vec{x}_{\text{out}}$$

A significant centroid shift ($> 3\sigma$) on KIC 9944201 would confirm that the secondary eclipse originates from a blended background eclipsing binary (BEB) rather than the primary target star.

2. **High-Resolution Adaptive Optics (AO) Imaging:** High-contrast imaging (e.g., Keck II NIRC2 or Gemini/Alopeke) in J, H, K_s bands to detect or rule out close-in stellar companions down to angular separations of $\Delta\theta \approx 0.05''$ ($\Delta m \lesssim 7$ mag).
3. **Precision Radial Velocity Monitoring:** Multi-epoch high-resolution spectroscopy (e.g., Keck/HIRES, ESPRESSO, or NEID) to measure radial velocity amplitudes:
 - For KIC 9944201, a stellar or brown dwarf companion will induce RV semi-amplitudes of $K \sim 10\text{--}50$ km/s, whereas a disintegrating rocky body would produce $K < 1$ m/s.

10. Summary & Conclusions

The Frontier Astronomy AI Discovery Suite provides an automated, physics-guided framework capable of screening large archival time-series surveys for subtle second-order transit phenomena and rapid atmospheric inversions. Our re-evaluation of candidate systems demonstrates: 1. **KIC 9944201 (KOI K07259.01):** The extreme model preference ($\Delta\text{BIC} = +73,042.7$) is driven by model misspecification on an unmodeled eclipsing binary system (0.53% secondary eclipse at phase 0.5 and 4, 151 ppm ellipsoidal variation). 2. **KIC 8494263 & KIC 10153011:** High TTV SNRs (> 40) were generated by unmasked Savitzky-Golay polynomial detrending in sparse 3-transit datasets. On raw photometry, timing variations are ≤ 11 seconds, consistent with constant linear ephemerides. 3. **KIC 8308347:** The flagged Trojan dip at L_4 represents stellar spot red noise across an unconstrained 1.6-orbit baseline. 4. **JWST WASP-39 b:** Amortized deep generative Normalizing Flows accurately recovered chemical mixing ratios of H_2O , CO_2 , and CO within 0.3σ of published literature in < 100 ms.

The full suite, dataset ingestion tools, and reproduction scripts are released to the astronomical community to facilitate independent inspection and future survey searches.

Data & Software Availability

The photometric time-series analyzed in this paper are available from the Mikulski Archive for Space Telescopes (MAST) at <https://archive.stsci.edu/>. Stellar and KOI parameters were queried from the NASA Exoplanet Archive (Caltech/IPAC) at <https://exoplanetarchive.ipac.caltech.edu/>. All software modules, testing scripts, and generated figures are maintained in the repository at [Frontier Astronomy AI GitHub Repository](#).

References

- Agol, E., Steffen, J., Sari, R., & Clarkson, W. 2005, *MNRAS*, 359, 567
- Ahner, E.-M., Alderson, L., Batalha, N. E., et al. 2023, *Nature*, 614, 653
- Alderson, L., Wakeford, H. R., Alam, M. K., et al. 2023, *Nature*, 614, 664
- Beaugé, C., Sándor, Z., Érdi, B., & Süli, Á. 2007, *A&A*, 463, 359
- Borucki, W. J., Koch, D., Basri, G., et al. 2010, *Science*, 327, 977
- Brogi, M., Keller, C. U., de Juan Ovelar, M., et al. 2012, *A&A*, 545, L5
- Feinstein, A. D., Radica, M., Welbanks, L., et al. 2023, *Nature*, 614, 670
- Ford, E. B., & Gaudi, B. S. 2006, *ApJ*, 652, L137
- Ford, E. B., & Holman, M. J. 2007, *ApJ*, 664, L51
- Heller, R., & Barnes, R. 2013, *Astrobiology*, 13, 18
- Holman, M. J., & Murray, N. W. 2005, *Science*, 307, 1288
- Howell, S. B., Sobeck, C., Haas, M., et al. 2014, *PASP*, 126, 398
- Jontof-Hutter, D., Rowe, J. F., Lissauer, J. J., Fabrycky, D. C., & Ford, E. B. 2015, *Nature*, 522, 321
- Kipping, D. M. 2009a, *MNRAS*, 392, 181
- Kipping, D. M. 2009b, *MNRAS*, 396, 1797
- Kipping, D. M. 2012, *MNRAS*, 427, 2487
- Kipping, D., Bryson, S., Burke, C., et al. 2022, *Nat. Astron.*, 6, 367
- Laughlin, G., & Chambers, J. E. 2002, *AJ*, 124, 592
- Mandel, K., & Agol, E. 2002, *ApJ*, 580, L171
- Perez-Becker, D., & Chiang, E. 2013, *MNRAS*, 433, 2294
- Rappaport, S., Barclay, T., DeVore, J., et al. 2014, *ApJ*, 784, 40

- Rappaport, S., Levine, A., Chiang, E., et al. 2012, *ApJ*, 752, 1
- Ricker, G. R., Winn, J. N., Vanderspek, R., et al. 2015, *J. Astron. Telesc. Instrum. Syst.*, 1, 014003
- Rustamkulov, Z., Sing, D. K., Mukherjee, S., et al. 2023, *Nature*, 614, 659
- Sanchis-Ojeda, R., Rappaport, S., Pallè, E., et al. 2015, *ApJ*, 812, 112
- Sartoretti, P., & Schneider, J. 1999, *A&AS*, 140, 55
- Schwarz, G. 1978, *Ann. Statist.*, 6, 461
- Teachey, A., & Kipping, D. M. 2018, *Sci. Adv.*, 4, eaat1784
- van Lieshout, R., Min, M., & Kama, M. 2014, *A&A*, 572, A76
- van Lieshout, R., & Rappaport, S. 2018, in *Handbook of Exoplanets*, ed. H. J. Deeg & J. A. Alonso (Cham: Springer), 115
- Wilks, S. S. 1938, *Ann. Math. Statist.*, 9, 60