

Predator and Prey Model

Mustafif Khan

MTH630-W2024

March 31, 2024

Toronto
Metropolitan
University

Outline

- 1 Lotka-Volterra Model
- 2 Dimensionless Equations
- 3 Equilibrium Points
- 4 Stability Analysis

Lotka-Volterra Model

The Lotka-Volterra Predator-Prey model consists of two first order non-linear differential equations. The model describes the interaction between two species, where one species is the predator y and the other is the prey x .

$$\frac{dx}{dt} = ax - bxy = x(a - by) \quad (1)$$

$$\frac{dy}{dt} = -cy + exy = y(-c + ex) \quad (2)$$

where a , b , c , and e are ≥ 0 .

Understanding the Parameters

Inside the equations, the parameters have the following meanings:

- x : the population of the prey species (e.g. rabbits)
- y : the population of the predator species (e.g. foxes)
- a : the natural growth rate of prey in the absence of predation
- b : the death rate per encounter of prey due to predation
- c : the natural death rate of the predator in the absence of prey
- e : the efficiency of turning prey into predator offspring

Dimensionless Equations

To simplify the equations, we can make them dimensionless.

$$u(\tau) = \frac{ex(t)}{c}$$

$$v(\tau) = -\frac{by(t)}{a}$$

$$\tau = at$$

Then we have the two following differential equations:

$$\frac{du}{d\tau} = u(1 - v) \quad (3)$$

$$\frac{dv}{d\tau} = \alpha v(u - 1) \quad (4)$$

Where $\alpha = c/a$.

Equilibrium Points

The equilibrium points of the system are found by setting the right hand side of the equations to zero. This gives us the following equilibrium points, $(0,0)$ and $(1,1)$.

The Jacobian matrix of the system is given by:

$$J(u, v) = \begin{bmatrix} 1 - v & -u \\ \alpha v & \alpha(u - 1) \end{bmatrix}$$

The eigenvalues of the Jacobian matrix at the equilibrium points are

$$J(0,0) = \begin{bmatrix} 1 & 0 \\ 0 & -\alpha \end{bmatrix} \text{ and } J(1,1) = \begin{bmatrix} 0 & -1 \\ \alpha & 0 \end{bmatrix}$$

The eigenvalues of $J(0,0)$ are 1 and $-\alpha$, and the eigenvalues of $J(1,1)$ are $\pm i\sqrt{\alpha}$.

Stability Analysis

Since the eigenvalues of $J(0,0)$ are both real and that $\lambda_1\lambda_2 < 0$, the equilibrium point $(0,0)$ is a saddle point.

Since both of the eigenvalues of $J(1,1)$ are complex, the equilibrium point $(1,1)$ is a center, or stable/unstable spiral.

- The equilibrium point $(0,0)$ represents when both species are extinct.
- The equilibrium point $(1,1)$ represents when both species coexist in time.
- A center can lead to a regular, cyclic behavior in both populations that oscillate around the equilibrium point in a stable manner.
- A stable spiral indicates that there can be a perturbation in one of the populations, but they will eventually return to the equilibrium state.
- An unstable spiral suggests that the populations can experience population collapses or rapid growth due to a small perturbation.

Consider the following example, where we have the following values for the constants: $a = 0.1$, $b = 0.02$, $c = 0.2$, and $e = 0.01$. We then get the following dimensionless equations:

$$\begin{aligned}\frac{du}{d\tau} &= u(1 - v) \\ \frac{dv}{d\tau} &= 2v(u - 1)\end{aligned}$$

This has the following Jacobian matrices at the equilibrium points:

$$J(0,0) = \begin{bmatrix} 1 & 0 \\ 0 & -2 \end{bmatrix} \quad J(1,1) = \begin{bmatrix} 0 & -1 \\ 2 & 0 \end{bmatrix}$$

The Jacobian matrix at $(0, 0)$ has eigenvalues 1 and -2 , and the has the following phase plane analysis:

- Trace $= 1 - 2 = -1$
- Determinant $= 1 \cdot -2 = -2$
- Discriminant $= (-1)^2 - 4(-2) = 1 + 8 = 9$

Since the discriminant is positive and the determinant is negative, the equilibrium point $(0, 0)$ is a saddle point.

The Jacobian matrix at $(1, 1)$ has eigenvalues $\pm i\sqrt{2}$, and the phase plane analysis is as follows:

- Trace = 0
- Determinant = $0 - (-2) = 2$
- Discriminant = $0^2 - 8 = -8$

Since the discriminant is negative and the trace is zero, the equilibrium point $(1, 1)$ is a center.