

1. **BO2025** Compute

$$\frac{20}{2! + 0} + \frac{24}{2! + 4}$$

Written by: Benjamin Oh

Answer: 14.

Simplify to $10 + 4$

2. **PS1985** Six people are in a line, including Adam and Ben. Given that Adam is third in line, then what is the probability that Ben is next to him?

Written by: Pranshu Sharma

Answer: $\frac{2}{5}$.

There are 5 available slots for Ben, and 2 of them are next to Adam. So, the chances of Ben being next to Adam is $\frac{2}{5}$.

3. **NA1942** A triangle has side lengths $2x$, $3x$, and $4x$. If the perimeter of the triangle is equal to 18cm, find the length of the longest side in centimeters.

Written by: Nicholas Andrison

Answer: 8.

The perimeter is equal to $2x + 3x + 4x = 9x$, leaving us with the equation $9x = 18\text{cm}$, or $x = 2\text{cm}$. The longest side is $4x$, so its length is $2 \times 4 = \textspan style="border: 1px solid black; padding: 0 5px;">8 centimeters.$

4. **TY2061** Joseph received 62 pieces of candy from trick-or-treating at Halloween. He gives the largest possible number of candies to his 12 friends, such that each friend gets the same amount, and Joseph still has more candy than any one of them. How many candy pieces does Joseph have now?

Written by: Troy Yang

Answer: 14

$62 \div 12 = 5\text{R}2$, meaning if Joseph gives 5 candies to each friend, he will only have 2 left. So instead, he gives 4 candies to each friend, and he's left with $62 - 12 \cdot 4 = \textspan style="border: 1px solid black; padding: 0 5px;">14 pieces of candy.$

5. **TY2055** Harrison's closet contains a red, a blue, and a black shirt; a pair of black pants and a pair of white pants; and a pair of black shoes and a pair of white shoes. If Harrison wears 1 shirt, 1 pair of pants, and 1 pair of shoes, how many outfits can he

make if he doesn't want to wear all black?

Written by: Troy Yang

Answer: 11

Harrison has 3 shirts, 2 pairs of pants, and 2 pairs of shoes he can choose from. This gives $3 \cdot 2 \cdot 2 = 12$ outfits, but the all-black outfit doesn't work, giving $12 - 1 = \boxed{11}$ good outfits.

6. **RS1971** John has a glass of orange juice that is $\frac{5}{8}$ full. He pours all of it into another glass and only ends up filling it halfway. The second glass has a capacity of 40 fluid ounces. What is the total capacity of the first glass in fluid ounces?

Written by: Rohan Singh

Answer: 32.

Since the second glass has a capacity of 40 fl. oz., and it is halfway full, the volume of orange juice must be 20 fl. oz.. This is the same amount that filled the first glass $\frac{5}{8}$ full. If V is the volume of the first glass, then $\frac{5}{8} \cdot V = 20$, so $V = 32$.

7. **KV1883** What is the largest number of sides a regular polygon could have such that it can be inscribed in a square?

Written by: Keshav Venkatesh

Answer: 8.

We can see that no side of the square can contain more than two vertices of the polygon, so the polygon can have at most 8 sides. When we draw what a regular octagon should look like inscribed in a square, we can see that the four sides of the squares obviously have equal length. Therefore, our answer is 8.

8. **AP2016** Lucy has a bag of marbles containing red, blue, and green marbles. The ratio of red to blue marbles is $3 : 2$, and the ratio of blue to green marbles is $4 : 5$. If Lucy has 45 marbles in total, how many green marbles does she have?

Written by: Aratrika Pal

Answer: 15.

For every 5 green marbles, there are 4 blue marbles and 6 red marbles. The total amount of marbles is 15. However, Lucy has 45 marbles in total, so we multiply all the values by 3, to get 15 green marbles total.

9. **JK1954** Triangle ABC is a right triangle with $AB = 2$ and $\angle BCA = 30^\circ$. Find the product of the side lengths of triangle ABC .

Written by: Joshua Koo

Answer: $\boxed{16\sqrt{3}}$

It should be obvious that triangle ABC is a $30 - 60 - 90$ right triangle. Since $\angle BCA = 30^\circ$, AB (the opposite side to $\angle BCA$) must be the smallest side length of the triangle. It quickly follows that the two other sides of the triangle have length $2\sqrt{3}$ and 4. The product of these three side lengths is $\boxed{16\sqrt{3}}$.

10. **AZ1781** Jax has 10 pyramids, 2 cubes, 4 octahedrons, 12 cylinders, and 15 spheres. He places all of these objects in a drawer and draws an object randomly. If a is the minimum number of draws to guarantee that Jax has all of at least 1 type of object, and b is the minimum number of draws to guarantee that he has all 12 cylinders, what is $\frac{a}{b}$?

Written by: Alexander Zhao

Answer: $\boxed{\frac{39}{43}}$

First, if there are n of the same type of object, it's possible for Jax to draw $n - 1$ of these objects before he will have all of them. So, the minimum amount of draws would be $(10 - 1) + (2 - 1) + (4 - 1) + (12 - 1) + (15 - 1) + 1 = 39$. The minimum amount of draws to guarantee to have all the cylinders is $10 + 2 + 4 + 12 + 15 = 43$.

The answer is $\boxed{\frac{39}{43}}$.

11. **AP2019** A rectangular garden is 15 feet long and 10 feet wide. If a path of uniform width is built around the garden on all four sides, increasing the total area to 300 square feet, what is the width of the path as a decimal in feet?

Written by: Aratrika Pal

Answer: $\boxed{2.5}$

Let the width of the path be x feet. The original area of the garden is $15 \times 10 = 150$ square feet. When the path is added, the new length of the garden including the path becomes $15 + 2x$ feet, and the new width becomes $10 + 2x$ feet. The new area of the garden including the path is $(15 + 2x)(10 + 2x) = 300$. Expanding this equation: $150 + 20x + 30x + 4x^2 = 300$. Simplifying: $4x^2 + 50x - 150 = 0$. Solving for x , we get $x = 2.5$ or $x = -15$. We can disregard the second solution, since distance must be positive. The width of the path, therefore, is 2.5 feet.

12. **KV1991** How many regular polygons have an integer value for their angle measure?

Written by: Keshav Venkatesh

Answer: $\boxed{22}$

If the interior angle measure of a regular polygon is an integer, then its exterior angle measure is also an integer. Since the exterior angle measure of an n -gon is $\frac{360}{n}$, we have $\frac{360}{n} = k$, where k is the exterior angle measure. Since $0 < k < 180$, all the factors of 360 besides 1 and 2 are valid values for n . Therefore, our answer is $\boxed{22}$.


13. **NA1978** Find the units digit of $2^{2024} \times 3^{2024^{2024}} \times 7^{2024^{2024^{2024}}}$.

Written by: Nicholas Andrison

Answer: $\boxed{6}$.

The last digit of the powers of 2, 3, and 7 all repeat in a cycle that depends on the remainder of the power when divided by 4. Since 2024 is divisible by 4, it will always have a remainder of 0 no matter what its power is. Using the pattern cycles, we can conclude that the last digit of 2^{2024} is 6, the last digit of $3^{2024^{2024}}$ is 1, and the last digit of $7^{2024^{2024^{2024}}}$ is 1. Multiplying these numbers gives us 6, so the last digit of the original number is $\boxed{6}$.

14. **TY1889** A Pac-Man ghost is made by joining semicircle $\widehat{AB}$ of radius 3 with rectangle $ABCD$ with 3 congruent isosceles triangles cut out, as shown below. Given that $BC = 5$ and $CE = \sqrt{5}$, find the area of the ghost in terms of π .



Written by: Troy Yang

Answer: $\boxed{\frac{9\pi}{2} + 24}$

Let P be the foot of the altitude from E to $\overline{CD}$. From rectangle $ABCD$, $CP \cdot 6 = 3 \cdot 2$,

so $CP = 1$. By the Pythagorean Theorem, $EP = \sqrt{\sqrt{5}^2 - 1^2} = 2$. Thus, the area of each isosceles triangle is $\frac{1}{2} \cdot (2 \cdot CP) \cdot EP = 2$. The area of the ghost is then given by $\frac{1}{2} \cdot 3^2 \cdot \pi + 6 \cdot 5 - 3 \cdot 2 = \boxed{\frac{9\pi}{2} + 24}$.

15. **DS1780** At Mustang High, there are 3 different grade levels (10th to 12th) with an equal number of students in each grade. At the start of every year, around 10% of the students from each grade level receive a trophy that they keep throughout high school. Assume that trophy-winners are chosen uniformly and randomly, and note that a student may have more than one. After 5 years have passed, what percent of the entire school population is expected to have at least one trophy? The school population does not include those that have graduated.

Written by: Dev Saxena

Answer: $\boxed{18.7}$

We can split this up into each grade level. First off, only 10% of the 10th graders received a trophy. The 11th graders had two different years in which they could have earned a trophy. The first year, around 10% of them received a trophy, and the second year, another 10% received one. However, we must subtract off people who have received more than one trophy. This occurs with around 1% of people (since there is a 10% chance the first year and a 10% chance the second year). Therefore, around $10\% + 10\% - 1\% = 19\%$ of the 11th graders have a trophy. Finally, we replicate the process for 12th graders. Around 19% of them have a trophy from the first two years, and another $10\% - (10\% \cdot 19\%) = 8.1\%$ received one their third year.

To summarize, 10% of the 10th, 19% of the 11th, and 27.1% of the 12th graders have a trophy. Since there are an equal number of students per grade, we average these values to get our answer, which is $\boxed{18.7\%}$.

16. **EX1877** Elaine starts with 100,000 strands of hair, each with a mass of 1 mg. In her quest for more massive hair, she purchases a magic brush. Each stroke of the brush through her hair adds 0.1 mg to each strand but causes 1000 strands to fall. How many times should she run the brush through her hair to achieve the most massive hair?

Written by: Elaine Xu

Answer: $\boxed{45}$

Let n be the number of times she brushes her hair. The mass of each hair is $1 + 0.1n$ and the number of hairs is $100,000 - 1000n$. Thus the total mass is $(1 + 0.1n)(100,000 - 1000n)$, which has maximum at $n = \boxed{45}$.

17. **AM1945** Say that the values of x and y for which the expression $x^2 + y^2 - 6x + 14y - 40$ is at it's minimal value (for real x and y) is for $x = a$ and $y = b$. What is $a + b$?

Written by: Anastasia Makarevich

Answer: $\boxed{-4.}$

The expression $x^2 + y^2 - 6x + 14y - 40$ can be rewritten as $(x^2 - 6x + 9) - 9 + (y^2 + 14y + 49) - 49 - 40$, or $(x - 3)^2 + (y + 7)^2 - 98$. Now, the first 2 parts of the expression, namely $(x - 3)^2$ and $(y + 7)^2$ are clearly perfect squares. Since we are only working with real numbers, the minimum value any perfect square can be is 0. Therefore, we want for both $(x - 3)^2$ and $(y + 7)^2$ to equal 0. This means that both $x - 3 = 0$ and $y + 7 = 0$. Solving for x and y , we get that $x = 3$ and $y = -7$. Therefore, $a + b = 3 + (-7) = 3 - 7 = \boxed{-4.}$

18. **SM1882** Positive integers x and y satisfy the equation

$$\frac{15}{3+x} = \frac{12+y}{7}$$

Find the sum of all possible values of xy .

Written by: Sergei Makarevich

Answer: $\boxed{30.}$

We get that $(x + 3)(y + 12) = 3 * 5 * 7$, since x, y , are positive integers, then we get that $(x, y) = (2, 9), (x, y) = (4, 3)$. This gives us $18 + 12 = 30$.

19. **TY1743** Jack and Jill are playing a not-so-fair game. They have a deck of 9 cards, each card numbered 1 through 9. Jack randomly takes 6 cards from the deck and gives the other 3 to Jill. Jack wins if the sum of his cards is greater than the sum of Jill's cards, and loses otherwise. How many different hands of 6 cards are winning for Jack?

Written by: Troy Yang

Answer: $\boxed{82}$

Jack can have a total of $\binom{9}{6} = 84$ different hands. Let us count the number of hands in which Jill wins, and subtract that from 84 to find how many hands Jack wins. The sum of all cards in the deck is equal to $1 + 2 + \dots + 9 = \frac{9 \cdot 10}{2} = 45$. Note that 45 is odd, so Jack and Jill cannot tie. If Jill has a sum of x , then Jack has a sum of $45 - x$. For Jill to win, $x > 45 - x \Rightarrow x > 22.5$. Jill's biggest possible hand is $(7, 8, 9)$, which has a sum of 24. The only other possible hand is $(6, 8, 9)$ since lowering the value of any of these 3 cards will result in a sum less than 22.5. Thus, Jill has 2 winning hands, and Jack has $84 - 2 = \boxed{82}$ winning hands.

20. **TL1880** What are the last two digits of 131^{45} ?

Written by: Tianlin Liu

Answer: $\boxed{51.}$

Using the binomial theorem and expanding, we obtain $(130 + 1)^{45} = 130^{45} + 45C1 * 130^{44} * 1^1 + 45C2 * 130^{43} * 1^2 + \dots + 45C2 * 130^2 * 1^{43} + 45C1 * 130^1 * 1^{44} + 1^{45}$. We see that we only need to consider the last 2 terms, since all other terms end with at least

2 zeros. The last two terms are $45C1 * 30^1 * 1^{44} + 1^{45}$, the first term ends with 50 and the second term ends with 1. Therefore, our final answer is 51.

21. **EL2057** Alice and Bob are blowing bubbles together. Alice can blow 2 bubbles per second for 10 seconds, then catches her breath for 2 seconds before continuing. Her bubbles last 15 seconds before popping. Bob can blow 3 bubbles per second for 9 seconds, then catches his breath for 6 seconds. His bubbles last 16 seconds before popping. If Alice and Bob start blowing bubbles at the same time, after 2 minutes, what is the positive difference between the number of Alice's bubbles and the number of Bob's bubbles still in the air?

Written by: Emma Li

Answer: 5

Notice that we only need to worry about the bubbles that were blown 15 seconds before time for Alice and 16 seconds before for Bob, as any bubbles blown before that would have popped already.

For Alice, her bubbles are blown in 'cycles' of $10 + 2 = 12$ seconds, with $2 * 10 = 20$ bubbles blown per cycle. Going 15 seconds backwards, she can fit in a full cycle and the last three seconds of another. Two of those seconds will be spent resting, but she still blows 2 bubbles in the third second. Thus, she blows $20 + 2 = 22$ bubbles that will still be in the air.

For Bob, his bubbles are blown in 'cycles' of $9 + 6 = 15$ seconds, with $3 * 9 = 27$ bubbles blown per cycle. Going 16 seconds backwards, he can fit in a full cycle and the last second of another, of which will be spent resting. Thus, he blows 27 bubbles that will still be in the air.

Then, the positive difference between them is $|22 - 27| =$ 5 $$ bubbles.

22. **TY1907** In a standard tic-tac-toe game, 5 moves have been made, i.e. there are 3 X's and 2 O's on the grid. How many board configurations exist where neither player has won yet?


Written by: Troy Yang

Answer: 1140

For the person playing X's, there are $\binom{9}{3} = 84$ ways to place the 3 X's. However, 8 of these do not work because they form 3 in a row; specifically, there are the 3 horizontal lines, 3 vertical lines, and 2 diagonals. Thus, the X-player has $84 - 8 = 76$ valid placements. The person playing O's must place 2 O's in the remaining 6 squares, giving $\binom{6}{2} = 15$ valid placements. Then, there are $76 \cdot 15 =$ 1140 $$ total board configurations.

23. **LT279** Refer to the diagram below. The rectangle has a width of 16 and a height of 18. Two congruent circles, ω_1 and ω_2 , are tangent to each other, and each is tangent to two sides of the rectangle. A third circle, ω_3 , is tangent to ω_1 and to two sides of the

rectangle. Find the radius of ω_3 .



Written by: Linus Tang

Answer: $\boxed{21 - 8\sqrt{5}}$

Let the rectangle have opposite vertices $(0,0)$ and $(16,18)$ with sides parallel to the coordinate axes, and define O_1, O_2 , and O_3 to be the centers of the circles. Let R be the common radius of ω_1 and ω_2 and r be the radius of ω_3 . We can first solve for R : we know that $O_1 = (R, R)$ and $O_2 = (16 - R, 18 - R)$. Since circles ω_1 and ω_2 are tangent to each other, the distance between O_1 and O_2 is $2R$. Using the Euclidean distance formula or the Pythagorean Theorem, $(16 - 2R)^2 + (18 - 2R)^2 = (2R)^2$. This can be expanded to give a quadratic in R whose positive solutions are $R = 5$ and $R = 29$. The latter solution is obviously too large, so we have $R = 5$.

Now, we turn our attention to tangent circles ω_1 and ω_3 . We know that the distance between O_1 and O_3 is $R + r$. Furthermore, $O_1 = (R, R)$ and $O_3 = (16 - r, r)$. Again, using the Euclidean distance formula, $(16 - r - R)^2 + (r - R)^2 = (R + r)^2$. $16 - r - R = 2\sqrt{rR}$ $R + 2\sqrt{rR} + r = 16$ $(\sqrt{R} + \sqrt{r})^2 = 16$ $\sqrt{R} + \sqrt{r} = 4$ Thus, $\sqrt{r} = 4 - \sqrt{R} = 4 - \sqrt{5}$. The answer is $r = (4 - \sqrt{5})^2 = \boxed{21 - 8\sqrt{5}}$.

24. **RRP2021** Let $f(x) = x^4 - 2x^3 + 8x^2 + x + 2$. Suppose $f(x)$ has roots r_1, r_2, r_3, r_4 . Compute

$$(r_1 + r_2 + r_3)(r_1 + r_2 + r_4)(r_1 + r_3 + r_4)(r_2 + r_3 + r_4).$$

Written by: Ryan Pascual

Answer: $\boxed{2}$

Note $r_1 + r_2 + r_3 = (r_1 + r_2 + r_3 + r_4) - r_4 = 2 - r_4$ by Vieta's. Then, the desired product equals $(2 - r_4)(2 - r_3)(2 - r_2)(2 - r_1) = f(2) = 36$.

25. **RT1732** A four digit number $\overline{abcd}$ is suspicious if $\overline{da} + \overline{bc} = \overline{ab} + \overline{cd}$. How many suspicious numbers are there?

Written by: Ryan Tang

Answer: 615

Note that a suspicious number satisfies $10d + a + 10b + c = 10a + b + 10c + d$. Simplifying this gives $a + c = d + b$. Let this common sum be k .

For $k \leq 9$, the number of ways to choose a and c is k and the number of ways to choose b and d is $k + 1$, so the total ways is $k(k + 1) = 2\binom{k+1}{2}$. The sum of this from $k = 1$ to 9 is $2\binom{11}{3} = 330$ by hockey stick.

For $k \geq 10$, the number of ways to choose a and c is $9 - (k - 9) + 1 = 19 - k$ and the number of ways to choose b and d is also $19 - k$, so the total ways is $(19 - k)^2$. The sum of this from 10 to 18 is equal to $1^2 + 2^2 + \cdots + 9^2 = \frac{9 \cdot 10 \cdot 19}{6} = 285$

The sum of these numbers equals to $330 + 285 =$ 615