

Solution homework exercise 3.4:

Statement: For

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}, f(x, y) = x^2 - 2xy + y^2$$

are all $(x, x)^T$ with $x \in \mathbb{R}$ nonstrict global minima. Other minima or maxima do not exist.

Proof: Assume that f has a minimum or a maximum $(\hat{x}, \hat{y})^T \in \mathbb{R}^2$. Due to the necessary first order optimality condition,

$$\nabla f(\hat{x}, \hat{y}) = 0.$$

For all $(x, y)^T \in \mathbb{R}^2$,

$$\nabla f(x, y) = \begin{pmatrix} 2x - 2y \\ -2x + 2y \end{pmatrix}.$$

Thus it follows

$$\hat{x} = \hat{y}.$$

Furthermore

$$\nabla^2 f(x, y) = \begin{pmatrix} 2 & -2 \\ -2 & 2 \end{pmatrix}$$

for all $(x, y)^T \in \mathbb{R}^2$. Since

$$\frac{\partial^2 f}{\partial x^2}(x, y) = 2 \geq 0,$$

$$\frac{\partial^2 f}{\partial y^2}(x, y) = 2 \geq 0$$

and

$$\det \left(\frac{\partial^2 f}{\partial x^2}(\hat{x}, \hat{y}) \right) = 2^2 - (-2)^2 = 0 \geq 0$$

for all $(x, y)^T \in \mathbb{R}^2$, it follows that $\nabla^2 f(x, y)^T$ is positive semidefinite for all $(x, y)^T \in \mathbb{R}^2$ due to Sylvester's criterion. Thus $(x, x)^T$ is a minimum of f for every $x \in \mathbb{R}$ due to the second order sufficient optimality condition. Since

$$f(x, x) = x^2 - 2xx + x^2 = 0 \leq (x - y)^2 = f(x, y)$$

for all $(x, y)^T \in \mathbb{R}^2$, all these minima are nonstrict global minima. □