


¹¹¹Cd (Cd I), nuclear spin $I = 1/2$. Strongest classified transitions below 2 μm (NIST): 15 levels, 21 lines.

Cd^+ ionisation limit $72540.05 \text{ cm}^{-1} = 8.99382 \text{ eV} (137.855 \text{ nm})$

Arrow label: vacuum wavelength | reduced dipole matrix element $\langle J \| er \| J' \rangle$ in ϵ_0
 Matrix elements are measured or NIST values; * = high-accuracy theory, $\approx$ = model calculation.
 Arrow width grows with the Einstein coefficient A. Dotted: forbidden (clock, M1, E2). Dashed: Rydberg excitation, $n = 70$.
 Level caption: energy (NIST, or a measurement of the last 20 years where one exists) and measured lifetime (where one exists).



Transition	λ vac (nm)	λ air (nm)	ν (THz)	$ \langle J e J' \rangle $ (ea ₀)	A (s ⁻¹)	$\Gamma/2\pi$ up. (MHz)	branch (%)	Use
$5s^2\ ^1S_0 - 5s5p\ ^1P^o_1$	228.8728	228.8023	1309.8647	3.331	6.250e+08	99.47	100	228.9 nm resonance line: first-stage MOT / slowing / imag...
$5s^2\ ^1S_0 - 5s5p\ ^3P^o_2$	314.1991	314.1081	954.1480	-	-	3.789e-09	-	314 nm ultra-narrow
$5s^2\ ^1S_0 - 5s5p\ ^3P^o_1$	326.1995	326.1055	919.0464	0.147	4.184e+05	0.06659	100	326.1 nm intercombination line: narrow-line MOT
$5s^2\ ^1S_0 - 5s5p\ ^3P^o_0$	332.0716	331.9760	902.7947	-	-	-	-	332 nm optical lattice clock transition
$5s5p\ ^3P^o_2 - 5s5d\ ^3D_3$	361.1538	361.0508	830.0965	4.600	1.300e+08	-	-	361.2 nm metastable MOT
$5s5p\ ^3P^o_2 - 5s5d\ ^3D_1$	361.5484	361.4453	829.1904	0.562*	4.510e+06	-	-	361.5 nm
$5s5p\ ^3P^o_1 - 5s6s\ ^3S_1$	480.1254	479.9912	624.4045	2.592	4.100e+07	-	-	480.0 nm optical pumping / repumping

λ , ν from NIST level energies unless a measured frequency for this isotope is available. A: measured value or NIST. $\Gamma/2\pi$: natural linewidth of the upper level, $1/(2\pi\tau)$.

Level	A (MHz)	B (MHz)	Shift of each F level from the centre of gravity (MHz)
5s5p $^3P_1^o$	-4123.813(13)	-	F=1/2: +4123.81 F=3/2: -2061.91
5s5p $^3P_2^o$	-3292.9364(8)	-	F=3/2: +4939.40 F=5/2: -3292.94
5s5p $^1P_1^o$	188.5(22)	-	F=1/2: -188.50 F=3/2: +94.25
5s6s 3S_1	-7750.8(9)	-	F=1/2: +7750.80 F=3/2: -3875.40
5s6d 3D_3	-2386.297(26)	-	F=5/2: +4772.59 F=7/2: -3579.45

Measured A, B constants; sources in the data file.

Isotope shifts			
Transition	λ vac (nm)	Isotopes	Shift (MHz)
$5s^2\ ^1S_0 - 5s5p\ ^1P_1^o$	228.873	106-114	1818(4)
$5s^2\ ^1S_0 - 5s5p\ ^1P_1^o$	228.873	108-114	1336(3)
$5s^2\ ^1S_0 - 5s5p\ ^1P_1^o$	228.873	110-114	865(3)
$5s^2\ ^1S_0 - 5s5p\ ^1P_1^o$	228.873	112-114	408(3)
$5s^2\ ^1S_0 - 5s5p\ ^1P_1^o$	228.873	116-114	-316(3)
$5s^2\ ^1S_0 - 5s5p\ ^1P_1^o$	228.873	111-114	805(0.3)
$5s^2\ ^1S_0 - 5s5p\ ^1P_1^o$	228.873	113-114	344.9(5)
$5s^2\ ^1S_0 - 5s5p\ ^3P_2^o$	314.199	106-114	1916(4)
$5s^2\ ^1S_0 - 5s5p\ ^3P_2^o$	314.199	108-114	1404(3)
$5s^2\ ^1S_0 - 5s5p\ ^3P_2^o$	314.199	110-114	913.7(21)
$5s^2\ ^1S_0 - 5s5p\ ^3P_2^o$	314.199	112-114	429.7(16)
$5s^2\ ^1S_0 - 5s5p\ ^3P_2^o$	314.199	116-114	-320.2(17)
$5s^2\ ^1S_0 - 5s5p\ ^3P_2^o$	326.200	106-114	1913.0(10)
$5s^2\ ^1S_0 - 5s5p\ ^3P_2^o$	326.200	108-114	1402.4(10)
$5s^2\ ^1S_0 - 5s5p\ ^3P_1^o$	326.200	110-114	914.7(10)
$5s^2\ ^1S_0 - 5s5p\ ^3P_1^o$	326.200	112-114	429.9(10)
$5s^2\ ^1S_0 - 5s5p\ ^3P_1^o$	326.200	116-114	-321.5(10)
$5s^2\ ^1S_0 - 5s5p\ ^3P_1^o$	326.200	111-114	861.6(10)
$5s^2\ ^1S_0 - 5s5p\ ^3P_1^o$	326.200	113-114	370.8(10)
$5s^2\ ^1S_0 - 5s5p\ ^3P_0^o$	332.072	106-114	1911(5)
$5s^2\ ^1S_0 - 5s5p\ ^3P_0^o$	332.072	108-114	1404(4)
$5s^2\ ^1S_0 - 5s5p\ ^3P_0^o$	332.072	110-114	914.1(21)

$$\text{Shift} = \nu(\text{first isotope}) - \nu(\text{second isotope})$$