

I (II). Strongest classified transitions below 2 μm (NIST): 29 levels, 50 lines.

$$\text{I}^+ \text{ ionisation limit } 84294.90 \text{ cm}^{-1} = 10.45124 \text{ eV (118.631 nm)}$$

Arrow label: vacuum wavelength | reduced dipole matrix element $\langle j || e r || j' \rangle$ in $e a_0$
 Matrix elements are measured or NIST values; * = high-accuracy theory, $\approx$ = model calculation.
 Arrow width grows with the Einstein coefficient A. Dotted: forbidden (clock, M1, E2). Dashed: Rydberg excitation, $n = 70$.
 Level caption: energy (NIST, or a measurement of the last 20 years where one exists) and measured lifetime (where one exists).



Transition	λ vac (nm)	λ air (nm)	ν (THz)	$\langle [J e J'] \rangle$ (ea)	A (s ⁻¹)	$\Gamma/2\pi$ up. (MHz)	branch (%)	Use
$2^1P_{3/2} - 2[1]_{3/2}^0$	149.1158	-	2010.4670(3)	-	-	4.483	-	TALIF detection of ground-state I: 2 x 298.2 nm, fluoresc...
$2^1P_{3/2}^0 - 2[3]_{3/2}^0$	152.2770	-	1968.7316(3)	-	-	-	-	TALIF of ground-state I near 2 x 304.5 nm, fluorescence o...
$2^1P_{3/2}^0 - 2[3]_{3/2}^0$	152.3361	-	1967.9669	-	-	-	-	classic TP-LIF /
$2^1P_{3/2}^0 - 2[2]_{3/2}^0$	153.3648	-	1954.7675	-	-	-	-	TP-LIF detection of spin-orbit excited I*
$2^1P_{3/2}^0 - 2[2]_{3/2}^0$	178.2763	-	1681.6172	1.770	2.800e+08	44.21	100	178.28 nm resonance line
$2^1P_{3/2}^0 - 2[2]_{3/2}^0$	183.0386	-	1637.8646	0.305	5.140e+06	0.8162	100	183.04 nm resonance line
$2[2]_{3/2} - 2[1]_{3/2}^0$	804.5905	804.3693	372.6025	2.498	6.070e+06	4.483	21.6	804 nm fluorescence line used to detect I atoms after two...
$2[2]_{3/2} - 2[3]_{3/2}^0$	906.0811	905.8325	330.8671	9.740	3.230e+07	-	-	905 nm fluorescence line: the only decay channel of 6p 2[...
$2[2]_{3/2} - 2[1]_{3/2}^0$	911.6391	911.3890	328.8499	6.054	2.450e+07	4.483	87	911 nm fluorescence channel of the TALIF upper level 6p 2...
$2^1P_{3/2}^0 - 2^1P_{1/2}^0$	1315.2744	1314.9147	227.931501(15)	-	8.000e+00	1.253e-06	100	1315 nm iodine-laser line

Level	A (MHz)	B (MHz)	Shift of each F level from the centre of gravity (MHz)			
$2P_{3/2}^o$	827.265(3)	1146.356(10)	F=1: -3540.69	F=2: -2803.25	F=3: -837.31	F=4: +3388.83
$2P_{1/2}^o$	6587(4)	-	F=2: -11527.25	F=3: +8233.75		
$2[2]_{5/2}$	1417.100	-1065.500	F=0: -13145.48	F=1: -11472.66	F=2: -8222.91	F=3: -3588.03 F=4: +2144.30 F=5: +8590.50
$2[3]_{7/2}^o$	586.700	-1104.400	F=1: -7192.02	F=2: -5703.07	F=3: -3587.99	F=4: -988.75 F=5: +1905.30 F=6: +4857.52

Measured A, B constants; sources in the data file.

λ , ν from NIST level energies unless a measured frequency for this isotope is available. A : measured value or NIST. $\Gamma/2\pi$: natural linewidth of the upper level, $1/(2\pi\tau)$.