


Ytterbium-173

¹⁷³Yb (Yb II), nuclear spin *I* = 5/2. Energy levels and transitions below 2 μm: 18 levels, 26 lines.

Yb + ionisation limit 50443.20 cm⁻¹ = 6.25416 eV (198.243 nm)



Key transitions

Transition	λ vac (nm)	λ air (nm)	ν (THz)	⟨J er J'⟩ (ea0)	A (s ⁻¹)	Γ/2π up. (MHz)	branch (%)	Use
6s² 1S ₀ - 6s7p 1P ₁ °	246.5242	246.4497	1216.0772	1.489	1.000e+08	-	-	246.5 nm
6s² 1S ₀ - (37415)1°	267.2754	267.1960	1121.6612	0.636	1.430e+07	-	-	267.3 nm
6s² 1S ₀ - 4f¹³5d6s² (7/2, 5/2)1°	346.5362	346.4370	865.1115	2.051	6.830e+07	-	-	346.5 nm
6s² 1S ₀ - 6s6p 1P ₁ °	398.9114	398.7986	751.5264	4.147	1.830e+08	29.13	100	399 nm: Zeeman slower, blue MOT, imaging
6s² 1S ₀ - 5d6s 3D ₂	404.0086	403.8945	742.0447	-	-	0.346	-	404 nm forbidden transition
6s² 1S ₀ - 5d6s 3D ₁	408.3449	408.2296	734.1648	-	-	0.4794	-	408 nm highly forbidden
6s² 1S ₀ - 4f¹³5d6s² (7/2, 3/2)2°	431.2479	431.1266	695.1743	-	-	2.411e-09	-	431 nm inner-shell clock transition
6s² 1S ₀ - 6s6p 3P ₂ °	507.3467	507.2053	590.9026	-	1.120e-01	1.788e-08	99.7	507 nm M2 transition: shelving into 3P2
6s² 1S ₀ - 6s6p 3P ₀ °	555.8024	555.6480	539.3868	0.540	1.146e+06	0.1824	100	556 nm: green MOT, narrow-line cooling, imaging in tweeze...
6s² 1S ₀ - 6s6p 3P ₀ °	578.4212	578.2609	518.2944	-	4.800e-02	7.579e-09	100	578 nm optical clock transition
6s6p 3P ₁ ° - 6s7s 1S ₀	611.2976	611.1285	490.4198	0.220	4.293e+05	3.475	1.97	611 nm
6s6p 3P ₀ ° - 6s7s 3S ₁	649.0871	648.9078	461.8678	1.951	9.400e+06	11.53	13	649 nm clock-state repumper / 3P0 detection
6s6p 3P ₁ ° - 6s7s 3S ₁	680.1479	679.9602	440.7754	3.480	2.600e+07	11.53	36	680 nm decay channel of 3S1 during repumping
6s6p 3P ₂ ° - 6s7s 3S ₁	770.1606	769.9487	389.2596	5.003	3.700e+07	11.53	50.8	770 nm 3P2 repumper
6s6p 3P ₁ ° - 5d6s 1D ₂	1032.4544	1032.1715	290.3687	0.540*	1.074e+05	0.02375	71.9	1032 nm
6s6p 1P ₁ ° - 6s7s 1S ₀	1077.3043	1077.0093	278.2802	4.310*	3.010e+07	3.475	100	1077 nm main decay channel of 6s7s 1S0
6s6p 3P ₂ ° - 5d6s 1D ₂	1255.1340	1254.7907	238.8530	0.380*	2.959e+04	0.02375	19.8	1255 nm
6s6p 1P ₁ ° - 6s7s 3S ₁	1311.2226	1310.8641	228.6358	0.730*	1.596e+05	11.53	0.21	1311 nm weak decay channel of 3S1
6s6p 3P ₀ ° - 5d6s 3D ₁	1388.7610	1388.3814	215.8705	2.759	1.920e+06	0.4794	63.8	1389 nm clock-state repumper
6s6p 3P ₁ ° - 5d6s 3D ₂	1479.3029	1478.8986	202.6579	4.030*	2.033e+06	0.346	93.5	1480 nm
6s6p 3P ₁ ° - 5d6s 3D ₁	1539.1494	1538.7289	194.7780	2.392	1.060e+06	0.4794	35.2	1539 nm decay channel of 3D1
6s6p 3P ₂ ° - 5d6s 3D ₃	1798.3949	1797.9039	166.7000	6.120*	1.864e+06	0.2947	100	1798 nm cycling transition from 3P2
6s6p 3P ₂ ° - 5d6s 3D ₂	1983.5130	1982.9717	151.1422	2.390*	2.966e+05	0.346	13.6	1983 nm


λ, ν from NIST level energies unless a measured frequency for this isotope is available. A: measured value or NIST.
Γ/2π: natural linewidth of the upper level, 1/(2πτ).

Isotope shifts

Transition	λ vac (nm)	Isotopes	Shift (MHz)
6s² 1S ₀ - 6s6p 1P ₁ °	398.911	168-174	1887.64(23)
6s² 1S ₀ - 6s6p 1P ₁ °	398.911	170-174	1187.8(27)
6s² 1S ₀ - 6s6p 1P ₁ °	398.911	171-174	940.8(9)
6s² 1S ₀ - 6s6p 1P ₁ °	398.911	172-174	533.0(3)
6s² 1S ₀ - 6s6p 1P ₁ °	398.911	173-174	291.51(5)
6s² 1S ₀ - 6s6p 1P ₁ °	398.911	176-174	-509.25(7)
6s² 1S ₀ - 5d6s 3D ₂	404.009	170-174	2531(5)
6s² 1S ₀ - 5d6s 3D ₂	404.009	171-174	2019(6)
6s² 1S ₀ - 5d6s 3D ₂	404.009	172-174	1100(4)
6s² 1S ₀ - 5d6s 3D ₂	404.009	173-174	609(6)
6s² 1S ₀ - 5d6s 3D ₂	404.009	176-174	-1059(11)
6s² 1S ₀ - 5d6s 3D ₁	408.345	170-174	2542(12)
6s² 1S ₀ - 5d6s 3D ₁	408.345	171-174	2038.9(12)
6s² 1S ₀ - 5d6s 3D ₁	408.345	172-174	1112(8)
6s² 1S ₀ - 5d6s 3D ₁	408.345	173-174	619.4(8)
6s² 1S ₀ - 5d6s 3D ₁	408.345	176-174	-1063(6)
6s² 1S ₀ - 4f¹³5d6s² (7/2, 3/2)2°	431.248	171-176	-3406.79(3)
6s² 1S ₀ - 4f¹³5d6s² (7/2, 3/2)2°	431.248	173-176	-1797.802(26)
6s² 1S ₀ - 4f¹³5d6s² (7/2, 3/2)2°	431.248	174-176	-1115.787(25)
6s² 1S ₀ - 4f¹³5d6s² (7/2, 3/2)2°	431.248	172-176	-2296.403(25)
6s² 1S ₀ - 4f¹³5d6s² (7/2, 3/2)2°	431.248	170-176	-3926.431(26)
6s² 1S ₀ - 4f¹³5d6s² (7/2, 3/2)2°	431.248	168-176	-5680.383(25)

Shift = ν(first isotope) - ν(second isotope).

Arrow label: vacuum wavelength | reduced dipole matrix element |⟨J||er||J'⟩| in ea₀
Matrix elements are measured or NIST values; * = high-accuracy theory, = model or semi-empirical calculation.
Arrow width grows with the Einstein coefficient A. Dotted: forbidden (clock, M1, E2). Dashed: Rydberg excitation, *n* = 70.
Level caption: energy (NIST, or a measurement of the last 20 years where one exists) and measured lifetime (where one exists).



¹⁷³Yb hyperfine structure (*I* = 5/2)

Level	A (MHz)	B (MHz)	Shift of each F level from the centre of gravity (MHz)
6s6p 3P ₁ °	-1094.361(11)	-826.35(8)	F=3/2: +3251.82 F=5/2: +1755.44 F=7/2: -2942.49
6s6p 3P ₀ °	-741.98(9)	1347.3(9)	F=1/2: +6136.97 F=3/2: +4417.72 F=5/2: +1889.12 F=7/2: -943.59 F=9/2: -3373.07
4f¹³5d6s² (7/2, 3/2)2°	-309.4491(20)	-1700.631(22)	F=1/2: +975.70 F=3/2: +1276.81 F=5/2: +1353.51 F=7/2: +568.04 F=9/2: -1972.40
5d6s 3D ₁	563.50(20)	336.8(3)	F=3/2: -1736.49 F=5/2: -832.94 F=7/2: +1492.95
6s6p 1P ₁ °	59.52(20)	601.9(5)	F=3/2: +212.99 F=5/2: -541.02 F=7/2: +299.27
6s7s 3S ₁	-1891.46(7)	-0.62(16)	F=3/2: +6619.68 F=5/2: +1891.96 F=7/2: -4728.81
6p² 3P ₂	-123.5(12)	-629.9(11)	F=1/2: +423.57 F=3/2: +521.77 F=5/2: +527.98 F=7/2: +205.96 F=9/2: -774.98

Measured A, B constants; sources in the data file.