


Te (Te I). Strongest classified transitions below 2 μm (NIST): 35 levels, 46 lines.

Te⁺ ionisation limit 72669.01 cm⁻¹ = 9.00981 eV (137.610 nm)

Arrow label: vacuum wavelength | reduced dipole matrix element $|U_{\text{erf}}|$ in ea_0
 Matrix elements are measured or NIST values; * = high-accuracy theory, $\approx$ = model calculation.
 Arrow width grows with the Einstein coefficient A . Dotted: forbidden (clock, M1, E2). Dashed: Rydberg excitation, $n = 70$.
 Level caption: energy (NIST, or a measurement of the last 20 years where one exists) and measured lifetime (where one exists).



Key transitions

Transition	λ vac (nm)	λ air (nm)	ν (THz)	$\langle U e U \rangle$ (eao)	A (s^{-1})	$\Gamma/2\pi$ up. (MHz)	branch (%)	Use
$5p^2\ ^3P_2 - 5p^36s\ ^3S_0^o$	214.3497	214.2822	1398.6139	2.133	$3.120e+08$	-	-	first step of the ISOLDE RILIS resonance-ionization scheme...
$5p^2\ ^3S_1^o - 5p^7p\ ^3P_1$	573.5175	573.3585	522.7259	-	-	1.941	-	second step of the ISOLDE RILIS resonance-ionization sche...

λ , ν from NIST level energies unless a measured frequency for this isotope is available. A: measured value or NIST. $\Gamma/2\pi$: natural linewidth of the upper level, $1/(2\pi\tau)$.