


Lewis Acid-Catalyzed Intermolecular Annulation: Three-Component Reaction toward Imidazo[1,2-*a*]pyridine ThionesZhengwang Chen,<sup>\*,</sup> Pei Liang, Fan Xu, Rulin Qiu, Qi Tan, Lipeng Long, and Min Ye<sup>\*</sup>

Key Laboratory of Organo-Pharmaceutical Chemistry of Jiangxi Province, Gannan Normal University, Ganzhou 341000, P. R. China

## Supporting Information

**ABSTRACT:** A Lewis acid-catalyzed three-component annulation reaction of 2-aminopyridines and ynals with elemental sulfur was established. A series of imidazo[1,2-*a*]pyridine thiones was obtained in moderate to excellent yields. The merits of this transformation include easily available starting materials, multiple C-heteroatom bond formation in one pot, good functional group tolerance, elemental sulfur as S source, operational simplicity, etc.



Imidazopyridines are a class of important nitrogen-fused heterocycles and have attracted considerable attention owing to their outstanding application value in natural products and pharmaceuticals.<sup>1</sup> Many imidazo[1,2-*a*]pyridine derivatives exhibit broad biological activities, including anticancer, antifungal, antibacterial, and antiviral activities.<sup>2</sup> Moreover, several commercially available drugs such as zolpidem, alpidem, olprinone, zolimidine, necopidem, and saripidem all contain the key structure of imidazo[1,2-*a*]pyridines.<sup>3</sup> As a consequence, numerous synthetic strategies have been developed for the construction of these skeletal structures over the past decades. Two regular methods involve the direct functionalization of pre-existing imidazo[1,2-*a*]pyridine compounds<sup>4</sup> and the establishment of this framework from acyclic starting materials.<sup>5</sup> Aminopyridines as key building blocks have been applied widely to couple with carbonyl compounds,<sup>6</sup> alkenes,<sup>7</sup> or alkynes<sup>8</sup> to construct structurally diverse imidazo[1,2-*a*]pyridines. Despite the significant advancement, the development of novel and powerful transformations for imidazo[1,2-*a*]pyridine synthesis is vital to the synthetic chemistry community.


Sulfur-containing molecules have found extensive applications in materials science and pharmaceutical chemistry.<sup>9</sup> Among the various synthetic strategies toward sulfur-containing compounds, the most attractive protocols involve the direct functionalization of the stable and nontoxic elemental sulfur.<sup>10</sup> Recently, some strategies have been established for the preparation of imidazo[1,2-*a*]pyridine derivatives based on three-component reaction from 2-aminopyridines, ynals, and coupling partners. These reagents include alcohols or thiols,<sup>11</sup> molecular oxygen,<sup>12</sup> and sodium benzenesulfonates<sup>13</sup> (Schemes 1a–c). Herein, we described the first direct synthesis of imidazo[1,2-*a*]pyridine thiones from 2-aminopyridines, aromatic ynals, and elemental sulfur catalyzed by Lewis acids (Scheme 1d).

Initially, we employed 2-aminopyridine **1a** and phenylpropionaldehyde **2a** as a model reaction to optimize the reaction conditions, and the results are shown in Table 1.

When the reaction was carried out without any catalyst and additive, the desired product **3a** was formed in 35% yield (entry 1). To further improve the reaction yield, various Lewis acid catalysts were tested, and all of the catalysts had positive effects on the transformation (entries 2–10). To our surprise, the rare-earth-metal catalysts<sup>14</sup> were effective to the reaction, and Sc(OTf)<sub>3</sub> was found to provide the best results in 89% isolated yield (entries 8–10). Then, a variety of aprotic and protic solvents was evaluated, and the results showed that solvents were essential for the transformation. Aprotic solvents were beneficial to the reaction and afforded the product in good yields (entries 11–19). Finally, it was disadvantageous when the reaction was conducted at higher or lower temperatures (entries 20 and 21). After some attempts, the optimized conditions were considered as follows: **1a** (0.2 mmol) with **2a** (0.2 mmol) and Sc(OTf)<sub>3</sub> (5 mol %) in DMF (1 mL) at 100 °C for 4 h.

With the optimal reaction conditions established, the substrate scope was screened, and the results are summarized in Scheme 2. We first explored the scope of 2-aminopyridines by reacting with phenylpropionaldehyde **2a** and elemental sulfur under the standard conditions. In general, the three-component annulation reaction has good functional group tolerance. Both electron-rich and electron-deficient groups on the pyridine ring were well-tolerated and delivered the desired products in moderate to excellent yields (**3b–p**). The yields were not affected obviously with the methyl group on the different position of the 2-aminopyridine (**3b–d**). Interestingly, the reaction afforded excellent yield when a methoxy group substituted 2-aminopyridine was employed (**3f**). Halogen substituents such as fluoro, chloro, bromo, and iodo were smooth to generate the products in 48–75% yield (**3h–l**). Especially, the aryl bromide and iodide could be further functionalized under transition-metal-catalyzed conditions. In

Received: May 2, 2019

Scheme 1. Synthesis of Imidazo[1,2-*a*]pyridines from 2-Aminopyridines, Ynals, and Various Coupling ReagentsTable 1. Optimization of the Reaction Conditions<sup>a</sup>


entry	Lewis acid	solvent	temperature	yield (%)
1		DMF	100	35
2	LiCl	DMF	100	51
3	ZnCl <sub>2</sub>	DMF	100	74
4	FeCl <sub>3</sub>	DMF	100	82
5	CuI	DMF	100	62
6	AgNO <sub>3</sub>	DMF	100	76
7	BF <sub>3</sub> ·OEt <sub>2</sub>	DMF	100	80
8	Sc(OTf) <sub>3</sub>	DMF	100	89
9	Ce(OTf) <sub>3</sub>	DMF	100	83
10	Sm(OTf) <sub>3</sub>	DMF	100	72
11	Sc(OTf) <sub>3</sub>	DMSO	100	86
12	Sc(OTf) <sub>3</sub>	THF	100	82
13	Sc(OTf) <sub>3</sub>	MeCN	100	71
14	Sc(OTf) <sub>3</sub>	toluene	100	84
15	Sc(OTf) <sub>3</sub>	DCE	100	78
16	Sc(OTf) <sub>3</sub>	dioxane	100	82
17	Sc(OTf) <sub>3</sub>	H <sub>2</sub> O	100	np
18	Sc(OTf) <sub>3</sub>	HOAc	100	np
19	Sc(OTf) <sub>3</sub>	EtOH	100	np
20	Sc(OTf) <sub>3</sub>	DMF	120	87
21	Sc(OTf) <sub>3</sub>	DMF	80	81

<sup>a</sup>Reaction conditions: **1a** (0.2 mmol), **2a** (0.2 mmol), S<sub>8</sub> (12.8 mg, 0.05 mmol) with catalyst (5 mol %) in solvent (1.0 mL) under argon for 4 h. np = no product.

addition, moderate to good yields were obtained when the reactants containing the strong electron-withdrawing group were used (**3m–p**). We were quite pleased to observe that the trisubstituted 2-aminopyridine was also suitable and gave the product in 51% yield (**3q**). To our delight, the reaction could be carried out on large scale synthesis and delivered the product in good yield (**3a**).

In continuous evaluation of the scope and generality of the three-component reaction, a set of ynals proceeded smoothly

with 2-aminopyridine **1a** to yield the corresponding products under the given conditions (**4a–q**). As illustrated in Scheme 3, the annulation reaction was compatible with a handful of substituents such as alkyl, aryl, alkoxy, fluoro, chloro, trifluoromethyl, acetyl, and ester groups (**4a–p**). The ynal bearing a bulky *tert*-butyl group produced the product in good yield (**4e**). Satisfactory yields were obtained when two fluoro groups attached to the benzene ring, which achieved **4l** and **4m** in 63 and 72% yields, respectively. It is worth noting that the

Scheme 2. Substrate Scope of 2-Aminopyridines<sup>a</sup>

<sup>a</sup>Reaction conditions: **1** (0.2 mmol), **2a** (0.2 mmol),  $S_8$  (12.8 mg, 0.05 mmol) with  $Sc(OTf)_3$  (5 mol %) in DMF (1.0 mL) under argon at 100 °C for 4 h. <sup>b</sup>2 mmol scale of the reaction.


products bearing acetyl and ester groups could be further functionalized, which may be applicable in the pharmaceutical chemistry (**4o** and **4p**). In addition, the heteroaryl substituted ynal worked well to deliver the product in 75% yield (**4q**). To our delight, pyridine ring bearing electron-donating (methoxy group) and electron-withdrawing groups (iodo group) reacted successfully to render the corresponding products in good yields, irrespective of the ynals contained electron-rich or electron-deficient substituents on the benzene ring (**4r–u**). Those results clearly indicated that this three-component annulation reaction is general and effective for the imidazo[1,2-*a*]pyridine thione library. Unfortunately, we found the reaction did not form the corresponding product with the aliphatic ynal (**4v**).

Some control experiments were performed to investigate the reaction mechanism (Scheme 4). The reaction of imidazo[1,2-*a*]pyridin-3-yl(phenyl)methanone **5a** with elemental sulfur in the presence or absence of Lewis acid could not generate imidazo[1,2-*a*]pyridin-3-yl(phenyl)methanethione **3a**, which indicated that the direct thiation of the C=O bond was not involved in the reaction process. Taking the possible radical

pathway of elemental sulfur into account, different radical scavengers TEMPO, DPE, and BHT were introduced into the reaction system, and the generation of **3a** was not suppressed. These results suggested that the reaction did not proceed via a radical mechanism.


On the basis of the previous reports and above experimental results,<sup>10</sup> a plausible mechanism for this three-component annulation reaction has been proposed in Scheme 5. First, amination intermediate **A** was generated from the nucleophilic addition of **1a** and **2a** by the promotion of Lewis acid. Subsequently, successive nucleophilic attack of the pyridine nitrogen to the triple bond and the activation of sulfur powder was followed by a dehydration process that led to intermediate **B**. Finally, the elimination process of **B** achieved the imidazo[1,2-*a*]pyridine thione **3a**. It is likely that the Lewis acid accelerates the nucleophilic addition and nucleophilic attack of the pyridine nitrogen to the triple bond of the alkyne.

In summary, we described a novel and highly efficient annulation reaction of 2-aminopyridines, ynals, and elemental sulfur catalyzed by a Lewis acid. A wide range of imidazo[1,2-*a*]pyridine thiones bearing electron-rich or electron-deficient

Scheme 3. Substrate Scope of 2-Aminopyridines and Ynals<sup>a</sup>


<sup>a</sup>Reaction conditions: **1** (0.2 mmol), **2** (0.2 mmol),  $S_8$  (12.8 mg, 0.05 mmol) with  $Sc(OTf)_3$  (5 mol %) in DMF (1.0 mL) under argon at 100 °C for 4 h.

Scheme 4. Control Experiments



- (1) TEMPO (2 equiv), **3a**, 86%
- (2) DPE (2 equiv), **3a**, 82%
- (3) BHT (2 equiv), **3a**, 83%

Scheme 5. Possible Mechanism



substituents was obtained in moderate to excellent yields. This transformation features readily available raw materials, is additive-free, has good functional tolerance, simple operation, etc. In addition, it provides a new strategy for the use of elemental sulfur for sulfur-containing molecules. Further utilization of this approach in synthetic and pharmaceutical chemistry and mechanistic studies is currently underway in our laboratory.

## EXPERIMENTAL SECTION

**General Information.**  $^1\text{H}$  and  $^{13}\text{C}\{^1\text{H}\}$  NMR spectra were obtained on a 400 and 100 MHz NMR spectrometer. The chemical shifts are referenced to signals at 7.26 and 77.0 ppm, respectively, and chloroform was used as the solvent with TMS as the internal standard unless otherwise noted. Mass spectra were recorded on a GC-MS spectrometer at an ionization voltage of 70 eV equipped with a DB-WAX capillary column (internal diameter: 0.25 mm, length: 30 m). Elemental analyses were performed with a Vario EL elemental analyzer. High resolution mass spectra (HRMS) (TOF) were measured using an electrospray ionization (ESI) mass spectrometry. Silica gel (300–400 mesh) was used for flash column chromatography, eluting (unless otherwise stated) with ethyl acetate/petroleum ether (PE) (60–90 °C) mixture.

The 2-aminopyridines and propargyl aldehyde **2a** were commercially available from Sigma-Aldrich China. Others propargyl aldehydes were synthesized according to the literature<sup>15</sup> and data of known compounds were compared with the reported data.<sup>16</sup>

**General Procedure for the Preparation of Imidazo[1,2-*a*]pyridine Thiones.** A mixture of 2-aminopyridines (0.2 mmol), ynal (0.2 mmol), elemental sulfur (12.8 mg, 0.05 mmol), and  $\text{Sc}(\text{OTf})_3$  (0.01 mmol, 5 mol %) in DMF (1.0 mL) was stirred in a preheated oil bath at 100 °C for 4 h in a sealed tube under argon atmosphere. After the reaction was finished, water (5 mL) was added, and the solution was extracted with ethyl acetate (3 × 5 mL), and the combined extract was dried with anhydrous  $\text{MgSO}_4$ . Solvent was removed, and the residue was separated by column chromatography to give the pure sample.

**Large Scale Synthesis.** An oven-dried 25 mL screw cap test tube was charged with a magnetic stir bar, **1a** (188 mg, 2 mmol), **2a** (260 mg, 2 mmol), elemental sulfur (128 mg, 0.5 mmol),  $\text{Sc}(\text{OTf})_3$  (0.1 mmol, 5 mol %), and DMF (6.0 mL). The tube was then evacuated and backfilled with argon three times. Then, the tube was placed in a preheated oil bath at 100 °C for 4 h. After being cooled to room temperature, water (10 mL) was added; the solution was extracted with ethyl acetate (3 × 10 mL), and the combined extract was dried with anhydrous  $\text{MgSO}_4$ . Solvent was removed, and the residue was separated by column chromatography (ethyl acetate/petroleum ether = 1:3) to give **3a** (386 mg, 81%).

**Imidazo[1,2-*a*]pyridin-3-yl(phenyl)methanethione (**3a**).** Red solid (42 mg, 89%); mp 89–91 °C;  $R_f$  = 0.28 (ethyl acetate/

petroleum ether = 1:3);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  = 10.80 (dd,  $J$  = 6.9, 0.8 Hz, 1H), 8.07 (s, 1H), 7.82 (dd,  $J$  = 8.8, 0.8 Hz, 1H), 7.68–7.59 (m, 3H), 7.46 (dd,  $J$  = 10.6, 4.1 Hz, 1H), 7.39 (dd,  $J$  = 8.0, 7.2 Hz, 2H), 7.19 (t,  $J$  = 6.9 Hz, 1H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  210.7, 150.6, 148.3, 145.8, 134.6, 131.4, 130.3, 128.5, 127.9, 126.8, 117.8, 116.1. MS (EI)  $m/z$ : 238, 205, 161, 119, 78, 51. HRMS (ESI): calcd for  $\text{C}_{14}\text{H}_{11}\text{N}_2\text{S}$  [ $M + \text{H}$ ]<sup>+</sup> 239.0637; found 239.0634.

**(7-Methylimidazo[1,2-*a*]pyridin-3-yl)(phenyl)methanethione (**3b**).** Red solid (42 mg, 83%); mp 144–146 °C;  $R_f$  = 0.53 (ethyl acetate/petroleum ether = 1:2);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  10.70 (d,  $J$  = 7.0 Hz, 1H), 8.03 (s, 1H), 7.61 (d,  $J$  = 7.8 Hz, 3H), 7.47 (t,  $J$  = 7.3 Hz, 1H), 7.39 (t,  $J$  = 7.4 Hz, 2H), 7.03 (d,  $J$  = 7.0 Hz, 1H), 2.51 (s, 3H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  209.5, 151.3, 148.3, 146.3, 143.7, 134.7, 130.2, 128.5, 127.9, 126.3, 118.4, 116.8, 21.7. MS (EI)  $m/z$ : 252, 219, 175, 126, 92, 65, 39. HRMS (ESI): calcd for  $\text{C}_{15}\text{H}_{13}\text{N}_2\text{S}$  [ $M + \text{H}$ ]<sup>+</sup> 253.0794; found 253.0799.

**(6-Methylimidazo[1,2-*a*]pyridin-3-yl)(phenyl)methanethione (**3c**).** Red solid (43 mg, 86%); mp 139–141 °C;  $R_f$  = 0.51 (ethyl acetate/petroleum ether = 1:2);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  10.67 (s, 1H), 8.01 (s, 1H), 7.71 (d,  $J$  = 8.9 Hz, 1H), 7.62–7.57 (m, 2H), 7.52–7.43 (m, 2H), 7.38 (t,  $J$  = 7.4 Hz, 2H), 2.44 (s, 3H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  210.1, 149.6, 148.4, 145.8, 134.5, 134.2, 130.1, 128.5, 127.8, 126.5, 124.8, 117.0, 18.4. MS (EI)  $m/z$ : 252, 237, 219, 175, 126, 92, 65, 39. Anal. Calcd for  $\text{C}_{15}\text{H}_{12}\text{N}_2\text{S}$ : C, 71.40; H, 4.79; N, 11.10; Found: C, 71.06; H, 4.85; N, 11.16.

**(8-Methylimidazo[1,2-*a*]pyridin-3-yl)(phenyl)methanethione (**3d**).** Red solid (41 mg, 81%); mp 110–112 °C;  $R_f$  = 0.47 (ethyl acetate/petroleum ether = 1:2);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  10.67 (d,  $J$  = 6.9 Hz, 1H), 8.05 (s, 1H), 7.64–7.60 (m, 2H), 7.48 (dd,  $J$  = 10.6, 4.2 Hz, 2H), 7.40 (t,  $J$  = 7.5 Hz, 2H), 7.11 (t,  $J$  = 7.0 Hz, 1H), 2.70 (s, 3H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  210.7, 150.7, 148.5, 145.2, 135.2, 130.7, 130.3, 128.6, 127.9, 127.8, 124.7, 116.1, 16.8. MS (EI)  $m/z$ : 252, 219, 175, 126, 92, 65, 39. HRMS (ESI): calcd for  $\text{C}_{15}\text{H}_{13}\text{N}_2\text{S}$  [ $M + \text{H}$ ]<sup>+</sup> 253.0794; found 253.0794.

**Phenyl(6-phenylimidazo[1,2-*a*]pyridin-3-yl)methanethione (**3e**).** Red solid (39 mg, 62%); mp 125–127 °C;  $R_f$  = 0.51 (ethyl acetate/petroleum ether = 1:2);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  11.20 (s, 1H), 8.12 (s, 1H), 7.96 (dt,  $J$  = 17.8, 5.3 Hz, 2H), 7.67 (t,  $J$  = 6.7 Hz, 4H), 7.54 (t,  $J$  = 7.7 Hz, 3H), 7.45 (q,  $J$  = 7.6 Hz, 3H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  211.1, 149.9, 148.6, 146.2, 136.5, 135.0, 131.5, 131.0, 130.4, 129.3, 128.6, 128.5, 128.1, 127.2, 124.5, 117.7. MS (EI)  $m/z$ : 314, 281, 237, 157, 127, 102, 77, 51. HRMS (ESI): calcd for  $\text{C}_{20}\text{H}_{15}\text{N}_2\text{S}$  [ $M + \text{H}$ ]<sup>+</sup> 315.0950; found 315.0945.

**(7-Methoxyimidazo[1,2-*a*]pyridin-3-yl)(phenyl)methanethione (**3f**).** Red solid (49 mg, 92%); mp 150–152 °C;  $R_f$  = 0.23 (ethyl acetate/petroleum ether = 1:2);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  10.70 (d,  $J$  = 7.6 Hz, 1H), 7.99 (s, 1H), 7.65–7.60 (m, 2H), 7.50–7.45 (m, 1H), 7.40 (t,  $J$  = 7.4 Hz, 2H), 7.13 (d,  $J$  = 2.6 Hz, 1H), 6.87 (dd,  $J$  = 7.6, 2.6 Hz, 1H), 3.96 (s, 3H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  208.2, 162.4, 153.7, 148.3, 147.0, 134.8, 130.1, 128.6,



128.3, 127.9, 109.2, 96.7, 56.0. MS (EI)  $m/z$ : 268, 235, 224, 191, 176, 134, 121, 77, 51. HRMS (ESI): calcd for  $C_{15}H_{13}N_2OS$   $[M + H]^+$  269.0743; found 269.0749.

(8-(Benzyloxy)imidazo[1,2-*a*]pyridin-3-yl)(phenyl)methanethione (**3g**). Red solid (56 mg, 81%); mp 115–117 °C;  $R_f$  = 0.60 (ethyl acetate/petroleum ether = 1:2);  $^1H$  NMR (400 MHz,  $CDCl_3$ ):  $\delta$  10.36 (dd,  $J$  = 6.1, 1.6 Hz, 1H), 8.03 (s, 1H), 7.66–7.62 (m, 2H), 7.49 (d,  $J$  = 7.3 Hz, 3H), 7.43–7.31 (m, 5H), 7.06–6.99 (m, 2H), 5.39 (s, 2H).  $^{13}C\{^1H\}$  NMR (100 MHz,  $CDCl_3$ ):  $\delta$  211.8, 148.6, 147.7, 144.6, 144.3, 135.6, 135.4, 130.4, 128.6, 128.6, 128.2, 128.0, 127.3, 119.9, 116.0, 110.9, 71.2. MS (EI)  $m/z$ : 344, 267, 253, 172, 121, 91, 65, 39. HRMS (ESI): calcd for  $C_{21}H_{17}N_2OS$   $[M + H]^+$  345.1056; found 345.1058.

(6-Fluoroimidazo[1,2-*a*]pyridin-3-yl)(phenyl)methanethione (**3h**). Red solid (38 mg, 75%); mp 121–123 °C;  $R_f$  = 0.39 (ethyl acetate/petroleum ether = 1:2);  $^1H$  NMR (400 MHz,  $CDCl_3$ ):  $\delta$  10.94 (dd,  $J$  = 5.2, 2.1 Hz, 1H), 8.11 (s, 1H), 7.83 (dd,  $J$  = 9.6, 5.1 Hz, 1H), 7.64–7.57 (m, 3H), 7.54–7.48 (m, 1H), 7.41 (dd,  $J$  = 10.3, 4.7 Hz, 2H).  $^{13}C\{^1H\}$  NMR (100 MHz,  $CDCl_3$ ):  $\delta$  211.8, 155.2 (d,  $J$  = 239 Hz), 148.2, 148.1, 146.0 (d,  $J$  = 3 Hz), 135.0, 130.6, 128.6, 128.1, 122.3 (d,  $J$  = 25 Hz), 118.1 (d,  $J$  = 9 Hz), 114.2 (d,  $J$  = 22.5 Hz). MS (EI)  $m/z$ : 256, 223, 179, 128, 96, 51. HRMS (ESI): calcd for  $C_{14}H_{10}FN_2S$   $[M + H]^+$  257.0543; found 257.0540.

(7-Chloroimidazo[1,2-*a*]pyridin-3-yl)(phenyl)methanethione (**3i**). Red solid (40 mg, 73%); mp 137–139 °C;  $R_f$  = 0.64 (ethyl acetate/petroleum ether = 1:4);  $^1H$  NMR (400 MHz,  $CDCl_3$ ):  $\delta$  10.74 (d,  $J$  = 7.3 Hz, 1H), 8.07 (s, 1H), 7.83 (s, 1H), 7.62 (d,  $J$  = 7.3 Hz, 2H), 7.53–7.49 (m, 1H), 7.42 (t,  $J$  = 7.4 Hz, 2H), 7.18 (d,  $J$  = 6.1 Hz, 1H).  $^{13}C\{^1H\}$  NMR (100 MHz,  $CDCl_3$ ):  $\delta$  211.5, 150.9, 148.2, 146.1, 137.5, 134.5, 130.7, 128.6, 128.1, 127.2, 117.3, 117.1. MS (EI)  $m/z$ : 272, 239, 195, 112, 89, 76, 51. HRMS (ESI): calcd for  $C_{14}H_{10}ClN_2S$   $[M + H]^+$  273.0248; found 273.0245.

(6-Chloroimidazo[1,2-*a*]pyridin-3-yl)(phenyl)methanethione (**3j**). Red solid (31 mg, 57%); mp 175–177 °C;  $R_f$  = 0.39 (ethyl acetate/petroleum ether = 1:3);  $^1H$  NMR (400 MHz,  $CDCl_3$ ):  $\delta$  10.95 (d,  $J$  = 1.5 Hz, 1H), 8.09 (s, 1H), 7.81 (d,  $J$  = 9.4 Hz, 1H), 7.65 (dd,  $J$  = 11.7, 7.3 Hz, 3H), 7.52 (t,  $J$  = 7.4 Hz, 1H), 7.43 (t,  $J$  = 7.5 Hz, 2H).  $^{13}C\{^1H\}$  NMR (100 MHz,  $CDCl_3$ ):  $\delta$  212.1, 148.9, 148.3, 145.6, 134.4, 132.3, 130.8, 128.6, 128.1, 124.8, 124.6, 118.1. MS (EI)  $m/z$ : 272, 239, 195, 112, 89, 76, 51. HRMS (ESI): calcd for  $C_{14}H_{10}ClN_2S$   $[M + H]^+$  273.0248; found 273.0240.

(6-Bromoimidazo[1,2-*a*]pyridin-3-yl)(phenyl)methanethione (**3k**). Red solid (44 mg, 70%); mp 194–196 °C;  $R_f$  = 0.35 (ethyl acetate/petroleum ether = 1:3);  $^1H$  NMR (400 MHz,  $CDCl_3$ ):  $\delta$  11.03 (s, 1H), 8.07 (s, 1H), 7.75 (s, 2H), 7.63 (d,  $J$  = 7.2 Hz, 2H), 7.52 (t,  $J$  = 7.4 Hz, 1H), 7.43 (t,  $J$  = 7.5 Hz, 2H).  $^{13}C\{^1H\}$  NMR (100 MHz,  $CDCl_3$ ):  $\delta$  212.1, 149.1, 148.3, 145.5, 134.5, 134.3, 130.8, 128.6, 128.1, 126.9, 118.4, 111.1. MS (EI)  $m/z$ : 317, 283, 239, 158, 89, 77, 51. HRMS (ESI): calcd for  $C_{14}H_9BrN_2NaS$   $[M + Na]^+$  338.9562; found 338.9568.

(6-Iodoimidazo[1,2-*a*]pyridin-3-yl)(phenyl)methanethione (**3l**). Red solid (35 mg, 48%); mp 198–200 °C;  $R_f$  = 0.38 (ethyl acetate/petroleum ether = 1:3);  $^1H$  NMR (400 MHz,  $CDCl_3$ ):  $\delta$  11.09 (s, 1H), 8.01 (s, 1H), 7.87 (dd,  $J$  = 9.2, 1.1 Hz, 1H), 7.65–7.61 (m, 3H), 7.52 (t,  $J$  = 7.3 Hz, 1H), 7.42 (t,  $J$  = 7.5 Hz, 2H).  $^{13}C\{^1H\}$  NMR (100 MHz,  $CDCl_3$ ):  $\delta$  212.0, 149.3, 148.3, 145.1, 139.2, 133.9, 131.6, 130.7, 128.6, 128.1, 118.8, 79.6. MS (EI)  $m/z$ : 364, 287, 236, 204, 160, 118, 89, 77, 51. HRMS (ESI): calcd for  $C_{14}H_{10}IN_2S$   $[M + H]^+$  364.9604; found 364.9607.

Phenyl(7-(trifluoromethyl)imidazo[1,2-*a*]pyridin-3-yl)methanethione (**3m**). Red solid (44 mg, 72%); mp 99–101 °C;  $R_f$  = 0.56 (ethyl acetate/petroleum ether = 1:2);  $^1H$  NMR (400 MHz,  $CDCl_3$ ):  $\delta$  11.19 (s, 1H), 8.18 (s, 1H), 7.96 (d,  $J$  = 9.2 Hz, 1H), 7.85–7.80 (m, 1H), 7.65 (d,  $J$  = 7.2 Hz, 2H), 7.55 (t,  $J$  = 7.3 Hz, 1H), 7.44 (t,  $J$  = 7.6 Hz, 2H).  $^{13}C\{^1H\}$  NMR (100 MHz,  $CDCl_3$ ):  $\delta$  213.1, 150.3, 148.2, 146.1, 134.6, 131.1, 128.6, 128.2, 126.9 (q,  $J$  = 3 Hz), 125.7 (q,  $J$  = 5 Hz), 123.1 (q,  $J$  = 271 Hz), 120.2 (q,  $J$  = 34 Hz), 118.7. MS (EI)  $m/z$ : 306, 273, 229, 146, 121, 89, 69, 51. HRMS (ESI): calcd for  $C_{15}H_{10}F_3N_2S$   $[M + H]^+$  307.0511; found 307.0519.

(7-Nitroimidazo[1,2-*a*]pyridin-3-yl)(phenyl)methanethione (**3n**). Green solid (23 mg, 40%); mp 157–159 °C;  $R_f$  = 0.44 (ethyl acetate/petroleum ether = 1:2);  $^1H$  NMR (400 MHz,  $CDCl_3$ ): 11.77–11.73 (m, 1H), 8.41 (dd,  $J$  = 9.7, 2.2 Hz, 1H), 8.22 (s, 1H), 7.93 (d,  $J$  = 9.7 Hz, 1H), 7.67 (dd,  $J$  = 5.1, 3.3 Hz, 2H), 7.57 (dd,  $J$  = 10.6, 4.3 Hz, 1H), 7.45 (dd,  $J$  = 10.5, 4.7 Hz, 2H).  $^{13}C\{^1H\}$  NMR (100 MHz,  $CDCl_3$ ):  $\delta$  213.9, 150.3, 148.0, 146.6, 139.6, 134.9, 131.4, 128.7, 128.3, 126.6, 124.5, 117.7. MS (EI)  $m/z$ : 283, 266, 252, 236, 206, 192, 160, 134, 121, 89, 77, 51. Anal. Calcd for  $C_{14}H_9N_3O_2S$ : C, 59.35; H, 3.20; N, 14.83; Found: C, 59.02; H, 3.29; N, 14.76.

3-(Phenylcarbonothioyl)imidazo[1,2-*a*]pyridine-7-carbonitrile (**3o**). Green solid (24 mg, 45%); mp 162–164 °C;  $R_f$  = 0.44 (ethyl acetate/petroleum ether = 1:4);  $^1H$  NMR (400 MHz,  $CDCl_3$ ):  $\delta$  10.75 (d,  $J$  = 7.2 Hz, 1H), 8.21 (s, 2H), 7.67–7.64 (m, 2H), 7.57 (s, 1H), 7.45 (t,  $J$  = 7.7 Hz, 2H), 7.32 (dd,  $J$  = 7.2, 1.6 Hz, 1H).  $^{13}C\{^1H\}$  NMR (100 MHz,  $CDCl_3$ ):  $\delta$  214.2, 148.4, 148.2, 145.9, 134.6, 131.4, 128.7, 128.3, 127.2, 123.4, 116.6, 116.2, 113.1. MS (EI)  $m/z$ : 263, 230, 186, 159, 103, 77, 43. HRMS (ESI): calcd for  $C_{15}H_{10}N_3S$   $[M + H]^+$  264.0590; found 264.0596.

Methyl 3-(Phenylcarbonothioyl)imidazo[1,2-*a*]pyridine-7-carboxylate (**3p**). Green solid (40 mg, 68%); mp 166–168 °C;  $R_f$  = 0.29 (ethyl acetate/petroleum ether = 1:2);  $^1H$  NMR (400 MHz,  $CDCl_3$ ):  $\delta$  10.74 (d,  $J$  = 7.2 Hz, 1H), 8.51 (s, 1H), 8.17 (s, 1H), 7.77 (dd,  $J$  = 7.2, 1.5 Hz, 1H), 7.65 (d,  $J$  = 7.3 Hz, 2H), 7.52 (d,  $J$  = 7.5 Hz, 1H), 7.43 (t,  $J$  = 7.6 Hz, 2H), 4.02 (s, 3H).  $^{13}C\{^1H\}$  NMR (100 MHz,  $CDCl_3$ ):  $\delta$  213.0, 164.5, 149.8, 148.4, 146.2, 134.7, 131.8, 130.9, 128.7, 128.2, 126.4, 119.8, 115.3, 52.9. MS (EI)  $m/z$ : 296, 281, 263, 219, 121, 102, 77, 41. HRMS (ESI): calcd for  $C_{16}H_{13}N_3O_2S$   $[M + H]^+$  297.0692; found 297.0697.

(6,8-Dibromo-7-methylimidazo[1,2-*a*]pyridin-3-yl)(phenyl)methanethione (**3q**). Red solid (42 mg, 51%); mp 171–173 °C;  $R_f$  = 0.41 (ethyl acetate/petroleum ether = 1:5);  $^1H$  NMR (400 MHz,  $CDCl_3$ ):  $\delta$  10.96 (s, 1H), 8.05 (s, 1H), 7.63 (d,  $J$  = 7.3 Hz, 2H), 7.53 (t,  $J$  = 7.4 Hz, 1H), 7.42 (t,  $J$  = 7.6 Hz, 2H), 2.75 (s, 3H).  $^{13}C\{^1H\}$  NMR (100 MHz,  $CDCl_3$ ):  $\delta$  212.4, 148.2, 147.9, 144.9, 141.8, 135.1, 131.0, 128.7, 128.2, 125.6, 113.8, 113.1, 23.6. MS (EI)  $m/z$ : 409, 377, 333, 249, 165, 145, 125, 121, 102, 90, 77, 51. Anal. Calcd for  $C_{15}H_{10}Br_2N_2S$ : C, 43.93; H, 2.46; N, 6.83; Found: C, 43.70; H, 2.53; N, 6.77.

Imidazo[1,2-*a*]pyridin-3-yl(*p*-tolyl)methanethione (**4a**). Red solid (44 mg, 87%); mp 153–155 °C;  $R_f$  = 0.42 (ethyl acetate/petroleum ether = 1:2);  $^1H$  NMR (400 MHz,  $CDCl_3$ ):  $\delta$  10.80 (d,  $J$  = 6.9 Hz, 1H), 8.11 (s, 1H), 7.85 (d,  $J$  = 8.8 Hz, 1H), 7.67 (t,  $J$  = 7.9 Hz, 1H), 7.57 (d,  $J$  = 8.0 Hz, 2H), 7.21 (t,  $J$  = 7.0 Hz, 3H), 2.41 (s, 3H).  $^{13}C\{^1H\}$  NMR (100 MHz,  $CDCl_3$ ):  $\delta$  211.0, 150.7, 146.0, 145.6, 141.2, 134.6, 131.2, 128.9, 128.7, 127.0, 117.9, 116.0, 21.3. MS (EI)  $m/z$ : 252, 219, 161, 78, 51. HRMS (ESI): calcd for  $C_{15}H_{13}N_2S$   $[M + H]^+$  253.0794; found 253.0792.

Imidazo[1,2-*a*]pyridin-3-yl(*m*-tolyl)methanethione (**4b**). Red solid (42 mg, 84%); mp 110–112 °C;  $R_f$  = 0.52 (ethyl acetate/petroleum ether = 1:2);  $^1H$  NMR (400 MHz,  $CDCl_3$ ):  $\delta$  10.82 (d,  $J$  = 6.9 Hz, 1H), 8.09 (s, 1H), 7.84 (d,  $J$  = 8.8 Hz, 1H), 7.70–7.65 (m, 1H), 7.45 (s, 1H), 7.40 (dd,  $J$  = 6.3, 1.8 Hz, 1H), 7.29 (d,  $J$  = 6.3 Hz, 2H), 7.21 (td,  $J$  = 6.9, 1.1 Hz, 1H), 2.39 (s, 3H).  $^{13}C\{^1H\}$  NMR (100 MHz,  $CDCl_3$ ):  $\delta$  211.4, 150.8, 148.5, 145.9, 137.8, 134.7, 131.4, 131.2, 129.3, 127.8, 127.0, 125.8, 117.9, 116.2, 21.2. MS (EI)  $m/z$ : 252, 237, 219, 161, 126, 91, 78, 51. HRMS (ESI): calcd for  $C_{15}H_{13}N_2S$   $[M + H]^+$  253.0794; found 253.0799.

Imidazo[1,2-*a*]pyridin-3-yl(*o*-tolyl)methanethione (**4c**). Red solid (41 mg, 81%); mp 129–131 °C;  $R_f$  = 0.52 (ethyl acetate/petroleum ether = 1:2);  $^1H$  NMR (400 MHz,  $CDCl_3$ ):  $\delta$  11.00 (d,  $J$  = 6.9 Hz, 1H), 7.88–7.83 (m, 2H), 7.75–7.70 (m, 1H), 7.30 (ddd,  $J$  = 7.8, 4.8, 2.1 Hz, 2H), 7.23 (dd,  $J$  = 4.0, 3.3 Hz, 3H), 2.23 (s, 3H).  $^{13}C\{^1H\}$  NMR (100 MHz,  $CDCl_3$ ):  $\delta$  212.8, 150.8, 147.8, 146.1, 134.8, 133.1, 131.8, 130.4, 128.4, 127.1, 127.1, 125.4, 118.0, 116.7, 19.3. MS (EI)  $m/z$ : 252, 219, 161, 134, 115, 79, 51. Anal. Calcd for  $C_{15}H_{12}N_2S$ : C, 71.40; H, 4.79; N, 11.10; Found: C, 71.12; H, 4.86; N, 11.05.

(3,5-Dimethylphenyl)imidazo[1,2-*a*]pyridin-3-yl)methanethione (**4d**). Red solid (44 mg, 83%); mp 129–131 °C;  $R_f$  = 0.50 (ethyl acetate/petroleum ether = 1:2);  $^1H$  NMR (400 MHz,  $CDCl_3$ ):  $\delta$

10.81 (d,  $J = 6.9$  Hz, 1H), 8.10 (s, 1H), 7.83 (d,  $J = 8.8$  Hz, 1H), 7.69–7.63 (m, 1H), 7.24 (s, 2H), 7.20 (t,  $J = 6.9$  Hz, 1H), 7.13 (s, 1H), 2.35 (s, 6H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  211.6, 150.7, 148.6, 145.8, 137.5, 134.6, 132.1, 131.3, 126.9, 126.5, 117.8, 116.0, 21.1. MS (EI)  $m/z$ : 266, 233, 161, 78, 51. HRMS (ESI): calcd for  $\text{C}_{16}\text{H}_{15}\text{N}_2\text{S}$  [ $\text{M} + \text{H}$ ] $^+$  267.0950; found 267.0946.

**(4-(tert-Butyl)phenyl)(imidazo[1,2-a]pyridin-3-yl)methanethione (4e).** Red solid (46 mg, 78%); mp 120–122 °C;  $R_f = 0.52$  (ethyl acetate/petroleum ether = 1:2);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  10.82 (d,  $J = 6.9$  Hz, 1H), 8.14 (s, 1H), 7.85 (d,  $J = 8.8$  Hz, 1H), 7.70–7.65 (m, 1H), 7.61 (d,  $J = 8.4$  Hz, 2H), 7.43 (d,  $J = 8.4$  Hz, 2H), 7.21 (td,  $J = 6.9$ , 1.0 Hz, 1H), 1.36 (s, 9H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  211.0, 154.2, 150.7, 145.9, 145.7, 134.6, 131.2, 128.7, 127.0, 125.0, 117.9, 116.0, 34.8, 31.1. MS (EI)  $m/z$ : 294, 279, 263, 237, 161, 125, 78, 51. HRMS (ESI): calcd for  $\text{C}_{18}\text{H}_{19}\text{N}_2\text{S}$  [ $\text{M} + \text{H}$ ] $^+$  295.1263; found 295.1257.

**[1,1'-Biphenyl]-4-yl(imidazo[1,2-a]pyridin-3-yl)methanethione (4f).** Red solid (47 mg, 75%); mp 172–174 °C;  $R_f = 0.45$  (ethyl acetate/petroleum ether = 1:2);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  10.84 (d,  $J = 6.9$  Hz, 1H), 8.19 (s, 1H), 7.88 (d,  $J = 8.8$  Hz, 1H), 7.75 (d,  $J = 8.2$  Hz, 2H), 7.71 (d,  $J = 7.9$  Hz, 1H), 7.65 (d,  $J = 6.9$  Hz, 4H), 7.48 (t,  $J = 7.5$  Hz, 2H), 7.40 (t,  $J = 7.3$  Hz, 1H), 7.27–7.21 (m, 1H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  210.2, 150.8, 147.2, 145.7, 143.4, 139.9, 134.7, 131.4, 129.3, 128.8, 127.8, 127.0, 127.0, 126.7, 118.0, 116.2. MS (EI)  $m/z$ : 314, 281, 161, 78, 44. Anal. Calcd for  $\text{C}_{20}\text{H}_{14}\text{N}_2\text{S}$ : C, 76.40; H, 4.49; N, 8.91; Found: C, 76.03; H, 4.58; N, 8.82.

**Imidazo[1,2-a]pyridin-3-yl(naphthalen-1-yl)methanethione (4g).** Red solid (44 mg, 76%); mp 163–165 °C;  $R_f = 0.33$  (ethyl acetate/petroleum ether = 1:3);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  11.10 (d,  $J = 6.3$  Hz, 1H), 7.88 (dd,  $J = 18.6$ , 9.7 Hz, 4H), 7.78 (s, 1H), 7.72 (t,  $J = 7.8$  Hz, 1H), 7.50 (td,  $J = 15.1$ , 7.4 Hz, 3H), 7.38 (t,  $J = 7.5$  Hz, 1H), 7.29 (t,  $J = 6.9$  Hz, 1H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  210.8, 150.8, 146.6, 145.5, 135.5, 133.4, 131.9, 130.2, 128.8, 128.0, 127.1, 126.5, 126.0, 125.3, 124.8, 124.6, 118.0, 116.8. MS (EI)  $m/z$ : 288, 255, 209, 161, 144, 127, 78, 51. HRMS (ESI): calcd for  $\text{C}_{18}\text{H}_{13}\text{N}_2\text{S}$  [ $\text{M} + \text{H}$ ] $^+$  289.0794; found 289.0797.

**Imidazo[1,2-a]pyridin-3-yl(4-methoxyphenyl)methanethione (4h).** Red solid (29 mg, 55%); mp 160–162 °C;  $R_f = 0.35$  (ethyl acetate/petroleum ether = 1:2);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  10.71 (d,  $J = 6.9$  Hz, 1H), 8.10 (s, 1H), 7.86 (d,  $J = 8.8$  Hz, 1H), 7.68 (dd,  $J = 18.3$ , 8.4 Hz, 3H), 7.21 (t,  $J = 6.9$  Hz, 1H), 6.94 (d,  $J = 8.7$  Hz, 2H), 3.89 (s, 3H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  209.8, 162.3, 150.7, 145.1, 141.5, 134.5, 131.1, 131.0, 127.0, 117.9, 115.8, 113.4, 55.5. MS (EI)  $m/z$ : 268, 252, 235, 224, 192, 161, 134, 112, 89, 78, 51. Anal. Calcd for  $\text{C}_{15}\text{H}_{12}\text{N}_2\text{O}_2\text{S}$ : C, 67.14; H, 4.51; N, 10.44; Found: C, 66.80; H, 4.62; N, 10.51.

**Benzo[d][1,3]dioxol-5-yl(imidazo[1,2-a]pyridin-3-yl)methanethione (4i).** Red solid (37 mg, 65%); mp 195–197 °C;  $R_f = 0.33$  (ethyl acetate/petroleum ether = 1:2);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  10.63 (dt,  $J = 7.0$ , 1.1 Hz, 1H), 8.06 (s, 1H), 7.83–7.73 (m, 1H), 7.64–7.57 (m, 1H), 7.23 (d,  $J = 1.8$  Hz, 1H), 7.17–7.11 (m, 2H), 6.76 (d,  $J = 8.1$  Hz, 1H), 5.99 (s, 2H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  209.1, 150.8, 150.4, 147.8, 145.2, 143.0, 134.6, 131.1, 127.0, 124.1, 118.0, 115.9, 109.9, 107.4, 101.8. MS (EI)  $m/z$ : 282, 249, 223, 179, 161, 141, 112, 90, 78, 51. HRMS (ESI): calcd for  $\text{C}_{15}\text{H}_{11}\text{N}_2\text{O}_2\text{S}$  [ $\text{M} + \text{H}$ ] $^+$  283.0536; found 283.0536.

**(4-Fluorophenyl)(imidazo[1,2-a]pyridin-3-yl)methanethione (4j).** Red solid (41 mg, 81%); mp 165–167 °C;  $R_f = 0.22$  (ethyl acetate/petroleum ether = 1:3);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  10.77 (d,  $J = 6.9$  Hz, 1H), 8.07 (s, 1H), 7.85 (d,  $J = 8.8$  Hz, 1H), 7.72–7.67 (m, 1H), 7.67–7.62 (m, 2H), 7.23 (td,  $J = 6.9$ , 1.0 Hz, 1H), 7.13–7.06 (m, 2H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  208.9, 164.2 (d,  $J = 251$  Hz), 150.9, 145.6, 144.6 (d,  $J = 3$  Hz), 134.68 131.6, 130.7 (d,  $J = 9$  Hz), 126.9, 118.0, 116.3, 115.1 (d,  $J = 22$  Hz). MS (EI)  $m/z$ : 256, 223, 161, 128, 95, 78, 51. Anal. Calcd for  $\text{C}_{14}\text{H}_9\text{FN}_2\text{S}$ : C, 65.61; H, 3.54; N, 10.93; Found: C, 65.40; H, 3.59; N, 10.88.

**(4-Chlorophenyl)(imidazo[1,2-a]pyridin-3-yl)methanethione (4k).** Red solid (46 mg, 85%); mp 179–181 °C;  $R_f = 0.50$  (ethyl acetate/petroleum ether = 1:2);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$

10.79 (d,  $J = 6.9$  Hz, 1H), 8.08 (s, 1H), 7.87 (d,  $J = 8.8$  Hz, 1H), 7.74–7.68 (m, 1H), 7.60–7.55 (m, 2H), 7.39 (d,  $J = 8.4$  Hz, 2H), 7.27–7.22 (m, 1H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  208.8, 150.9, 146.7, 145.7, 136.7, 134.6, 131.7, 129.9, 128.3, 126.9, 118.0, 116.4. MS (EI)  $m/z$ : 272, 239, 161, 136, 118, 78, 51. Anal. Calcd for  $\text{C}_{14}\text{H}_9\text{ClN}_2\text{S}$ : C, 61.65; H, 3.33; N, 10.27; Found: C, 61.32; H, 3.42; N, 10.35.

**(3,5-Difluorophenyl)(imidazo[1,2-a]pyridin-3-yl)methanethione (4l).** Red solid (35 mg, 63%); mp 185–187 °C;  $R_f = 0.30$  (ethyl acetate/petroleum ether = 1:3);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  10.81 (dt,  $J = 7.0$ , 1.0 Hz, 1H), 8.12 (s, 1H), 7.92–7.87 (m, 1H), 7.79–7.74 (m, 1H), 7.29 (td,  $J = 7.0$ , 1.2 Hz, 1H), 7.19–7.13 (m, 2H), 6.94 (tt,  $J = 8.7$ , 2.3 Hz, 1H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  206.2, 162.2 (dd,  $J = 249$ , 12 Hz), 151.2, 150.6 (t,  $J = 9$  Hz), 146.0, 134.6, 132.2, 127.0, 118.2, 116.8, 111.7 (d,  $J = 26$  Hz), 105.3 (t,  $J = 25$  Hz). MS (EI)  $m/z$ : 274, 241, 161, 137, 113, 78, 51. HRMS (ESI): calcd for  $\text{C}_{14}\text{H}_9\text{F}_2\text{N}_2\text{S}$  [ $\text{M} + \text{H}$ ] $^+$  275.0449; found 275.0443.

**(3,4-Difluorophenyl)(imidazo[1,2-a]pyridin-3-yl)methanethione (4m).** Red solid (39 mg, 72%); mp 170–172 °C;  $R_f = 0.37$  (ethyl acetate/petroleum ether = 1:2);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  10.77 (d,  $J = 6.9$  Hz, 1H), 8.10 (s, 1H), 7.89 (d,  $J = 8.8$  Hz, 1H), 7.76–7.71 (m, 1H), 7.57–7.50 (m, 1H), 7.39 (ddd,  $J = 8.1$ , 3.9, 1.8 Hz, 1H), 7.30–7.26 (m, 1H), 7.25–7.18 (m, 1H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  206.9, 151.9 (dd,  $J = 252$ , 12 Hz), 151.1, 149.8 (dd,  $J = 249$ , 13 Hz), 145.7, 145.0 (dd,  $J = 5$ , 4 Hz), 134.7, 131.9, 127.0, 124.9 (dd,  $J = 6$ , 3 Hz), 118.2, 118.2 (d,  $J = 19$  Hz), 116.8 (d,  $J = 18$  Hz), 116.6. MS (EI)  $m/z$ : 274, 241, 161, 137, 78, 51. HRMS (ESI): calcd for  $\text{C}_{14}\text{H}_9\text{F}_2\text{N}_2\text{S}$  [ $\text{M} + \text{H}$ ] $^+$  275.0449; found 275.0460.

**Imidazo[1,2-a]pyridin-3-yl(3-(trifluoromethyl)phenyl)methanethione (4n).** Red solid (51 mg, 84%); mp 126–128 °C;  $R_f = 0.47$  (ethyl acetate/petroleum ether = 1:2);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  10.85–10.79 (m, 1H), 8.05 (s, 1H), 7.88 (dd,  $J = 5.1$ , 3.5 Hz, 2H), 7.80–7.71 (m, 3H), 7.55 (t,  $J = 7.8$  Hz, 1H), 7.30–7.25 (m, 1H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  208.0, 151.0, 148.7, 145.9, 134.8, 132.1, 131.5, 130.5 (q,  $J = 33$  Hz), 128.6, 126.9, 126.7 (q,  $J = 4$  Hz), 125.2 (q,  $J = 4$  Hz), 123.6 (q,  $J = 271$  Hz), 118.1, 116.7. MS (EI)  $m/z$ : 306, 273, 161, 78, 51. Anal. Calcd for  $\text{C}_{15}\text{H}_9\text{F}_3\text{N}_2\text{S}$ : C, 58.82; H, 2.96; N, 9.15; Found: C, 58.54; H, 3.05; N, 9.22.

**1-(4-(Imidazo[1,2-a]pyridine-3-carbonothioyl)phenyl)ethanone (4o).** Green solid (45 mg, 81%); mp 159–161 °C;  $R_f = 0.28$  (ethyl acetate/petroleum ether = 1:1);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  10.83 (d,  $J = 6.6$  Hz, 1H), 8.04 (s, 1H), 7.98 (d,  $J = 7.9$  Hz, 2H), 7.86 (d,  $J = 8.7$  Hz, 1H), 7.72 (t,  $J = 7.8$  Hz, 1H), 7.66 (d,  $J = 7.9$  Hz, 2H), 7.26 (t,  $J = 6.6$  Hz, 1H), 2.63 (s, 3H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  208.7, 197.3, 151.9, 150.9, 146.0, 137.7, 134.7, 132.0, 128.5, 128.0, 126.9, 118.0, 116.7, 26.7. MS (EI)  $m/z$ : 280, 236, 161, 132, 118, 89, 78, 51. HRMS (ESI): calcd for  $\text{C}_{16}\text{H}_{12}\text{N}_2\text{NaOS}$  [ $\text{M} + \text{Na}$ ] $^+$  303.0563; found 303.0559.

**Methyl 4-(Imidazo[1,2-a]pyridine-3-carbonothioyl)benzoate (4p).** Red solid (41 mg, 70%); mp 180–182 °C;  $R_f = 0.37$  (ethyl acetate/petroleum ether = 1:2);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  10.86 (d,  $J = 6.9$  Hz, 1H), 8.11–8.05 (m, 3H), 7.89 (d,  $J = 8.8$  Hz, 1H), 7.77–7.72 (m, 1H), 7.66 (d,  $J = 8.4$  Hz, 2H), 7.28 (td,  $J = 7.1$ , 1.2 Hz, 1H), 3.96 (s, 3H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  209.0, 166.3, 151.9, 146.0, 134.7, 132.0, 131.2, 129.3, 128.6, 128.3, 127.0, 118.1, 116.6, 52.3. MS (EI)  $m/z$ : 296, 280, 263, 236, 161, 132, 118, 89, 78, 51. Anal. Calcd for  $\text{C}_{16}\text{H}_{12}\text{N}_2\text{O}_2\text{S}$ : C, 64.85; H, 4.08; N, 9.45; Found: C, 64.59; H, 4.17; N, 9.51.

**Imidazo[1,2-a]pyridin-3-yl(thiophen-2-yl)methanethione (4q).** Red solid (37 mg, 75%); mp 180–182 °C;  $R_f = 0.31$  (ethyl acetate/petroleum ether = 1:2);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  10.30 (dt,  $J = 7.0$ , 1.1 Hz, 1H), 8.25 (s, 1H), 7.76 (dt,  $J = 8.9$ , 1.1 Hz, 1H), 7.64 (dd,  $J = 5.1$ , 1.2 Hz, 1H), 7.57–7.52 (m, 1H), 7.44 (dd,  $J = 3.8$ , 1.2 Hz, 1H), 7.09 (ddd,  $J = 11.3$ , 5.0, 1.2 Hz, 2H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  197.8, 154.5, 150.7, 143.6, 134.8, 133.2, 130.7, 129.3, 128.1, 126.9, 117.9, 115.4. MS (EI)  $m/z$ : 244, 211, 161, 122, 100, 78, 51. HRMS (ESI): calcd for  $\text{C}_{12}\text{H}_9\text{N}_2\text{S}_2$  [ $\text{M} + \text{H}$ ] $^+$  245.0202; found 245.0201.



(7-Methoxyimidazo[1,2-a]pyridin-3-yl)(o-tolyl)methanethione (**4r**). Red solid (50 mg, 88%); mp 120–122 °C;  $R_f$  = 0.24 (ethyl acetate/petroleum ether = 1:3);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  10.80 (d,  $J$  = 7.5 Hz, 1H), 7.71 (s, 1H), 7.28–7.24 (m, 1H), 7.22–7.18 (m, 3H), 7.12 (d,  $J$  = 2.6 Hz, 1H), 6.88 (dd,  $J$  = 7.6, 2.6 Hz, 1H), 3.93 (s, 3H), 2.23 (s, 3H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  209.4, 162.5, 153.6, 147.5, 146.9, 134.8, 133.1, 130.2, 128.3, 128.1, 127.1, 125.3, 109.3, 96.9, 55.9, 19.2. MS (EI)  $m/z$ : 282, 249, 206, 191, 134, 109, 79, 52. HRMS (ESI): calcd for  $\text{C}_{16}\text{H}_{15}\text{N}_2\text{OS}$  [ $\text{M} + \text{H}$ ] $^+$  283.0900; found 283.0903.

4-Chlorophenyl(7-methoxyimidazo[1,2-a]pyridin-3-yl)-methanethione (**4s**). Red solid (41 mg, 68%); mp 202–204 °C;  $R_f$  = 0.29 (ethyl acetate/petroleum ether = 1:3);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  10.66 (d,  $J$  = 7.6 Hz, 1H), 7.98 (s, 1H), 7.59–7.55 (m, 2H), 7.40–7.36 (m, 2H), 7.15 (d,  $J$  = 2.6 Hz, 1H), 6.88 (dd,  $J$  = 7.6, 2.6 Hz, 1H), 3.98 (s, 3H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  206.1, 162.7, 153.9, 146.7, 146.6, 136.4, 134.8, 129.9, 128.3, 128.2, 109.4, 96.9, 56.0. MS (EI)  $m/z$ : 302, 287, 269, 191, 176, 148, 108, 93, 69, 52. HRMS (ESI): calcd for  $\text{C}_{15}\text{H}_{12}\text{ClN}_2\text{OS}$  [ $\text{M} + \text{H}$ ] $^+$  303.0353; found 303.0358.

(6-Iodoimidazo[1,2-a]pyridin-3-yl)(o-tolyl)methanethione (**4t**). Red solid (54 mg, 72%); mp 107–109 °C;  $R_f$  = 0.43 (ethyl acetate/petroleum ether = 1:3);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  11.27 (s, 1H), 7.89 (dd,  $J$  = 9.2, 1.6 Hz, 1H), 7.75 (s, 1H), 7.62 (d,  $J$  = 9.2 Hz, 1H), 7.33–7.28 (m, 1H), 7.21 (dd,  $J$  = 10.5, 5.2 Hz, 3H), 2.22 (s, 3H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  213.8, 149.3, 147.6, 145.3, 139.4, 133.8, 133.0, 131.6, 130.4, 128.5, 127.1, 125.5, 118.9, 80.2, 19.4. MS (EI)  $m/z$ : 378, 345, 249, 218, 160, 134, 125, 115, 89, 77, 51. HRMS (ESI): calcd for  $\text{C}_{15}\text{H}_{12}\text{IN}_2\text{S}$  [ $\text{M} + \text{H}$ ] $^+$  378.9760; found 378.9765.

(4-Chlorophenyl)(6-iodoimidazo[1,2-a]pyridin-3-yl)-methanethione (**4u**). Red solid (52 mg, 65%); mp 165–167 °C;  $R_f$  = 0.56 (ethyl acetate/petroleum ether = 1:3);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  11.05 (d,  $J$  = 0.8 Hz, 1H), 8.01 (s, 1H), 7.89 (dd,  $J$  = 9.2, 1.6 Hz, 1H), 7.65 (d,  $J$  = 9.2 Hz, 1H), 7.57 (d,  $J$  = 8.5 Hz, 2H), 7.40 (d,  $J$  = 8.5 Hz, 2H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  209.8, 149.4, 146.5, 144.9, 139.4, 137.1, 133.8, 131.6, 129.9, 128.4, 118.9, 79.8. MS (EI)  $m/z$ : 397, 365, 287, 271, 235, 204, 160, 135, 118, 77, 50. HRMS (ESI): calcd for  $\text{C}_{14}\text{H}_9\text{ClIN}_2\text{S}$  [ $\text{M} + \text{H}$ ] $^+$  398.9214; found 398.9218.

## ■ ASSOCIATED CONTENT

### ● Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.joc.9b01188.

$^1\text{H}$  and  $^{13}\text{C}$  NMR spectra (PDF)

## ■ AUTHOR INFORMATION

### Corresponding Authors

\*E-mail: [chenzwang@126.com](mailto:chenzwang@126.com).

\*E-mail: [yemin811@gnnu.cn](mailto:yemin811@gnnu.cn)

### ORCID

Zhengwang Chen: 0000-0002-6795-9625

### Notes

The authors declare no competing financial interest.

## ■ ACKNOWLEDGMENTS

We are grateful for financial support from the NSF of Jiangxi Province (20171ACB21048), the NSF of Jiangxi Provincial Education Department (GJJ160924), and the Innovation Fund of Jiangxi Province (YC2018-S385).

## ■ REFERENCES

(1) (a) Bagdi, A. K.; Santra, S.; Monir, K.; Hajra, A. Synthesis of imidazo[1,2-a]pyridines: a decade update. *Chem. Commun.* **2015**, 51, 1555–1575. (b) Kumar, A.; Pericherla, K.; Kaswan, P.; Pandey, K.

Recent Developments in the Synthesis of Imidazo[1,2-a]pyridines. *Synthesis* **2015**, 47, 887–912. (c) Pires, M. J. D.; Poiera, D. L.; Marques, M. M. B. Metal-Catalyzed Cross-Coupling Reactions of Aminopyridines. *Eur. J. Org. Chem.* **2015**, 2015, 7197–7234.

(2) (a) Ulloora, S.; Shabaraya, R.; Adhikari, A. V. Facile synthesis of new imidazo[1,2-a]pyridines carrying 1,2,3-triazoles via click chemistry and their antiepileptic studies. *Bioorg. Med. Chem. Lett.* **2013**, 23, 3368–3372. (b) Ismail, M. A.; Brun, R.; Wenzler, T.; Tanius, F. A.; Wilson, W. D.; Boykin, D. W. Novel dicationic imidazo[1,2-a]pyridines and 5,6,7,8-tetrahydro-imidazo[1,2-a]pyridines as antiprotozoal agents. *J. Med. Chem.* **2004**, 47, 3658–3664. (c) Gudmundsson, K. S.; Johns, B. A. Synthesis of novel imidazo[1,2-a]pyridines with potent activity against herpesviruses. *Org. Lett.* **2003**, 5, 1369–1372. (d) Gudmundsson, K. S.; Williams, J. D.; Drach, J. C.; Townsend, L. B. Synthesis and antiviral activity of novel erythrofuransyl imidazo[1,2-a]pyridine C-nucleosides constructed via palladium coupling of iodoimidazo[1,2-a]pyridines and dihydrofuran. *J. Med. Chem.* **2003**, 46, 1449–1455.

(3) (a) Koo, H. L.; DuPont, H. L. Rifaximin: a unique gastrointestinal-selective antibiotic for enteric diseases. *Curr. Opin. Gastroenterol.* **2010**, 26, 17–25. (b) Mizushige, K.; Ueda, T.; Yukiiri, K.; Suzuki, H. Olprinone: a phosphodiesterase III inhibitor with positive inotropic and vasodilator effects. *Cardiovasc. Drug Rev.* **2002**, 20, 163–174. (c) Langer, S. Z.; Arbilla, S.; Benavides, J.; Scatton, B. Zolpidem and alpidem: two imidazopyridines with selectivity for omega 1- and omega 3-receptor subtypes. *Adv. Biochem. Psychopharmacol.* **1990**, 46, 61–72. (d) Almirante, L.; Polo, L.; Mugnaini, A.; Provinciali, E.; Rugarli, P.; Biancotti, A.; Gamba, A.; Murmann, W. Derivatives of Imidazole. I. Synthesis and Reactions of Imidazo[1,2-a]pyridines with Analgesic, Antiinflammatory, Antipyretic, and Anticonvulsant Activity. *J. Med. Chem.* **1965**, 8, 305–312.

(4) (a) Cao, H.; Lei, S.; Li, N.; Chen, L.; Liu, J.; Cai, H.; Qiu, S.; Tan, J. Cu-Catalyzed selective C3-formylation of imidazo[1,2-a]pyridine C-H bonds with DMSO using molecular oxygen. *Chem. Commun.* **2015**, 51, 1823–1825. (b) Kaswan, P.; Porter, A.; Pericherla, K.; Simone, M.; Peters, S.; Kumar, A.; DeBoef, B. Oxidative Cross-Coupling of sp<sup>3</sup>- and sp<sup>2</sup>-Hybridized C-H Bonds: Vanadium-Catalyzed Aminomethylation of Imidazo[1,2-a]pyridines. *Org. Lett.* **2015**, 17, 5208–5211. (c) Lei, S.; Cao, H.; Chen, L.; Liu, J.; Cai, H.; Tan, J. Regioselective Oxidative Homocoupling Reaction: An Efficient Copper-Catalyzed Synthesis of Biimidazo[1,2-a]pyridines. *Adv. Synth. Catal.* **2015**, 357, 3109–3114. (d) Lei, S.; Chen, G.; Mai, Y.; Chen, L.; Cai, H.; Tan, J.; Cao, H. Regioselective Copper-Catalyzed Oxidative Cross-Coupling of Imidazo[1,2-a]pyridines with Methyl Ketones: An Efficient Route for Synthesis of 1,2-Diketones. *Adv. Synth. Catal.* **2016**, 358, 67–73. (e) Lei, S.; Mai, Y.; Yan, C.; Mao, J.; Cao, H. A Carbonylation Approach Toward Activation of Csp<sup>2</sup>-H and Csp<sup>3</sup>-H Bonds: Cu-Catalyzed Regioselective Cross Coupling of Imidazo[1,2-a]pyridines with Methyl Hetarenes. *Org. Lett.* **2016**, 18, 3582–3585. (f) Wang, C.; Lei, S.; Cao, H.; Qiu, S.; Liu, J.; Deng, H.; Yan, C. Regioselective Copper-Catalyzed Dicarboxylation of Imidazo[1,2-a]pyridines with N,N-Disubstituted Acetamide or Acetone: An Approach to 1,2-Diketones Using Molecular Oxygen. *J. Org. Chem.* **2015**, 80, 12725–12732.

(5) (a) Chernyak, N.; Gevorgyan, V. General and efficient copper-catalyzed three-component coupling reaction towards imidazoheterocycles: one-pot synthesis of alpidem and zolpidem. *Angew. Chem., Int. Ed.* **2010**, 49, 2743–2746. (b) McDonald, I. M.; Peese, K. M. General Method for the Preparation of Electron-Deficient Imidazo[1,2-a]pyridines and Related Heterocycles. *Org. Lett.* **2015**, 17, 6002–6005. (c) Rassokhina, I. V.; Shirinian, V. Z.; Zavarzin, I. V.; Gevorgyan, V.; Volkova, Y. A. Copper(II)-Mediated Aerobic Synthesis of Imidazo[1,2-a]pyridines via Cascade Aminomethylation/Cycloisomerization of Alkynes. *J. Org. Chem.* **2015**, 80, 11212–11218. (d) Wang, H.; Wang, Y.; Liang, D.; Liu, L.; Zhang, J.; Zhu, Q. Copper-catalyzed intramolecular dehydrogenative aminooxygenation: direct access to formyl-substituted aromatic N-heterocycles. *Angew. Chem., Int. Ed.* **2011**, 50, 5678–5681. (e) Wang, H.; Wang, Y.; Peng, C.; Zhang, J.; Zhu, Q. Direct Intramolecular C-H Amination Reaction



Cocatalyzed by Copper(II) and Iron(III) as Part of an Efficient Route for the Synthesis of Pyrido[1,2-a]benzimidazoles from N-Aryl-2-aminopyridines. *J. Am. Chem. Soc.* **2010**, *132*, 13217–13219. (f) Zhou, X.; Yan, H.; Ma, C.; He, Y.; Li, Y.; Cao, J.; Yan, R.; Huang, G. Copper-Mediated Aerobic Oxidative Synthesis of 3-Bromo-imidazo[1,2-a]pyridines with Pyridines and Enamides. *J. Org. Chem.* **2016**, *81*, 25–31.

(6) For selected examples, see: (a) Nguyen, T. B.; Retaillieu, P. Elemental Sulfur as Reaction Medium for the Synthesis of Fused Nitrogen Heterocycles by Oxidative Coupling between Cycloalkanones and Nitrogen Nucleophiles. *Adv. Synth. Catal.* **2017**, *359*, 3843–3847. (b) Huo, C.; Tang, J.; Xie, H.; Wang, Y.; Dong, J. CBr<sub>4</sub> Mediated Oxidative C–N Bond Formation: Applied in the Synthesis of Imidazo[1,2-a]pyridines and Imidazo[1,2-a]pyrimidines. *Org. Lett.* **2016**, *18*, 1016–1019. (c) Xiao, X.; Xie, Y.; Bai, S.; Deng, Y.; Jiang, H.; Zeng, W. Transition-Metal-Free Tandem Chlorocyclization of Amines with Carboxylic Acids: Access to Chloroimidazo[1,2-a]pyridines. *Org. Lett.* **2015**, *17*, 3998–4001. (d) Nguyen, T. B.; Corbin, M.; Retaillieu, P.; Ermolenko, L.; Al-Mourabit, A. Elements as Direct Feedstocks for Organic Synthesis: Fe/I<sub>2</sub>/O<sub>2</sub> for Diamination of 2-Cyclohexenones with 2-Aminopyrimidine and 2-Aminopyridines. *Org. Lett.* **2015**, *17*, 4956–4959. (e) Lee, S. K.; Park, J. K. Room-temperature transition-metal-free one-pot synthesis of 3-aryl imidazo[1,2-a]pyridines via iodo-hemiaminal intermediate. *J. Org. Chem.* **2015**, *80*, 3723–3729. (f) Adiyala, P. R.; Mani, G. S.; Nanubolu, J. B.; Shekar, K. C.; Maurya, R. A. Access to Imidazo[1,2-a]pyridines via Annulation of  $\alpha$ -Keto Vinyl Azides and 2-Aminopyridines. *Org. Lett.* **2015**, *17*, 4308–4311. (g) Chandra Mohan, D.; Reddy Donthiri, R.; Nageswara Rao, S.; Adimurthy, S. Copper(I) Iodide-Catalysed Aerobic Oxidative Synthesis of Imidazo[1,2-a]pyridines from 2-Aminopyridines and Methyl Ketones. *Adv. Synth. Catal.* **2013**, *355*, 2217–2221. (h) Cai, Z.-J.; Wang, S.-Y.; Ji, S.-J. Copper(I) Iodide/Boron Trifluoride Etherate-Cocatalyzed Aerobic Dehydrogenative Reactions Applied in the Synthesis of Substituted Heteroaromatic Imidazo[1,2-a]pyridines. *Adv. Synth. Catal.* **2013**, *355*, 2686–2692. (i) Bagdi, A. K.; Rahman, M.; Santra, S.; Majee, A.; Hajra, A. Copper-Catalyzed Synthesis of Imidazo[1,2-a]pyridines through Tandem Imine Formation-Oxidative Cyclization under Ambient Air: One-Step Synthesis of Zolimidine on a Gram-Scale. *Adv. Synth. Catal.* **2013**, *355*, 1741–1747. (j) Ma, L.; Wang, X.; Yu, W.; Han, B. TBAI-catalyzed oxidative coupling of aminopyridines with  $\beta$ -keto esters and 1,3-diones-synthesis of imidazo[1,2-a]pyridines. *Chem. Commun.* **2011**, *47*, 11333–11335.

(7) (a) Yan, R. L.; Yan, H.; Ma, C.; Ren, Z. Y.; Gao, X. A.; Huang, G. S.; Liang, Y. M. Cu(I)-catalyzed synthesis of imidazo[1,2-a]pyridines from aminopyridines and nitroolefins using air as the oxidant. *J. Org. Chem.* **2012**, *77*, 2024–2028. (b) Santra, S.; Bagdi, A. K.; Majee, A.; Hajra, A. Iron(III)-Catalyzed Cascade Reaction between Nitroolefins and 2-Aminopyridines: Synthesis of Imidazo[1,2-a]pyridines and Easy Access towards Zolimidine. *Adv. Synth. Catal.* **2013**, *355*, 1065–1070. (c) Nair, D. K.; Mobin, S. M.; Namboothiri, I. N. N. Synthesis of Imidazopyridines from the Morita-Baylis-Hillman Acetates of Nitroalkenes and Convenient Access to Alpidem and Zolpidem. *Org. Lett.* **2012**, *14*, 4580–4583. (d) Monir, K.; Bagdi, A. K.; Ghosh, M.; Hajra, A. Unprecedented catalytic activity of Fe(NO<sub>3</sub>)<sub>3</sub>·9H<sub>2</sub>O: regioselective synthesis of 2-nitroimidazopyridines via oxidative amination. *Org. Lett.* **2014**, *16*, 4630–4633.

(8) (a) Gao, Y.; Yin, M.; Wu, W.; Huang, H.; Jiang, H. Copper-Catalyzed Intermolecular Oxidative Cyclization of Halo-alkynes: Synthesis of 2-Halo-substituted Imidazo[1,2-a]pyridines, Imidazo[1,2-a]pyrazines and Imidazo[1,2-a]pyrimidines. *Adv. Synth. Catal.* **2013**, *355*, 2263–2273. (b) Zeng, J.; Tan, Y. J.; Leow, M. L.; Liu, X.-W. Copper(II)/Iron(III) Co-catalyzed Intermolecular Diamination of Alkynes: Facile Synthesis of Imidazopyridines. *Org. Lett.* **2012**, *14*, 4386–4389. (c) He, C.; Hao, J.; Xu, H.; Mo, Y.; Liu, H.; Han, J.; Lei, A. Heteroaromatic imidazo[1,2-a]pyridines synthesis from C–H/N–H oxidative cross-coupling/cyclization. *Chem. Commun.* **2012**, *48*, 11073–11075.

(9) (a) Boyd, D. A. Sulfur and Its Role In Modern Materials Science. *Angew. Chem., Int. Ed.* **2016**, *55*, 15486–15502. (b) Cinar, M. E.; Ozturk, T. Thienothiophenes, dithienothiophenes, and thienoacenes: syntheses, oligomers, polymers, and properties. *Chem. Rev.* **2015**, *115*, 3036–3140. (c) Ilardi, E. A.; Vitaku, E.; Njardarson, J. T. Data-mining for sulfur and fluorine: an evaluation of pharmaceuticals to reveal opportunities for drug design and discovery. *J. Med. Chem.* **2014**, *57*, 2832–2842. (d) Jiang, W.; Li, Y.; Wang, Z. Heteroarenes as high performance organic semiconductors. *Chem. Soc. Rev.* **2013**, *42*, 6113–6127.

(10) For selected reviews and examples, see: (a) Nguyen, T. B. Recent Advances in Organic Reactions Involving Elemental Sulfur. *Adv. Synth. Catal.* **2017**, *359*, 1066–1130. (b) Qiao, Z.; Jiang, X. Recent developments in sulfur-carbon bond formation reaction involving thiosulfates. *Org. Biomol. Chem.* **2017**, *15*, 1942–1946. (c) Liu, H.; Jiang, X. Transfer of sulfur: from simple to diverse. *Chem. - Asian J.* **2013**, *8*, 2546–2563.

(11) (a) Cao, H.; Liu, X.; Zhao, L.; Cen, J.; Lin, J.; Zhu, Q.; Fu, M. One-pot regioselective synthesis of imidazo[1,2-a]pyridines: a novel, metal-free, three-component reaction for the formation of C–N, C–O, and C–S bonds. *Org. Lett.* **2014**, *16*, 146–149. (b) Zhan, H.; Cao, H.; Qiu, H.; Li, N.; Chen, L.; Liu, J.; Cai, H.; Tan, J. Microwave-assisted C–N and C–S bond-forming reactions: an efficient three-component domino sequence for the synthesis of sulfoether-decorated imidazo[1,2-a]pyridines. *RSC Adv.* **2015**, *5*, 32205–32209. (c) Zhang, H.; Jiang, L. Microwave-assisted solvent-free synthesis of imidazo[1,2-a]pyridines via a three-component reaction. *Tetrahedron Lett.* **2015**, *56*, 2777–2779.

(12) (a) Cao, H.; Liu, X.; Liao, J.; Huang, J.; Qiu, H.; Chen, Q.; Chen, Y. Transition metal-mediated C=O and C=C bond-forming reactions: a regioselective strategy for the synthesis of imidazo[1,2-a]pyridines and imidazo[1,2-a]pyrazines. *J. Org. Chem.* **2014**, *79*, 11209–11214. (b) Zhan, H.; Zhao, L.; Liao, J.; Li, N.; Chen, Q.; Qiu, S.; Cao, H. Gold-Catalyzed Synthesis of 3-Acylimidazo[1,2-a]pyridines via Carbene Oxidation. *Adv. Synth. Catal.* **2015**, *357*, 46–50. (c) Reddy, K. R.; Gupta, A. P.; Das, P. Metal-Free Amino-oxygenation of Alkynes: Efficient Synthesis of 3-Aroylimidazo[1,2-a]-N-Heterocycles. *Asian J. Org. Chem.* **2016**, *5*, 900–906. (d) Chen, Z.; Liu, B.; Liang, P.; Yang, Z.; Ye, M. Iron(III)-catalyzed synthesis of 3-aryylimidazo[1,2-a]pyridines from 2-aminopyridines and ynals. *Tetrahedron Lett.* **2018**, *59*, 667–670.

(13) Cui, Z.; Zhu, B.; Li, X.; Cao, H. Access to sulfonylated furans or imidazo[1,2-a]pyridines via a metal-free three-component, domino reaction. *Org. Chem. Front.* **2018**, *5*, 2219–2223.

(14) Kobayashi, S.; Sugiura, M.; Kitagawa, H.; Lam, W. W. L. Rare-earth metal triflates in organic synthesis. *Chem. Rev.* **2002**, *102*, 2227–2302.

(15) Qiu, Y. F.; Yang, F.; Qiu, Z. H.; Zhong, M. J.; Wang, L. J.; Ye, Y. Y.; Song, B.; Liang, Y. M. Bronsted acid catalyzed and NIS-promoted cyclization of diynones: selective synthesis of 4-pyrone, 4-pyridone, and 3-pyrrolone derivatives. *J. Org. Chem.* **2013**, *78*, 12018–12028.

(16) (a) Zhu, C.-Z.; Sun, Y. L.; Wei, Y.; Shi, M. Phosphine-Mediated Dimerization of Conjugated Ene-Yne Ketones: Stereoselective Construction of Dihydrobenzofurans. *Adv. Synth. Catal.* **2017**, *359*, 1263–1270. (b) Song, B.; Li, L. H.; Song, X. R.; Qiu, Y. F.; Zhong, M. J.; Zhou, P. X.; Liang, Y. M. Zinc-catalyzed [4 + 3] cycloaddition with concomitant furan annulation: formation of cyclohepta[b]furans. *Chem. - Eur. J.* **2014**, *20*, 5910–5913. (c) Ji, G.; Wang, X.; Zhang, S.; Xu, Y.; Ye, Y.; Li, M.; Zhang, Y.; Wang, J. Synthesis of 3-trifluoromethylpyrazoles via trifluoromethylation/cyclization of  $\alpha,\beta$ -alkynic hydrazones using a hypervalent iodine reagent. *Chem. Commun.* **2014**, *50*, 4361–4363. (d) Romanov-Mikhailidis, F.; Besnard, C.; Alexakis, A. N-Heterocyclic Carbene-Catalyzed Annulation of  $\alpha$ -Cyano-1,4-diketones with Ynals. *Org. Lett.* **2012**, *14*, 4906–4909. (e) Kim, H. Y.; Lantrip, D. A.; Fuchs, P. L. Relative Reactivity of *anti*- and *syn*-Oximino Carbonates and Carbamates of 2-Pyridylacetic Acid Esters. *Org. Lett.* **2001**, *3*, 2137–2140.