

---

# **BASICS OF DIFFERENTIAL GEOMETRY 2**

---

**Notes of BIMSA course**

**Misuzu/Yuxuan Liu**

February 24, 2025

## Contents

<b>1</b>	<b>Principle Bundles</b>	<b>2</b>
1.1	Lie Groups . . . . .	2
1.2	Definition of Principle Bundles . . . . .	7
1.3	Connections on Principle Bundles . . . . .	9

## Introduction

Last semester:

- Geometry of vector bundles
- Basic Riemannian geometry
- Differential operators on manifolds

We will learn this semester:

- Theory of principle bundles
- characteristic classes
- Basics of complex manifold, Kähler manifold, symplectic manifold.

# 1 Principle Bundles

In this section, we introduce the connections of principle bundles, it's closely related to the connections of vector bundles.

## 1.1 Lie Groups

**Definition 1.1.** Let  $G$  be a smooth manifold.  $G$  is a *Lie group* if  $G$  is a group s.t. multiplication and inverse are smooth.

Let  $G$  be a Lie group,  $g \in G$ , we denote:

- $L_g : G \rightarrow G, h \mapsto gh$  (left translation)
- $R_g : G \rightarrow G, h \mapsto hg$  (right translation)
- $\mathfrak{X}^L(G) = \{X \in \mathfrak{X}(G) \mid \forall g \in G, (L_g)_*X = X\}$  (left invariant vector fields)

For  $X \in \mathfrak{X}^L(G)$ ,  $L_{g*}X = X$  means that  $X$  is  $L_g$ -related to  $X$ . Then for  $\forall X, Y \in \mathfrak{X}^L(G)$ ,  $L_{g*}([X, Y]) = [L_{g*}X, L_{g*}Y] = [X, Y]$ , so  $\mathfrak{X}^L(G)$  is closed under  $[\cdot, \cdot]$

**Definition 1.2.** Set  $\mathbb{K} = \mathbb{R}$  or  $\mathbb{C}$ . Given a  $\mathbb{K}$ -vector space  $\mathfrak{g}$  and a bilinear map  $[\cdot, \cdot] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$ , we say  $\mathfrak{g}$  is a *Lie algebra* if:

- (1)  $\forall X, Y \in \mathfrak{g}, [X, Y] = -[Y, X]$
  - (2)  $\forall X, Y, Z \in \mathfrak{g}, [[X, Y], Z] + [[Y, Z], X] + [[Z, X], Y] = 0$
- $[\cdot, \cdot]$  is called Lie bracket.

So by definition we have  $(\mathfrak{X}^L(G), [\cdot, \cdot])$  is a Lie algebra.

**Definition 1.3.** For Lie algebra  $\mathfrak{g}, \mathfrak{h}$ , a linear map  $f : \mathfrak{g} \rightarrow \mathfrak{h}$  is called the *Lie algebra homomorphism* if:  $\forall X, Y \in \mathfrak{g}, f([X, Y]) = [f(X), f(Y)]$

If  $f$  is in addition an isomorphism, then  $f$  is called a *Lie algebra isomorphism*.

Let  $e \in G$  be the unit of  $G$ . Set  $\iota : \mathfrak{X}^L(G) \rightarrow T_eG, X \mapsto X_e$ . Then  $\iota$  is a linear isomorphism. Let  $\mathfrak{g} = T_eG$ , so we can define the Lie bracket on  $\mathfrak{g}$  s.t.  $\iota$  is a Lie algebra isomorphism, i.e. setting  $X^\# = \iota^{-1}(X)$ ,  $[X, Y] = [X^\#, Y^\#]_e$ . Note that  $X_g^\# = (L_g)_*X_e$ ,  $g \in G$ .

**Definition 1.4.** Let  $G$  be Lie group,  $\mathfrak{g} = T_eG$  with  $[\cdot, \cdot]$  is called the *Lie algebra of  $G$* .  $(\mathfrak{X}^L(G), [\cdot, \cdot])$  is also called the Lie algebra of  $G$

**Definition 1.5.** Let  $G, H$  be Lie groups. A map  $\rho : G \rightarrow H$  is a *Lie group homomorphism* if  $\rho$  is a smooth map and a group homomorphism. For the special

case  $(\mathbb{R}, +) \rightarrow G, t \mapsto g_t, \{g_t\}_{t \in \mathbb{R}}$  is called *one parameter subgroup* of  $G$ .

**Proposition 1.1.** Let  $G$  be Lie group and  $\mathfrak{g}$  its Lie algebra. Then

- (1)  $\forall X \in \mathfrak{g}, X^\# = \iota^{-1}(X)$  is complete, i.e.  $X^\#$  generates a flow  $\{\varphi_t\}_{t \in \mathbb{R}}$ .
- (2) Set  $\exp_G(tX) = \varphi_t(e) \in G$ . Then  $\varphi_t = R_{\exp_G(tX)}$ .
- (3) For  $s, t \in \mathbb{R}, \exp_G(sX) \exp_G(tX) = \exp_G((s+t)X)$ , i.e.  $\{\exp_G(tX)\}_{t \in \mathbb{R}}$  is one parameter subgroup of  $G$ .
- (4)  $\mathfrak{g} \rightarrow \{\text{one parameter subgroup of } G\}, X \mapsto \{\exp_G(tX)\}_{t \in \mathbb{R}}$  is bijective.

*Proof.* (1) By ODE theory,  $\exists \epsilon > 0, \gamma_e : (-\epsilon, \epsilon) \rightarrow G$  s.t.  $\gamma_e(0) = e, \frac{d\gamma_e}{dt} = X^\#_{\gamma_e(t)}$ .

**Claim 1.**  $\forall g \in G$ , define  $\gamma_g : (-\epsilon, \epsilon) \rightarrow G, t \mapsto g\gamma_e(t)$  is the integral curve of  $X^\#$  with  $\gamma_g(0) = g$ .

Indeed,  $\forall t \in (-\epsilon, \epsilon), \frac{d\gamma_g}{dt}(t) = (L_g)_* \gamma_e(t) \frac{d\gamma_e}{dt}(t) = X^\#_{g\gamma_e(t)}$ .

**Claim 2.**  $\gamma_e : (-\epsilon, \epsilon) \rightarrow G$  can be extended to integral curve  $\gamma_e : \mathbb{R} \rightarrow G$  of  $X^\#$  with  $\gamma_e(0) = e$ .

Set  $\varphi_t = R_{\gamma_e(t)}$ , then  $\{\varphi_t\}_{t \in \mathbb{R}}$  is the flow generated by  $X^\#$ . So the following are easy.  $\square$

By this proposition, we can define the exponential map  $\exp_G : \mathfrak{g} \rightarrow G$ .

**Proposition 1.2.** Let  $G, H$  be Lie groups with Lie algebra  $\mathfrak{g}, \mathfrak{h}$ . If  $f : G \rightarrow H$  is Lie group homomorphism, then  $f_* : \mathfrak{g} \rightarrow \mathfrak{h}$  is a Lie algebra homomorphism.

*Proof.* We only need to show that  $X^\#$  and  $(f_*X)^\#$  are  $f$ -related. Since  $X = \frac{d}{dt} \exp_G(tX)|_{t=0}$ , we have  $f_*(X^\#) = \frac{d}{dt} f(g \cdot \exp_G(tX))|_{t=0} = \frac{d}{dt} f(g) f(\exp_G(tX))|_{t=0} = (L_{f(g)})_* (f_*X) = (f_*X)^\#_{f(g)}$ .  $\square$

**Example 1.1.** Let  $V$  be a  $\mathbb{R}$ -vector space,  $G = GL(V)$ ,  $\mathfrak{g}$  Lie algebra of  $G$ . Then  $\mathfrak{g} = \text{End}(V)$ , the bracket is given as follows:

**Proposition 1.3.**  $\forall X, Y \in \text{End}(V), [X, Y] = XY - YX$ .

*Proof.* For  $X \in \text{End}(V)$ , set matrix exponential  $e^{tX} = \sum_{k=0}^{\infty} \frac{(tX)^k}{k!}$ . Then  $\{e^{tX}\}_{t \in \mathbb{R}}$  is a one parameter subgroup of  $G$  and  $\frac{d}{dt} e^{tX}|_{t=0} = X$ . So  $\exp_G(tX) = e^{tX}$ .

Then  $[X, Y] = [X^\#, Y^\#]_e = (\mathcal{L}_{X^\#} Y^\#)_e = \frac{d}{dt} (\varphi_{-t})_* e^{tX} (Y^\#_{e^{tX}})|_{t=0} = \frac{d}{dt} \frac{d}{ds} \varphi_{-t} (e^{tX} e^{sY})|_{s=t=0} = XY - YX$ .  $\square$

**Example 1.2.** Set

- $O(n) = \{g \in GL(n; \mathbb{R}) \mid g^t g = E_n\}$  (orthogonal group)
- $SO(n) = \{g \in O(n) \mid \det g = 1\}$  (special orthogonal group)

we can check that  $O(n), SO(n)$  are Lie subgroups of  $GL(n; \mathbb{R})$ .

$SO(n)$  is the unit component of  $O(n)$ , so  $\mathfrak{o}(n) = \mathfrak{so}(n)$  (Lie algebra of  $(O(n))$  and  $SO(n)$ ). This is a Lie subalgebra of  $End(\mathbb{R}^n)$  given by

$$\mathfrak{o}(n) = \mathfrak{so}(n) = \{X \in End(\mathbb{R}^n) \mid X^t + X = O_n\}$$

where  $O_n$  is the zero matrix of size  $n$ .

Similarly, set

- $U(n) = \{g \in GL(n; \mathbb{C}) \mid g^* g = E_n\}$  (unitary group) where  $g^* = \overline{g^t}$
- $SU(n) = \{g \in U(n) \mid \det g = 1\}$  (special unitary group)

We can check that

- $U(n), SU(n)$  are Lie subgroups of  $GL(n; \mathbb{C})$
- $\mathfrak{u}(n) = \{X \in End(\mathbb{C}^n) \mid X^* + X = O\}$  (Lie algebra of  $U(n)$ )
- $\mathfrak{su}(n) = \{X \in End(\mathbb{C}^n) \mid X^* + X = O, \text{tr} X = 0\}$  (Lie algebra of  $SU(n)$ )

**Note.** A Lie subgroup  $H$  of  $G$  is a Lie group s.t.

- $H$  is a subset of  $G$
- inclusion map  $H \hookrightarrow G$  is an embedding and group homomorphism

**Fact.** A closed subgroup of  $G$  is a Lie subgroup of  $G$ .

**Definition 1.6.** Let  $V$  be a  $\mathbb{K}$ -vector space,  $G$  a Lie group. A Lie group homomorphism  $\rho : G \rightarrow GL(V)$  is called a *representation of  $V$* . The Lie algebra homomorphism  $\rho_{*e} : \mathfrak{g} \rightarrow End(V)$  is called a *differential representation*.

**Example 1.3.** Let  $G$  be a Lie group,  $\mathfrak{g}$  its Lie algebra.  $\forall g \in G$ , define a homomorphism

$$F_g : G \rightarrow G, h \mapsto ghg^{-1}$$

Note that  $F_g \circ F_{g'} = F_{gg'}$ . This induces a Lie algebra homomorphism  $(F_g)_{*e} : \mathfrak{g} \rightarrow \mathfrak{g}$  which satisfies  $(F_g)_{*e} \circ (F_{g'})_{*e} = (F_{gg'})_{*e}$ . So we obtain a representation

$$Ad : G \rightarrow GL(\mathfrak{g}), g \mapsto (F_g)_{*e}$$

called adjoint representation of  $G$ . The differential representation  $ad : \mathfrak{g} \rightarrow End(\mathfrak{g})$  of  $Ad$  is given as follows.

**Proposition 1.4.**  $\forall X, Y \in \mathfrak{g}, ad(X)(Y) = [X, Y]$ .

*Proof.* Note that  $F_g = R_{g^{-1}} \circ L_g$ . Then

$$ad(X)(Y) = \frac{d}{dt} Ad(\exp_G(tX))(Y)|_{t=0} = \frac{d}{dt} (R_{\exp_G(-tX)})_{*\exp_G(tX)} (L_{\exp_G(tX)})_{*e} (Y)|_{t=0} = [X^\sharp, Y^\sharp]_e = [X, Y]. \quad \square$$

Recall that there is a exponential map in Riemannian geometry. The Riemannian exp and the Lie group exp are related as follows.

**Definition 1.7.** A Riemannian metric  $\langle \cdot, \cdot \rangle$  on a Lie group  $G$  is said to be *bi-invariant* if  $\forall g, h \in G, L_g^* R_h^* \langle \cdot, \cdot \rangle = \langle \cdot, \cdot \rangle$ .

**Theorem 1.1.** Let  $G$  be a Lie group with a bi-invariant metric  $\langle \cdot, \cdot \rangle$ . Then  $\exp_e = \exp_G$ .

To show this we describe the Levi-Civita connection  $\nabla$  of  $\langle \cdot, \cdot \rangle$ .

**Lemma 1.1.**  $\forall X, Y \in \mathfrak{g}, \nabla_{X^\sharp} Y^\sharp = \frac{1}{2}[X, Y]^\sharp$ .

*Proof.* By Koszul formula, we have

$$\begin{aligned} \langle \nabla_{X^\sharp} Y^\sharp, Z^\sharp \rangle &= \frac{1}{2} \left( X^\sharp \langle Y^\sharp, Z^\sharp \rangle + Y^\sharp \langle Z^\sharp, X^\sharp \rangle - Z^\sharp \langle X^\sharp, Y^\sharp \rangle \right. \\ &\quad \left. - \langle Y^\sharp, [X^\sharp, Z^\sharp] \rangle - \langle Z^\sharp, [Y^\sharp, X^\sharp] \rangle + \langle X^\sharp, [Z^\sharp, Y^\sharp] \rangle \right) \end{aligned}$$

Since for  $\forall g \in G, X_g^\sharp = \frac{d}{dt} g \cdot \exp_G(tX)|_{t=0}$ , we have

$$X^\sharp \langle Y^\sharp, Z^\sharp \rangle = \frac{d}{dt} \langle Y_{g \cdot \exp_G(tX)}^\sharp, Z_{g \cdot \exp_G(tX)}^\sharp \rangle_{g \cdot \exp_G(tX)}|_{t=0} = \frac{d}{dt} \langle Y, Z \rangle_e|_{t=0} = 0$$

Since  $\langle \cdot, \cdot \rangle$  is bi-invariant,

$$L_g^* R_{g^{-1}}^* \langle \cdot, \cdot \rangle_e = \langle \cdot, \cdot \rangle_e \text{ for } \forall g \in G \iff \langle Ad(g)(\cdot), Ad(g)(\cdot) \rangle_e = \langle \cdot, \cdot \rangle_e$$

Setting  $g = \exp_G(tZ)$  and  $\frac{d}{dt}|_{t=0}$ , we have  $\langle ad(Z)(\cdot), \cdot \rangle_e + \langle \cdot, ad(Z)(\cdot) \rangle_e = 0$ , which shows that  $\langle Y^\sharp, [X^\sharp, Z^\sharp] \rangle + \langle X^\sharp, [Z^\sharp, Y^\sharp] \rangle = 0$ , so we have  $\nabla_{X^\sharp} Y^\sharp = \frac{1}{2}[X, Y]^\sharp$ .  $\square$

The proof of the theorem completes once shown that  $\exp_G(tX)$  is geodesic, which is left as an exercise.

**Exercise 1.1.** Prove the theorem.

**Remark 1.1.** Existence/uniqueness of bi-invariant metrics? Some facts from representation theory are needed, the argument here is not used after this remark.

Existence When  $G$  is compact,  $\exists$  bi-invariant metric using “averaging trick”.

- We first define  $Ad$ -invariant inner product on  $\mathfrak{g}$ .
- Then extend it to the whole  $G$  by pulling back  $L_g$ .

**Note:**  $\exists$  bi-invariant on  $G \iff \exists$   $Ad$ -invariant inner product on  $\mathfrak{g}$ .

$\left\{ \begin{array}{l} (\Rightarrow) \text{ Trivial.} \\ (\Leftarrow) \text{ Given } Ad\text{-invariant inner product on } \mathfrak{g}, \text{ we can extend it to left-invariant metric} \\ \text{on } G, \text{ this is also right-invariant by pullback of } R_h = R_h \circ L_{h^{-1}} \circ L_h = Ad(h^{-1}) \circ L_h \end{array} \right.$

**Uniqueness** When  $G$  is abelian, then  $L_g = R_g$ , so  $\exists$  many bi-invariant metrics on  $G$  (Any inner product on  $\mathfrak{g}$  induces left-invariant metric on  $\mathfrak{g}$ , by the note above it is bi-invariant). Suppose that  $\exists$   $Ad$ -invariant inner product  $\langle \cdot, \cdot \rangle$  on  $\mathfrak{g}$ . By  $\langle \cdot, \cdot \rangle$ , we have an irreducible decomposition of  $(\mathfrak{g}, Ad)$ :  $\mathfrak{g} = \mathfrak{g}_1^{\oplus n_1} \oplus \cdots \oplus \mathfrak{g}_r^{\oplus n_r}$ , where  $\mathfrak{g}_i$  is irreducible representation of  $G$  and  $\mathfrak{g}_i \neq \mathfrak{g}_j$  for  $i \neq j$ . Then

$$\dim \{ Ad\text{-invariant symmetric bilinear map } \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{R} \} = \sum_{i=1}^r n_i^2$$

To see this, take  $T \in \{ Ad\text{-invariant symmetric bilinear map} \}$  and use Schur's lemma to

$$T_{ij} : \mathfrak{g}_i \hookrightarrow \mathfrak{g} \xrightarrow{x \mapsto T(x, \cdot)} \mathfrak{g}^* \xrightarrow{\langle \cdot, \cdot \rangle} \mathfrak{g} \xrightarrow{proj.} \mathfrak{g}_j$$

Then  $T_{ij} = \begin{cases} 0 & (i \neq j) \\ c \cdot id & (i = j) \text{ for } \exists c \in \mathbb{R} \end{cases}$ , so uniqueness up to scalar multiplication holds

only when  $r = 1, n = 1$ , i.e.  $(\mathfrak{g}, Ad)$  is irreducible  $\iff G$  is simple Lie group.

**Definition 1.8.** Let  $M$  be smooth manifold,  $G$  be Lie group with unit  $e$ . A smooth map

$$A : M \times G \rightarrow M, (x, g) \mapsto xg$$

is called the *right action of  $G$  on  $M$*  if

- (1)  $\forall x \in M, xe = x$
- (2)  $\forall x \in M, \forall g, h \in G, (xg)h = x(gh)$

We write the right action as  $M \curvearrowright G$ .

**Definition 1.9.** Suppose  $M \curvearrowright G$ .

- (1) For  $\forall g \in G$ , set  $R_g : M \rightarrow M, X \mapsto xg$  (*right translation*).

- (2) For  $\forall X \in \mathfrak{g}$ , define the *fundamental vector field*  $X^\# \in \mathfrak{X}(M)$  by  $X_x^\# = \frac{d}{dt} x \cdot \exp_G(tX) |_{t=0} = dA(x, \cdot)_e(X)$ .

Here the notation  $X^\#$  is the same as the left-invariant vector field on Lie group, we'll show that they have the same property:



**Remark 1.2.** (1)  $\forall g \in G, \forall X \in \mathfrak{g}, (R_g)_* X^\# = (Ad(g^{-1})X)^\#$ .  
 (2)  $\forall X, Y \in \mathfrak{g}, [X^\#, Y^\#] = [X, Y]^\#$ .

*Proof.* (1)  $\forall x \in M, ((R_g)_* X^\#)_x = (R_g)_* X^\#_{xg^{-1}} = \frac{d}{dt} xg^{-1} \exp_G(tX)g \big|_{t=0}$ . Since  $\{g^{-1} \exp_G(tX)g\}_{t \in \mathbb{R}}$  is a one parameter subgroup of  $G$  with  $\frac{d}{dt} g^{-1} \exp_G(tX)g \big|_{t=0} = Ad(g^{-1})X$ , then  $g^{-1} \exp_G(tX)g = \exp_G(tAd(g^{-1})X)$ , which gives (1).

(2) By definition,  $\{\varphi_t = R_{\exp_G(tX)}\}_{t \in \mathbb{R}}$  is flow of  $X^\#$ . So

$$[X^\#, Y^\#] = \frac{d}{dt} (\varphi_{-t})_* Y^\# \big|_{t=0} = \frac{d}{dt} (Ad(\exp_G(tX)) Y)^\# \big|_{t=0} = (ad(X)(Y))^\# = [X, Y]^\#.$$

□

**Remark 1.3.** We can define the left action

$$A^L : G \times M \rightarrow M, (g, x) \mapsto gx$$

and also the fundamental vector field  $X_L^\# \in \mathfrak{X}(M)$ . The left and right actions are essentially the same, since the right action is given from the left action. Indeed, given  $A^L$  above, define  $A$  by  $A(x, g) = A^L(g^{-1}, x) = g^{-1}x$ , then  $X_L^\# = -X^\#$  for  $X \in \mathfrak{g}$ .  $[X_L^\#, Y_L^\#] = [X, Y]^\# = -[X, Y]^\#$ .

**Definition 1.10.** Suppose  $M \curvearrowright G$ .

- (1) For  $p \in M$ , define  $G_p = \{g \in G \mid pg = p\}$  (*isotropy subgroup at p*).
- (2) The  $G$  action is *free* if  $G_p = \{e\}$  for  $\forall p \in M$ .
- (3) The  $G$  action is *effective* if  $\bigcap_{p \in M} G_p = \{e\}$ . In other words,  $G \rightarrow \text{Diff}(M)$  is injective.

## 1.2 Definition of Principle Bundles

**Definition 1.11.** Let  $P, M$  be smooth manifolds and  $G$  be Lie group. The map  $\pi_P : P \rightarrow M$  is a *principle  $G$ -bundle* or *principle bundle with structure group  $G$*  if:

- (1)  $P \curvearrowright G$ .
- (2) There exists an open cover  $\{U_\alpha\}_{\alpha \in A}$  of  $M$  and diffeomorphisms called local trivialization

$$\phi_\alpha : \pi_P^{-1}(U_\alpha) \xrightarrow{\cong} U_\alpha \times G$$

such that

- (2.1) Denoting by  $p_1 : U_\alpha \times G \rightarrow U_\alpha$  the projection, then  $\pi_P = p_1 \circ \phi_\alpha$
- (2.2) The  $G$ -action preserves each  $\pi_P^{-1}(U_\alpha)$ . Denoting the right  $G$ -action

on  $U_\alpha \times G$  by

$$(U_\alpha \times G) \times G \rightarrow U_\alpha \times G, ((x, h), g) \mapsto (x, h) \cdot g = (x, hg)$$

Then  $\phi_\alpha$  is  $G$ -equivalent, i.e.  $\forall \xi \in \pi_P^{-1}(U_\alpha), \forall g \in G, \phi_\alpha(\xi g) = \phi_\alpha(\xi)g$ . Note that the  $G$ -action is free.

We often write  $P|_U = \pi_P^{-1}(U)$  for open subset  $U \subseteq M$  and  $P_x = \pi_P^{-1}(x)$  for  $x \in M$ ,  $P_x$  is called the fiber of  $P$  at  $x$ .

Recall that  $e \in G$  is the unit, define a section  $p_\alpha \in \Gamma(P|_{U_\alpha})$  on  $U_\alpha$ :  $\phi_\alpha(p_\alpha(x)) = (x, e)$ , which is equivalent to  $p_\alpha(x) = \phi_\alpha^{-1}(x, e)$ . Define  $g_{\alpha\beta} : U_\alpha \cap U_\beta \rightarrow G$  by  $p_\alpha(x)g_{\alpha\beta}(x) = p_\beta(x)$ ,  $\{g_{\alpha\beta}\}_{\alpha\beta}$  is called the transition map of  $\pi_P : P \rightarrow M$ . Note that  $\forall x \in U_\alpha \cap U_\beta \cap U_\gamma$ , we have  $g_{\alpha\beta}(x)g_{\beta\gamma}(x) = g_{\alpha\gamma}(x)$ . Conversely, given open covering  $\{U_\alpha\}_{\alpha \in A}$  of  $M$  and transition maps, we can recover principle  $G$ -bundle  $\pi_P : P \rightarrow M$ .

As before, for  $g \in G$ , we can define  $R_g : P \rightarrow P$  the right translation and the fundamental vector field  $X^\sharp$  generated by  $X \in \mathfrak{g}$ .

**Definition 1.12.** Let  $\pi_P : P \rightarrow M$  be a principle  $G$ -bundle,  $\rho : G \rightarrow GL(V)$  representation of  $G$ . Define the right  $G$ -action on  $P \times V$  by

$$(P \times V) \times G \rightarrow P \times V, ((\xi, v), g) \mapsto (\xi g, \rho(g)^{-1}v)$$

$P \times V = (P \times V)/G$  is called the *associated vector bundle to  $P$* .

Set  $\xi \times_\rho v$  the equivalence class of  $(\xi, v) \in P \times V$ . Set  $E = P \times_\rho V$ ,  $\pi_E : E \rightarrow M$ ,  $\xi \times_\rho v \mapsto \pi_P(\xi)$ . Then  $\pi_E : E \rightarrow M$  is a vector bundle.

The local trivialization of  $E$  are induced from those of  $P$ :

$$\phi_\alpha^E : E|_{U_\alpha} \xrightarrow{\cong} U_\alpha \times V, p_\alpha(x) \times_\rho v \mapsto (x, v)$$

For  $x \in U_\alpha \cap U_\beta$  and  $v_\beta \in V$ ,  $p_\beta(x) \times_\rho v_\beta = p_\alpha g_{\alpha\beta}(x) \times_\rho v_\beta = p_\alpha(x) \times_\rho \rho(g_{\alpha\beta}(x)) v_\beta$ . The transition functions of  $E$  are given by  $\{\rho(g_{\alpha\beta}) : U_\alpha \cap U_\beta \rightarrow GL(V)\}$ .

We will explain some relations between  $P$  and  $E$ .

- First note that  $\forall \xi \in P$ , we have  $\xi : V \xrightarrow{\cong} E_{\pi_P(\xi)}, v \mapsto \xi \times_\rho v$  is an isomorphism. For  $\xi' \in P$  with  $\xi' = \xi g$  for  $g \in G$ , we have  $\xi^{-1} \left( \xi' \times_\rho v' \right) = \xi^{-1} \left( \xi \times_\rho \rho(g) v' \right) = \rho(g) v'$  for  $v' \in V$ .
- $\pi_P^* E$  is a trivial bundle. Indeed,

$$P \times V \xrightarrow[\substack{(\xi, \xi^{-1}(e)) \mapsto (\xi, e)}]{\substack{(\xi, v) \mapsto (\xi, \xi \times_\rho v)}} \pi_P^* E = \{(\xi, e) \in P \times E \mid \pi_P(\xi) = \pi_E(e)\} \text{ is isomorphism.}$$

- Next, for  $s \in \Omega^q(E) = \Gamma(\Lambda^q T^* M \otimes E)$ , define  $\pi_P^* s \in \Omega^q(P; V)$  as follows ( $V$ -

valued  $q$ -form on  $P$ )

- For  $q = 0$ ,  $(\pi_P^* s)(\xi) = \xi^{-1}(s(\pi_P(\xi)))$
- For  $q > 1$ ,  $\forall \alpha \in \Omega^q(M)$ ,  $\forall s \in \Omega^0(E) = \Gamma(E)$ ,

$$\pi_P^*(\alpha \otimes s) = \pi_P^* \alpha \otimes \pi_P^* s$$

The left one is pullback and the right one is define above. In other words,  
 $\forall \xi \in P$ ,  $\forall v_1, \dots, v_q \in T_\xi P$ ,

$$(\pi_P^* s)_\xi(v_1, \dots, v_q) = \xi^{-1}(s_{\pi_P(\xi)}(\pi_{P*}(v_1), \dots, \pi_{P*}(v_q)))$$

Notation: denote  $\Omega_B^q(P; V)$  to be the elements  $\tilde{s}$  in  $\Omega^q(P; V)$  satisfying:

- $\forall X \in \mathfrak{g}$ ,  $i(X^\sharp)\tilde{s} = 0$ .
- $\forall g \in G$ ,  $R_g^* \tilde{s} = \rho(g)^{-1} \tilde{s}$ .

called the *space of basic  $q$ -forms*. Note that  $\Omega_B^q(P; V)$  depends on representation  $\rho$ .

**Proposition 1.5. (Important to study the relations between  $P$  and  $E$ )**

- (1)  $\pi_P^*(\Omega^q(E)) \subseteq \Omega_B^q(P; V)$  and  $\pi_P^* : \Omega^q(E) \xrightarrow{\cong} \Omega_B^q(P; V)$ .  $E$ -valued  $q$ -forms on  $M$  are identified with basic  $q$ -forms on  $P$ .
- (2) Recall the local trivialization  $\phi_\alpha^E : E|_{U_\alpha} \xrightarrow{\cong} U_\alpha \times V$ . For  $s \in \Omega^q(E)$ , suppose that  $s|_{U_\alpha}$  corresponds to  $s_\alpha \in \Omega^q(U_\alpha; V)$ . Then  $s_\alpha = p_\alpha^*(\pi_P^* s)$ . So we regard  $s \in \Omega^q(E)$  as a basic form, and then pullback by  $p_\alpha$  is  $s_\alpha$ .

*Proof.* We show  $\pi_P^*(\Omega^q(E)) \subseteq \Omega_B^q(P; V)$ . Take  $\forall s \in \Omega^q(E)$ , □

**Example 1.4.** Let  $\pi_E : E \rightarrow M$  be a vector bundle with rank  $r$ . For  $x \in M$ ,

### 1.3 Connections on Principle Bundles

In this subsection we study properties of connection on principle bundle and its relation between connection on associated vector bundle.

**Definition 1.13.** Let  $\pi_P : P \rightarrow M$  be principle  $G$ -bundle.

- (1) A distribution  $\{H_\xi \subseteq T_\xi P\}_{\xi \in P}$  is a *connection* on  $P$  if
  - (1-1)  $\forall \xi \in P$ ,  $T_\xi P = \ker(\pi_P)_* \oplus H_\xi$ .
  - (1-2)  $\{H_\xi \subseteq T_\xi P\}_{\xi \in P}$  is  $G$ -invariant, i.e.  $\forall \xi \in P$ ,  $\forall g \in G$ ,  $(R_g)_* H_\xi = H_{\xi g}$ . $H_\xi$ ,  $\ker(\pi_P)_*$  are called *horizontal/vertical subspaces*.
- (2) A  $\mathfrak{g}$ -valued 1-form  $\theta \in \Omega^1(P; \mathfrak{g})$  on  $P$  is a *connection form* if
  - (2-1)  $\forall X \in \mathfrak{g}$ ,  $\theta(X^\sharp) = X$ .
  - (2-2)  $\forall g \in G$ ,  $R_g^* \theta = \text{Ad}(g^{-1})\theta$ .

These 2 notions are the same in the following sense:

**Theorem 1.2.** Let  $\pi_P : P \rightarrow M$  be principle  $G$ -bundle.

- (1) If  $\theta \in \Omega^1(P; \mathfrak{g})$  is a connection form, a distribution  $\{\ker \theta_\xi\}_{\xi \in P} = \{v \in T_\xi P \mid \theta_\xi(v) = 0\}_{\xi \in P}$  is a connection on  $P$ .  
 (2)  $\{\text{connection form}\} \rightarrow \{\text{connection on } P\}$ ,  $\theta \mapsto \{\ker \theta_\xi\}_{\xi \in P}$  is bijective.

*Proof.* (1) We check that  $\{\ker \theta_\xi\}_{\xi \in P}$  satisfies (1-1), (1-2):

(1-1) Note that  $\ker(\pi_P)_{*\xi} = \{X_\xi^\# \in T_\xi P \mid X \in \mathfrak{g}\}$ , then for  $\forall v \in T_\xi P$ , we have  $\theta(v) \in \mathfrak{g}$  and  $v = \theta(v)_\xi^\# + (v - \theta(v)_\xi^\#)$ , which implies that  $T_\xi P = \ker(\pi_P)_{*\xi} \oplus \ker \theta_\xi$  ( $\ker(\pi_P)_{*\xi} \cap \ker \theta_\xi = \{0\}$  is obvious).

(1-2) Take  $\forall v \in \ker \theta_\xi$ . By (2-2),  $\forall g \in G$ , we have  $(R_g^* \theta)_\xi = \text{Ad}(g^{-1})\theta_\xi$ , the left hand side is  $\theta_{\xi g}((R_g)_*\xi(\cdot))$ , so we have  $(R_g)_*\xi(v) \in \ker \theta_{\xi g}$ , hence  $(R_g)_*\xi(\ker \theta_\xi) \subseteq \ker \theta_{\xi g}$ . Replacing  $(g, \xi)$  with  $(g^{-1}, \xi g)$ , we have  $(R_{g^{-1}})_*\xi g(\ker \theta_{\xi g}) \subseteq \ker \theta_\xi$ . So  $(R_g)_*\xi(\ker \theta_\xi) = \ker \theta_{\xi g}$ ,  $\{\ker \theta_\xi\}_{\xi \in P}$  is a connection on  $P$ .

(2) **Injectivity** Let  $\theta, \theta'$  be connection forms with  $\ker \theta_\xi = \ker \theta'_\xi \forall \xi \in P$ . We show that  $\forall v \in T_\xi P$ ,  $\theta_\xi(v) = \theta'_\xi(v)$ . By (1),  $v$  is described as  $v = X_\xi^\# + w$  for  $X_\xi^\# \in \ker(\pi_P)_{*\xi}$  and  $w \in \ker \theta_\xi = \ker \theta'_\xi$ . So  $\theta_\xi(v) = \theta_\xi(X_\xi^\#) = X = \theta'_\xi(v)$ .

**Surjectivity** Take  $\forall \{H_\xi\}_{\xi \in P}$  a connection on  $P$ . By (1-1), we can define  $\theta \in \Omega^1(P; \mathfrak{g})$  by

$$\theta_\xi(v) = \begin{cases} 0 & (v \in H_\xi) \\ X & (v = X_\xi^\# \text{ for } X \in \mathfrak{g}) \end{cases}$$

By definition,  $\ker \theta_\xi = H_\xi$ , we check (2-1), (2-2).

(2-1) Holds by definition of  $\theta_\xi$ .

(2-2)  $\forall \xi \in P$ ,  $\forall g \in G$ , we show that  $\theta_{\xi g}((R_g)_*\xi(\cdot)) = \text{Ad}(g^{-1})\theta_\xi$  on  $T_\xi P$ . Recall that  $T_\xi P = \ker(\pi_P)_{*\xi} \oplus H_\xi$ , if  $v \in H_\xi$ , the equality holds by definition and (1-2); for  $\forall X \in \mathfrak{g}$ ,

$$(R_g)_*\xi(X_\xi^\#) = (R_g)_*\xi \frac{d}{dt} \xi \exp_G(tX) \big|_{t=0} = \frac{d}{dt} \xi g \cdot g^{-1} \exp_G(tX) g \big|_{t=0} = (\text{Ad}(g^{-1})X)_\xi^\#$$

So  $\theta_{\xi g}((R_g)_*\xi(X_\xi^\#)) = \text{Ad}(g^{-1})X = \text{Ad}(g^{-1})\theta_\xi(X_\xi^\#)$ , hence the equality holds. So we have  $\theta_{\xi g}((R_g)_*\xi(\cdot)) = \text{Ad}(g^{-1})\theta_\xi$  on  $T_\xi P$ .  $\square$

The next proposition says that a connection form  $\theta$  on  $P$  induces a connection  $\nabla^E$  of the associated vector bundle  $E$ . The relation between  $\theta$  and local connection form of  $\nabla^E$  is also given.

**Proposition 1.6.** Let  $\pi_P : P \rightarrow M$  be a principle bundle,  $\rho : G \rightarrow GL(V)$  a representation of  $G$  with differential representation  $\rho_* : \mathfrak{g} \rightarrow \text{End}(V)$ . Denote

by  $\theta \in \Omega^1(P; \mathfrak{g})$  a connection form. Set  $E = P \times_{\rho} V$  its associated vector bundle.

Then,

(1)  $(d + \rho_*(\theta) \wedge) \Omega_B^q(P; V) \subseteq \Omega_B^{q+1}(P; V)$ . Here

- $d$ : standard exterior derivative.
- $\rho_*(\theta) \in \Omega^1(P; \text{End}(V))$  acts on  $\Omega_B^q(P; V)$  by wedging on differential form parts and composing  $\text{End}(V)$ ,  $V$ -parts.

(2) Recall that  $\pi_P^* : \Omega^q(E) \xrightarrow{\cong} \Omega_B^q(P; V)$ . Then we can define  $\nabla^E : \Omega^0(E) \rightarrow \Omega^1(E)$  by  $(\pi_P^*)^{-1} \circ (d + \rho_*(\theta) \wedge) \circ \pi_P^*$ .

(3) Recall that a local section  $p_\alpha \in \Gamma(P|_{U_\alpha})$  induces a local trivialization  $\phi_\alpha^E : E|_{U_\alpha} \xrightarrow{\cong} U_\alpha \times V$ . Then

$$\begin{array}{ccc} \Omega^0(E|_{U_\alpha}) & \xrightarrow{\nabla^E|_{U_\alpha}} & \Omega^1(E|_{U_\alpha}) \\ \parallel & \curvearrowright & \parallel \\ \Omega^0(U_\alpha; V) & \xrightarrow{d + \rho_\alpha^*(\theta) \wedge} & \Omega^1(U_\alpha; V) \end{array}$$

(4) Recall that a connection  $\nabla^E$  induces the exterior derivative  $d^{\nabla^E} : \Omega^q(E) \rightarrow \Omega^{q+1}(E)$ . Then

$$\begin{array}{ccc} \Omega^q(E) & \xrightarrow{d^{\nabla^E}} & \Omega^{q+1}(E) \\ \pi_P^* \downarrow \cong & \curvearrowright & \pi_P^* \downarrow \cong \\ \Omega_B^q(P; V) & \xrightarrow{d + \rho_*(\theta) \wedge} & \Omega_B^{q+1}(P; V) \end{array}$$

**Remark 1.4.** In [Kobayashi-Nomizu, *Foundation of differential geometry* Vol 1, chapter 2, section 5], for any principle  $G$ -bundle with a connection form  $\theta \in \Omega^1(P; \mathfrak{g})$ ,  $\forall V$  vector space, the *exterior covariant derivative*  $D : \Omega^q(P; V) \rightarrow \Omega^{q+1}(P; V)$  is defined by  $(D\tilde{s})(v_0, \dots, v_q) = (d\tilde{s})(hv_0, \dots, hv_q)$  for  $v_i \in TP$ , where  $h : TP \rightarrow \ker \theta$  is the projection. If in addition, given a representation  $\rho : G \rightarrow GL(V)$  and  $\tilde{s} \in \Omega_B^q(P; V)$ , we have  $D\tilde{s} = (d + \rho_*(\theta) \wedge)(\tilde{s})$ .

*Proof.* (1) Take  $\forall \tilde{s} \in \Omega_B^q(P; V)$ , recall that  $\begin{cases} \forall X \in \mathfrak{g}, i(X^\#)\tilde{s} = 0. \\ \forall g \in G, R_g^*\tilde{s} = \rho(g)^{-1}\tilde{s}. \end{cases}$ . We show that  $(d + \rho_*(\theta) \wedge)\tilde{s}$  also satisfies the same property.

- $\forall X \in \mathfrak{g}$ , we have  $\mathcal{L}_{X^\#}\tilde{s} = \frac{d}{dt} R_{\exp_G(tX)}^* \tilde{s} |_{t=0} = \frac{d}{dt} \rho(\exp_G(tX))^{-1} \tilde{s} |_{t=0} = -\rho_*(X)\tilde{s}$ . Since  $\mathcal{L}_{X^\#}\tilde{s} = i(X^\#)d\tilde{s} + d(i(X^\#)\tilde{s})$  and  $i(X^\#)\tilde{s} = 0$ , we have  $i(X^\#)d\tilde{s} = -\rho_*(X)\tilde{s}$ . Hence  $i(X^\#)((d + \rho_*(\theta) \wedge)(\tilde{s})) = i(X^\#)d\tilde{s} + \rho_*(\theta(X^\#))\tilde{s} - \rho_*(\theta) \wedge i(X^\#)\tilde{s} = 0$ .
- For  $\forall g \in G$ ,  $R_g^*((d + \rho_*(\theta) \wedge)(\tilde{s})) = dR_g^*\tilde{s} + \rho_*(R_g^*\theta) \wedge R_g^*\tilde{s} = d(\rho(g)^{-1}\tilde{s}) +$

$\rho_*(Ad(g^{-1})\theta) \wedge \rho(g)^{-1}\tilde{s}$ . Since  $\rho(g)^{-1}$  acts only on  $V$ -part,  $d(\rho(g)^{-1}\tilde{s}) = \rho(g)^{-1}d\tilde{s}$ . Note that  $\forall X \in \mathfrak{g}$ ,  $\frac{d}{dt}\rho(g^{-1}\exp_G(tX)g)\rho(g)^{-1}|_{t=0} = \frac{d}{dt}\rho(g^{-1}\exp_G(tX))|_{t=0}$  and  $g^{-1}\exp_G(tX)g = \exp_G(tAd(g^{-1})X)$ , we have  $\rho_*(Ad(g^{-1})X)\rho(g)^{-1} = \rho(g)^{-1}\rho_*(X)$ . This implies that  $\rho_*(Ad(g^{-1})\theta) \wedge \rho(g)^{-1}\tilde{s} = \rho(g)^{-1}(\rho_*(\theta) \wedge \tilde{s})$ . Then we obtain  $R_g^*((d + \rho_*(\theta) \wedge)(\tilde{s})) = \rho(g)^{-1}((d + \rho_*(\theta) \wedge)(\tilde{s}))$ , so  $(d + \rho_*(\theta) \wedge)(\tilde{s}) \in \Omega_B^{q+1}(P; V)$ .

(2)  $\nabla^E = (\pi_P^*)^{-1} \circ (d + \rho_*(\theta) \wedge) \circ \pi_P^*$ , we check the Leibniz rule, i.e. for  $\forall f \in C^\infty(M)$ ,  $\forall s \in \Gamma(E)$ , we show  $\nabla^E(fs) = df \otimes s + f\nabla^E s$ . This is left as an exercise.

(3) Since for  $s \in \Omega^q(E)$ ,  $s|_{U_\alpha}$  corresponds to  $p_\alpha^*(\pi_P^*s)$ . We compute  $p_\alpha^*\pi_P^*(\nabla^E s) = p_\alpha^*((d + \rho_*(\theta) \wedge)\pi_P^*s) = p_\alpha^*d(\pi_P^*s) + \rho_*(p_\alpha^*\theta) \wedge p_\alpha^*\pi_P^*s = (d + \rho_*(p_\alpha^*\theta) \wedge)(p_\alpha^*\pi_P^*s)$ .

(4) Since  $d^{\nabla^E}$  is given by  $d^{\nabla^E}(s \otimes \alpha) = \nabla^E s \wedge \alpha + s \otimes d\alpha$  for  $s \in \Gamma(E)$ ,  $\alpha \in \Omega^q(M)$ , we have  $\pi_P^*(d^{\nabla^E}(s \otimes \alpha)) = \pi_P^*(\nabla^E s \wedge \alpha + s \otimes d\alpha) = (d + \rho_*(\theta) \wedge)\pi_P^*s \wedge \pi_P^*\alpha + \pi_P^*s \otimes \pi_P^*d\alpha = d(\pi_P^*s \otimes \pi_P^*\alpha) + \rho_*(\theta) \wedge (\pi_P^*s \otimes \pi_P^*\alpha) = (d + \rho_*(\theta) \wedge)(\pi_P^*(s \otimes \alpha))$ .

□

**Exercise 1.2.** Prove that  $\nabla^E$  defined above is a connection.

**Example 1.5.** Given a vector bundle  $\pi_E : E \rightarrow M$ , let  $\pi_P : P \rightarrow M$  be the frame bundle. Consider the trivial representation  $id : GL(r; \mathbb{K}) \rightarrow GL(r; \mathbb{K})$ . Then