


Principal Component Analysis

5 questions



1
point



1.
Consider the following 2D dataset:



Which of the following figures correspond to possible values that PCA may return for $u^{(1)}$ (the first eigenvector / first principal component)? Check all that apply (you may have to check more than one figure).







1
point

2.

Which of the following is a reasonable way to select the number of principal components k ?

(Recall that n is the dimensionality of the input data and m is the number of input examples.)

- ☐ Choose the value of k that minimizes the approximation error $\frac{1}{m} \sum_{i=1}^m ||x^{(i)} - x_{\text{approx}}^{(i)}||^2$.
 - ☐ Choose k to be the smallest value so that at least 1% of the variance is retained.
 - ☐ Choose k to be the smallest value so that at least 99% of the variance is retained.
 - ☐ Choose k to be 99% of n (i.e., $k = 0.99 * n$, rounded to the nearest integer).
-

1
point

3.

Suppose someone tells you that they ran PCA in such a way that "95% of the variance was retained." What is an equivalent statement to this?

- ☐ $\frac{\frac{1}{m} \sum_{i=1}^m ||x^{(i)}||^2}{\frac{1}{m} \sum_{i=1}^m ||x^{(i)} - x_{\text{approx}}^{(i)}||^2} \geq 0.05$
 - ☐ $\frac{\frac{1}{m} \sum_{i=1}^m ||x^{(i)}||^2}{\frac{1}{m} \sum_{i=1}^m ||x^{(i)} - x_{\text{approx}}^{(i)}||^2} \leq 0.05$
 - ☐ $\frac{\frac{1}{m} \sum_{i=1}^m ||x^{(i)}||^2}{\frac{1}{m} \sum_{i=1}^m ||x^{(i)} - x_{\text{approx}}^{(i)}||^2} \leq 0.95$
 - ☐ $\frac{\frac{1}{m} \sum_{i=1}^m ||x^{(i)} - x_{\text{approx}}^{(i)}||^2}{\frac{1}{m} \sum_{i=1}^m ||x^{(i)}||^2} \leq 0.05$
-

1
point

4.

Which of the following statements are true? Check all that apply.



PCA can be used only to reduce the dimensionality of data by 1 (such as 3D to 2D, or 2D to 1D).

- ☐ Given an input $x \in \mathbb{R}^n$, PCA compresses it to a lower-dimensional vector $z \in \mathbb{R}^k$.
- ☐ Feature scaling is not useful for PCA, since the eigenvector calculation (such as using Octave's `svd(Sigma)` routine) takes care of this automatically.
- ☐ If the input features are on very different scales, it is a good idea to perform feature scaling before applying PCA.

1
point

5.

Which of the following are recommended applications of PCA? Select all that apply.

- ☐ As a replacement for (or alternative to) linear regression: For most learning applications, PCA and linear regression give substantially similar results.
- ☐ Data compression: Reduce the dimension of your input data $x^{(i)}$, which will be used in a supervised learning algorithm (i.e., use PCA so that your supervised learning algorithm runs faster).
- ☐ Data visualization: To take 2D data, and find a different way of plotting it in 2D (using $k=2$).
- ☐ Data compression: Reduce the dimension of your data, so that it takes up less memory / disk space.



I understand that submitting work that isn't my own may result in permanent failure of this course or deactivation of my Coursera account. Learn more about Coursera's Honor Code

Anderson Banihirwe

Submit Quiz