DSC 210, Fall 2026 — Homework 2 Nicholas Wendt (200478549)

Question 1: Gauss elimination (20 points)

Use Gauss elimination to solve the following equations for $\mathbf{x} = \begin{bmatrix} x_1 & x_2 & x_3 \end{bmatrix}^\top$:

$$\begin{aligned} x_1 + 5x_2 + 3x_3 &= 1 \\ 2x_1 + 4x_2 - 3x_3 &= -1 \\ -2x_1 + 2x_2 + 3x_3 &= -5 \end{aligned}$$

Solution: First set up an augmented matrix

$$\left[\begin{array}{ccc|c} 1 & 5 & 3 & 1 \\ 2 & 4 & -3 & -1 \\ -2 & 2 & 3 & -5 \end{array}\right]$$

Now solve the system $Ax = b$ by row reducing the matrix.
Apply the row operations $2r_1 - r_2 \to r_2$ and $2r_1 + r_3 \to r_3$

$$\left[\begin{array}{ccc|c} 1 & 5 & 3 & 1 \\ 0 & 6 & 9 & 3 \\ 0 & 12 & 9 & -3 \end{array}\right]$$

Now use row 2's pivot column $2r_2 - r_3 \to r_3$

$$\left[\begin{array}{ccc|c} 1 & 5 & 3 & 1 \\ 0 & 6 & 9 & 3 \\ 0 & 0 & 9 & 9 \end{array}\right]$$

Now use back substitution to solve for $\mathbf{x}$ Starting with the last row we get the equation $9x_3 = 9$ so $x_3 = 1$.
Now the second row gives us the equation $6x_2 + 9x_3 = 3 \to 6x_2 + 9 = 3$. So we see $x_2 = -1$.
Finally, the first equation gives us $1x_1 + 5x_2 + 3x_3 = 1 \to x_1 - 5 + 3 = 1$. So we see $x_1 = 3$.

$$\mathbf{x} = \begin{bmatrix} x_1 & x_2 & x_3 \end{bmatrix}^\top = \begin{bmatrix} 3 & -1 & 1 \end{bmatrix}$$

Question 2: LU decomposition (20 points)

Perform LU decomposition for the matrix corresponding to the equations in **Question 1**. Using the matrices L and U, do forward and backward substitution to solve for $\mathbf{x}$. Match your solution with that of **Question 1**.

Solution:

$$A = \begin{bmatrix} 1 & 5 & 3 \\ 2 & 4 & -3 \\ -2 & 2 & 3 \end{bmatrix}$$

We can factor A into a lower triangular matrix (L) and an upper triagular matrix (U) such that $A = LU$
We row reduce A to an upper triangular matrix without using row interchanges.
We keep track of the scalers in L.

First 2 operations $r_2 - 2r_1$, $r_3 + 2r_1$. We keep track of the scalers of the pivot row in L according to $r_i - cr_j$ where c is the row i column j entry in L.

Now we have:

$$A = \begin{bmatrix} 1 & 5 & 3 \\ 0 & -6 & -9 \\ 0 & 12 & 9 \end{bmatrix} \quad , \quad L = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ -2 & _ & 1 \end{bmatrix}$$

The last row operation is $r_3 + 2r_2$. Since A in now upper triangular, we write U

$$U = \begin{bmatrix} 1 & 5 & 3 \\ 0 & -6 & -9 \\ 0 & 0 & -9 \end{bmatrix} \quad , \quad L = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ -2 & -2 & 1 \end{bmatrix}$$

Firs use forward substitution to solve $Ly = b$:

$$\begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ -2 & -2 & 1 \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \\ -5 \end{bmatrix}$$

$$\begin{aligned} y_1 &= 1 \\ 2y_1 + y_2 = -1 &\implies y_2 = -1 - 2(1) = -3 \\ -2y_1 - 2y_2 + y_3 = -5 &\implies y_3 = -5 + 2(1) + 2(-3) = -9 \end{aligned}$$

So $\mathbf{y} = \begin{bmatrix} 1 \\ -3 \\ -9 \end{bmatrix}$.

Then use backward substitution to solve $Ux = y$:

$$\begin{bmatrix} 1 & 5 & 3 \\ 0 & -6 & -9 \\ 0 & 0 & -9 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 1 \\ -3 \\ -9 \end{bmatrix}$$

$$
\begin{aligned}
-9x_3 = -9 &\implies x_3 = 1 \\
-6x_2 - 9x_3 = -3 &\implies x_2 = \frac{-3 + 9(1)}{-6} = -1 \\
x_1 + 5x_2 + 3x_3 = 1 &\implies x_1 = 1 - 5(-1) - 3(1) = 3
\end{aligned}
$$

$$
\mathbf{x} = \begin{bmatrix} 3 \\ -1 \\ 1 \end{bmatrix}
$$

This matches the solution from question 1.

Question 3: QR decomposition (20 points)

Perform QR decomposition for the matrix corresponding to the equations in **Question 1** using the Gram-Schmidt algorithm. Using the decomposition, solve for $\mathbf{x}$. Match your solution with that of **Question 1** and **Question 2**.

Solution:

Question 4: Gram-Schmidt process (20 points)

Show that the residual vector $\mathbf{a}_i^{\perp}$ is orthogonal to $\mathbf{q}_1, \mathbf{q}_2, \ldots, \mathbf{q}_{i-1}$ in the Gram-Schmidt process.

Solution:

Question 5: Rank deficient matrices (20 points)

Compute the QR decomposition of the following two matrices using the Gram-Schmidt process and properties of the matrix $\mathbf{Q}$.

(i) $\mathbf{A} = \begin{bmatrix} 1 & 1 & -1 \\ 1 & -1 & 1 \\ -1 & -1 & 1 \end{bmatrix}$

(ii) $\mathbf{B} = \begin{bmatrix} 1 & -1 & 1 \\ -1 & 1 & -1 \\ -1 & 1 & -1 \end{bmatrix}$

Note: the matrices $\mathbf{Q}$ and $\mathbf{R}$ must be square matrices.

Solution:

(i)

(ii)