

# Calc 1500, Week 5

Chain Rule

Tangent  
Planes and  
Normal Lines

## 1 Chain Rule

## 2 Tangent Planes and Normal Lines

## Chain Rule

Chain Rule  
Tangent  
Planes and  
Normal Lines

Assume  $f(x, y)$ ,  $x(s, t)$ , and  $y(s, t)$  are of class  $\mathcal{C}^1$ .

$$f(x(s, t), y(s, t))$$

is also of class  $\mathcal{C}^1$  and

$$\frac{\partial}{\partial s} f(x(s, t), y(s, t)) = \frac{\partial f}{\partial x}(x, y) \cdot \frac{\partial x}{\partial s}(s, t) + \frac{\partial f}{\partial y}(x, y) \cdot \frac{\partial y}{\partial s}(s, t),$$

$$\frac{\partial}{\partial t} f(x(s, t), y(s, t)) = \frac{\partial f}{\partial x}(x, y) \cdot \frac{\partial x}{\partial t}(s, t) + \frac{\partial f}{\partial y}(x, y) \cdot \frac{\partial y}{\partial t}(s, t).$$

$$f = f(x, y)$$

$$x = x(s, t)$$


$$y = y(s, t)$$

$$\frac{\partial f}{\partial s}(s, t) = \frac{\partial f}{\partial x}(x, y) \cdot \frac{\partial x}{\partial s}(s, t) + \frac{\partial f}{\partial y}(x, y) \cdot \frac{\partial y}{\partial s}(s, t)$$


We also write

$$\frac{\partial f}{\partial s} = \frac{\partial f}{\partial x} \frac{\partial x}{\partial s} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial s}$$

$$\text{or } f_s = f_x x_s + f_y y_s$$



Dependency diagram



$$\frac{\partial f}{\partial t}(s,t) = \frac{\partial f}{\partial x}(x,y) \cdot \frac{\partial x}{\partial t}(s,t) + \frac{\partial f}{\partial y}(x,y) \cdot \frac{\partial y}{\partial t}(s,t)$$

## Example: Chain Rule


Chain Rule  
Tangent  
Planes and  
Normal Lines

Find *Take the partial derivative wrt s* of this function

$$\frac{\partial}{\partial s} f(x(s, t), y(s, t))$$

for  $f(x, y) = x^2 + y^2$ ,  $x = st$ , and  $y = s \cos t$ .

$$\begin{aligned}\frac{\partial f}{\partial s} &= \frac{\partial f}{\partial x} \cdot \frac{\partial x}{\partial s} + \frac{\partial f}{\partial y} \cdot \frac{\partial y}{\partial s} \\ &= (2x)(t) + (2y)(\cos t) \\ &= 2xt + 2y \cos t\end{aligned}$$



$$\begin{aligned}
 & \frac{\partial}{\partial s} f(x(s,t), y(s,t)) \\
 & \text{let } F(s,t) = f(x(s,t), y(s,t)) \\
 & \begin{bmatrix} s \\ t \end{bmatrix} \mapsto \begin{bmatrix} x(s,t) \\ y(s,t) \end{bmatrix} \mapsto [f(x, y)] \\
 & x = st \\
 & y = s \cos t \\
 & f = x^2 + y^2
 \end{aligned}$$

$$\frac{\partial}{\partial s} f(x(s,t), y(s,t)) = \frac{\partial F}{\partial s}(s,t)$$

$L : \mathbb{R}^n \rightarrow \mathbb{R}^m$   $[L]$  is an  $m \times n$  matr.x

$$\begin{bmatrix} \frac{\partial F}{\partial s} & \frac{\partial F}{\partial t} \end{bmatrix} = \begin{bmatrix} \frac{\partial f}{\partial x} & \frac{\partial f}{\partial y} \end{bmatrix} \begin{bmatrix} \frac{\partial x}{\partial s} & \frac{\partial x}{\partial t} \\ \frac{\partial y}{\partial s} & \frac{\partial y}{\partial t} \end{bmatrix}$$

## Notation for Chain Rule

Chain Rule

Tangent  
Planes and  
Normal Lines

Assume  $f(x, y)$ ,  $x(s, t)$ , and  $y(s, t)$  are of class  $\mathcal{C}^1$ . Instead of writing

$$\frac{\partial}{\partial s} f(x(s, t), y(s, t)) = \frac{\partial f}{\partial x}(x, y) \cdot \frac{\partial x}{\partial s}(s, t) + \frac{\partial f}{\partial y}(x, y) \cdot \frac{\partial y}{\partial s}(s, t),$$

we may write

$$\frac{\partial F}{\partial s} = \frac{\partial f}{\partial x} \cdot \frac{\partial x}{\partial s} + \frac{\partial f}{\partial y} \cdot \frac{\partial y}{\partial s},$$

or

$$F_s = f_x x_s + f_y y_s,$$

where  $F$  is the composite function  $F(s, t) = f(x(s, t), y(s, t))$ .


## Example: Chain Rule (FRY Exercise III.2.4.5.1c)

Chain Rule  
Tangent  
Planes and  
Normal Lines

*Composite function*  
Let  $h(x, y, z) = f(u(x, y, z), v(x, y), w(x))$ . Find  $\frac{\partial h}{\partial x}$ .

(Show the dependency diagram.)

$$\frac{\partial h}{\partial x} = \frac{\partial}{\partial x} f(u(x, y, z), v(x, y), w(x))$$



$$\frac{\partial h}{\partial x} = \frac{\partial f}{\partial u} \frac{\partial u}{\partial x} + \frac{\partial f}{\partial v} \frac{\partial v}{\partial x} + \frac{\partial f}{\partial w} \frac{\partial w}{\partial x}$$

We use the ordinary (non-curly)  $d$  for the derivative of  $w$  with respect to  $x$  because  $w$  is a function of a single variable (and not a multivariable function).


## Live Poll: Chain Rule

Chain Rule  
Tangent  
Planes and  
Normal Lines

Let  $f = f(x, y)$ ,  $x = x(s, t)$ , and  $y = y(t)$ .

(a) Draw a dependency diagram for the variables.

(b) Write an expression for  $\frac{\partial}{\partial t} f(x(s, t), y(t))$ .



$$\frac{\partial f}{\partial t} = \frac{\partial f}{\partial x} \cdot \frac{\partial x}{\partial t} + \frac{\partial f}{\partial y} \cdot \frac{dy}{dt}$$

## Example: Chain Rule (Like FRY Exercise III.2.4.5.3)


Chain Rule


Tangent  
Planes and  
Normal Lines

Let  $w = f(x, t)$  where  $x$  depends on  $t$ . Suppose that at some point  $x_0$  and at some time  $t_0$ ,

- the partial derivatives  $f_x$  and  $f_t$  equal  $-2$  and  $-3$ , respectively; and
- $\frac{dx}{dt} = 2$ .

What is  $\frac{dw}{dt}$ ?





Given

- $f_x(x_0, t_0) = -2$
- $f_t(x_0, t_0) = -3$

Goal : Find  $\frac{dw}{dt}(t_0) = ?$

$$\cdot \frac{dx}{dt}(t_0) = 2$$

$$\begin{aligned}\frac{dw}{dt}(t_0) &= \frac{\partial f}{\partial x}(x_0, t_0) \frac{dx}{dt}(t_0) + \frac{\partial f}{\partial t}(x_0, t_0) \\ &= (-2)(2) + (-3) \\ &= -7\end{aligned}$$

## Example: Chain Rule (FRY Exercise III.2.4.5.21a)

Suppose that  $f(x, y)$  is twice differentiable with  $f_{xy} = f_{yx}$  and  $x = r \cos \theta$  and  $y = r \sin \theta$ . Evaluate  $f_\theta$ ,  $f_r$ , and  $f_{r\theta}$  in terms of  $r$ ,  $\theta$ ,  $f_x$  and  $f_y$ .

Observe that

$$x_\theta = -r \sin \theta, \quad y_\theta = r \cos \theta, \quad x_r = \cos \theta, \quad y_r = \sin \theta.$$

and

$$f_\theta = f_x \cdot x_\theta + f_y \cdot y_\theta = -r \sin \theta f_x + r \cos \theta f_y$$

$$f_r = f_x \cdot x_r + f_y \cdot y_r = \cos \theta f_x + \sin \theta f_y$$

Chain Rule

Tangent  
Planes and  
Normal Lines

## Example: Chain Rule (FRY Exercise III.2.4.5.21a)

$$\begin{aligned} f_{r\theta} &= (f_r)_\theta = \left( \cos \theta f_x + \sin \theta f_y \right)_\theta \\ &= -\sin \theta f_x + \cos \theta (f_x)_\theta + \cos \theta f_y + \sin \theta (f_y)_\theta \\ &= -\sin \theta f_x + \cos \theta (f_{xx} x_\theta + f_{xy} y_\theta) + \cos \theta f_y + \sin \theta (f_{yx} x_\theta + f_{yy} y_\theta) \\ &= -\sin \theta f_x + \cos \theta \left( f_{xx} (-r \sin \theta) + f_{xy} r \cos \theta \right) \\ &\quad + \cos \theta f_y + \sin \theta \left( f_{yx} (-r \sin \theta) + f_{yy} r \cos \theta \right) \\ &= -\sin \theta f_x - r \sin \theta \cos \theta f_{xx} + r \cos^2 \theta f_{xy} \\ &\quad + \cos \theta f_y - r \sin^2 \theta f_{yx} + r \sin \theta \cos \theta f_{yy} \\ &= -\sin \theta f_x - \frac{1}{2} r \sin(2\theta) f_{xx} + r (\cos^2 \theta - \sin^2 \theta) f_{xy} \\ &\quad + \cos \theta f_y + \frac{1}{2} r \sin(2\theta) f_{yy} \\ &= -\sin \theta f_x - \frac{1}{2} r \sin(2\theta) f_{xx} + r \cos(2\theta) f_{xy} + \cos \theta f_y + \frac{1}{2} r \sin(2\theta) f_{yy} \end{aligned}$$

Chain Rule  
Tangent  
Planes and  
Normal Lines

## Tangent Plane and Normal Line to Surface $G(x, y, z) = K$ at the Point $(x_0, y_0, z_0)$

Let  $\mathbf{r}(t) = \langle x(t), y(t), z(t) \rangle$  be path on  $G(x, y, z) = K$  that passes through  $(x_0, y_0, z_0)$  at  $t = 0$ . Assume  $x = x(t)$ ,  $y = y(t)$ ,  $z = z(t)$  are differentiable at  $t = 0$ .

Since  $G(\mathbf{r}(t)) = G(x(t), y(t), z(t)) = K$ , differentiating both sides wrt  $t$  gives:

$$\frac{\partial G}{\partial x} \frac{dx}{dt} + \frac{\partial G}{\partial y} \frac{dy}{dt} + \frac{\partial G}{\partial z} \frac{dz}{dt} = 0.$$

At  $t = 0$ , we get

$$\frac{\partial G}{\partial x}(x_0, y_0, z_0) \frac{dx}{dt}(0) + \frac{\partial G}{\partial y}(x_0, y_0, z_0) \frac{dy}{dt}(0) + \frac{\partial G}{\partial z}(x_0, y_0, z_0) \frac{dz}{dt}(0) = 0.$$

## Tangent Plane and Normal Line to Surface $G(x, y, z) = K$ at the Point $(x_0, y_0, z_0)$

Written as a dot product:

$$\left\langle \frac{\partial G}{\partial x}(x_0, y_0, z_0), \frac{\partial G}{\partial y}(x_0, y_0, z_0), \frac{\partial G}{\partial z}(x_0, y_0, z_0) \right\rangle \cdot \left\langle \frac{dx}{dt}(0), \frac{dy}{dt}(0), \frac{dz}{dt}(0) \right\rangle = 0.$$

The left-hand side is such an important vector that we give it a special name.

The **gradient** of a function  $f : \mathbb{R}^3 \rightarrow \mathbb{R}$  at whose partial derivatives exist at a point  $\mathbf{a} \in \mathbb{R}^3$  is the vector

$$\nabla f(\mathbf{a}) = \left\langle \frac{\partial f}{\partial x}(\mathbf{a}), \frac{\partial f}{\partial y}(\mathbf{a}), \frac{\partial f}{\partial z}(\mathbf{a}) \right\rangle.$$

Chain Rule

Tangent  
Planes and  
Normal Lines

## Tangent Plane and Normal Line to Surface $G(x, y, z) = K$ at the Point $(x_0, y_0, z_0)$

Chain Rule

Tangent  
Planes and  
Normal Lines

Thus

$$\left\langle \frac{\partial G}{\partial x}(x_0, y_0, z_0), \frac{\partial G}{\partial y}(x_0, y_0, z_0), \frac{\partial G}{\partial z}(x_0, y_0, z_0) \right\rangle \cdot \left\langle \frac{dx}{dt}(0), \frac{dy}{dt}(0), \frac{dz}{dt}(0) \right\rangle = 0.$$

can be expressed using the gradient of  $G$  at the point  $(x_0, y_0, z_0)$ :

$$\nabla G(x_0, y_0, z_0) \cdot \mathbf{r}'(t) = 0.$$

## Tangent Plane and Normal Line to Surface $G(x, y, z) = K$ at the Point $(x_0, y_0, z_0)$

Chain Rule

Tangent  
Planes and  
Normal Lines

$$\nabla G(x_0, y_0, z_0) \cdot \mathbf{r}'(t) = 0.$$

Since our choice of path along the surface  $G(x, y, z) = K$  passing through  $(x_0, y_0, z_0)$  was arbitrary, the gradient  $\nabla G(x_0, y_0, z_0)$  is perpendicular to every path through  $(x_0, y_0, z_0)$ .

Thus, *the gradient vector  $\nabla G(x_0, y_0, z_0)$  is also a normal vector to a tangent plane to the surface  $G(x, y, z) = K$  at the point  $(x_0, y_0, z_0)$ .*

## Tangent Plane and Normal Line to Surface $G(x, y, z) = K$ at the Point $(x_0, y_0, z_0)$

Let  $K$  be a constant and  $(x_0, y_0, z_0)$  be a point on the surface  $G(x, y, z) = K$ . Assume that the gradient

Chain Rule  
Tangent  
Planes and  
Normal Lines

$$\nabla G(x_0, y_0, z_0) = \left\langle \frac{\partial G}{\partial x}(x_0, y_0, z_0), \frac{\partial G}{\partial y}(x_0, y_0, z_0), \frac{\partial G}{\partial z}(x_0, y_0, z_0) \right\rangle$$

of  $G$  at  $(x_0, y_0, z_0)$  is nonzero. Then,

- 1** The equation of the tangent plane to the surface  $G(x, y, z) = K$  at  $(x_0, y_0, z_0)$  is

$$\nabla G(x_0, y_0, z_0) \cdot \langle x - x_0, y - y_0, z - z_0 \rangle = 0.$$

- 2** The vector equation of the normal line to the surface  $G(x, y, z) = K$  at  $(x_0, y_0, z_0)$  is

$$\langle x, y, z \rangle = \langle x_0, y_0, z_0 \rangle + t \nabla G(x_0, y_0, z_0), \quad t \in \mathbb{R}.$$

## Example: Equation of Tangent Plane (Exercise III.2.5.3.8)

Chain Rule

Tangent  
Planes and  
Normal Lines

Find the equation of the tangent plane and normal line (in vector, parametric, and symmetric forms) to the surface given by

$$xyz^2 + y^2z^3 = 3 + x^2$$

at the point  $(-1, 1, 2)$ .

$\cancel{xyz^2 + y^2z^3 - x^2 = 3}$   
let  $G(x, y, z) = xyz^2 + y^2z^3 - x^2$ . The surface  
has the form  $\underline{G(x, y, z) = K}$ , where  $K=3$

eqn of tangent plane:  $\nabla G(-1, 1, 2) \cdot \langle x - (-1), y - 1, z - 2 \rangle = 0$

$$\nabla G(x_0, y_0, z_0) \cdot \langle x - x_0, y - y_0, z - z_0 \rangle = 0$$

$$\nabla G(x, y, z) = \left\langle \frac{\partial G}{\partial x}, \frac{\partial G}{\partial y}, \frac{\partial G}{\partial z} \right\rangle$$

$$= \left\langle yz^2 - 2x, xz^2 + 2yz^3, 2xyz + 3y^2z^2 \right\rangle$$

$$G(x, y, z) = xyz^2 + y^2z^3 - x^2$$

So  $\nabla G(-1, 1, 2) = \left\langle (1)(2)^2 - 2(-1), (-1)(2)^2 + 2(1)(2)^3, 2(-1)(1)(2), + 3(1)^2(2)^2 \right\rangle$

$$= \langle 6, 12, 8 \rangle$$

*{Eqn of tangent plane}*

$$\nabla G(-1, 1, 2) \cdot \langle x - (-1), y - 1, z - 2 \rangle = 0$$

$$\langle 6, 12, 8 \rangle \cdot \langle x+1, y-1, z-2 \rangle = 0$$

$$6x + 6 + 12y - 12 + 8z - 16 = 0$$

$$6x + 12y + 8z = 22$$

$$3x + 6y + 4z = 11$$

← Equation of  
Tangent Plane  
to the  
given surface  
at  $(-1, 1, 2)$ .

---

**Vector equation of normal line at  $(-1, 1, 2)$**

$$\langle x, y, z \rangle = \langle -1, 1, 2 \rangle + t \langle 6, 12, 8 \rangle, t \in \mathbb{R}$$

↑  
direction vector  
is  $\sqrt{6}(-1, 1, 2)$

## Parametric Eqn of Normal Line

$$x = -1 + 6t$$

$$y = 1 + 12t \quad , t \in \mathbb{R}$$

$$z = 2 + 8t$$

↖ Isolate for  $t$

## Symmetric Eqn of Normal Line

$$\frac{x+1}{6} = \frac{y-1}{12} = \frac{z-2}{8}$$

Remark:  $\nabla G(-1, 1, 2) = \langle 6, 12, 8 \rangle$

Only nonzero multiple of  $\nabla G(-1, 1, 2)$  is a "good" normal vector. For example,

$$\vec{n} = \frac{1}{\sqrt{6^2 + 12^2 + 8^2}} \langle 6, 12, 8 \rangle \quad \text{is also a normal vector.}$$