

# RELATIVE-MOTION ANALYSIS OF TWO PARTICLES USING TRANSLATING AXES

## Today's Objectives:

Students will be able to:

1. Understand translating frames of reference.
2. Use translating frames of reference to analyze relative motion.



## In-Class Activities:

- Check Homework,
- Reading Quiz
- Applications
- Relative Position, Velocity and Acceleration
- Vector & Graphical Methods
- Concept Quiz
- Group Problem Solving
- Attention Quiz

## READING QUIZ

1. The velocity of B relative to A is defined as

A)  $\mathbf{v}_B - \mathbf{v}_A$ .

B)  $\mathbf{v}_A - \mathbf{v}_B$ .

C)  $\mathbf{v}_B + \mathbf{v}_A$ .

D)  $\mathbf{v}_A + \mathbf{v}_B$ .

2. Since two-dimensional vector addition forms a triangle, there can be at most \_\_\_\_\_ unknowns (either magnitudes and/or directions of the vectors).


A) one

B) two

C) three

D) four

# APPLICATIONS




When you try to hit a moving object, the position, velocity, and acceleration of the object all have to be accounted for by your mind.

You are smarter than you thought!

Here, the boy on the ground is at  $d = 10$  ft when the girl in the window throws the ball to him.

If the boy on the ground is running at a constant speed of 4 ft/s, how fast should the ball be thrown?

## APPLICATIONS (continued)




When fighter jets take off or land on an aircraft carrier, the velocity of the carrier becomes an issue.

If the aircraft carrier is underway with a forward velocity of 50 km/hr and plane A takes off at a horizontal air speed of 200 km/hr (measured by someone on the water), how do we find the velocity of the plane relative to the carrier?

How would you find the same thing for airplane B?

How does the wind impact this sort of situation?

## RELATIVE POSITION (Section 12.10)



The **absolute positions** of two particles  $A$  and  $B$  with respect to the fixed  $x$ ,  $y$ ,  $z$ -reference frame are given by  $\mathbf{r}_A$  and  $\mathbf{r}_B$ . The **position of  $B$  relative to  $A$**  is represented by


$$\mathbf{r}_{B/A} = \mathbf{r}_B - \mathbf{r}_A$$

Therefore, if  $\mathbf{r}_B = (10 \mathbf{i} + 2 \mathbf{j}) \text{ m}$

and  $\mathbf{r}_A = (4 \mathbf{i} + 5 \mathbf{j}) \text{ m}$ ,

then  $\mathbf{r}_{B/A} = \mathbf{r}_B - \mathbf{r}_A = (6 \mathbf{i} - 3 \mathbf{j}) \text{ m}$ .

# RELATIVE VELOCITY



To determine the **relative velocity** of B with respect to A, the time derivative of the relative position equation is taken.

$$\mathbf{v}_{B/A} = \mathbf{v}_B - \mathbf{v}_A$$


or

$$\mathbf{v}_B = \mathbf{v}_A + \mathbf{v}_{B/A}$$

In these equations,  $\mathbf{v}_B$  and  $\mathbf{v}_A$  are called **absolute velocities** and  $\mathbf{v}_{B/A}$  is the **relative velocity** of B with respect to A.

Note that  $\mathbf{v}_{B/A} = -\mathbf{v}_{A/B}$ .

# RELATIVE ACCELERATION



The time derivative of the relative velocity equation yields a similar vector relationship between the **absolute** and **relative accelerations** of particles A and B.

These derivatives yield:  $\mathbf{a}_{B/A} = \mathbf{a}_B - \mathbf{a}_A$

or

$$\mathbf{a}_B = \mathbf{a}_A + \mathbf{a}_{B/A}$$

# SOLVING PROBLEMS

Since the relative motion equations are **vector equations**, problems involving them may be solved in one of two ways.


For instance, the velocity vectors in  $\mathbf{v}_B = \mathbf{v}_A + \mathbf{v}_{B/A}$  could be written as two dimensional (2-D) **Cartesian vectors** and the resulting 2-D scalar component equations solved for up to two unknowns.

Alternatively, vector problems can be solved “**graphically**” by use of trigonometry. This approach usually makes use of the **law of sines** or the **law of cosines**.

Could a CAD system be used to solve these types of problems?



# LAWS OF SINES AND COSINES




Since vector addition or subtraction forms a triangle, **sine and cosine laws** can be applied to solve for relative or absolute velocities and accelerations. As a review, their formulations are provided below.

$$\text{Law of Sines: } \frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$\begin{aligned}\text{Law of Cosines: } a^2 &= b^2 + c^2 - 2bc \cos A \\ b^2 &= a^2 + c^2 - 2ac \cos B \\ c^2 &= a^2 + b^2 - 2ab \cos C\end{aligned}$$

## EXAMPLE



**Given:** Two aircraft as shown.

$$v_A = 650 \text{ km/h}$$

$$v_B = 800 \text{ km/h}$$

**Find:**  $v_{B/A}$

**Plan:**

- 1) Vector Method: Write vectors  $v_A$  and  $v_B$  in Cartesian form, then determine  $v_B - v_A$
- 2) Graphical Method: Draw vectors  $v_A$  and  $v_B$  from a common point. Apply the laws of sines and cosines to determine  $v_{B/A}$ .

## EXAMPLE (continued)

### Solution:

#### 1) Vector Method



$$\mathbf{v}_A = (650 \mathbf{i}) \text{ km/h}$$

$$\begin{aligned}\mathbf{v}_B &= -800 \cos 60 \mathbf{i} - 800 \sin 60 \mathbf{j} \\ &= (-400 \mathbf{i} - 692.8 \mathbf{j}) \text{ km/h}\end{aligned}$$

$$\mathbf{v}_{B/A} = \mathbf{v}_B - \mathbf{v}_A = (-1050 \mathbf{i} - 692.8 \mathbf{j}) \text{ km/h}$$

$$v_{B/A} = \sqrt{(-1050)^2 + (-692.8)^2} = \underline{1258 \text{ km/h}}$$



$$\theta = \tan^{-1}\left(\frac{692.8}{1050}\right) = \underline{33.4^\circ}$$



## EXAMPLE (continued)

### 2) Graphical Method:

Note that the vector that measures the tip of B relative to A is  $\mathbf{v}_{B/A}$ .



### Law of Cosines:

$$(v_{B/A})^2 = (800)^2 + (650)^2 - (800)(650) \cos 120^\circ$$
$$v_{B/A} = \underline{1258 \text{ km/h}}$$

### Law of Sines:

$$\frac{v_{B/A}}{\sin(120^\circ)} = \frac{v_A}{\sin \theta} \quad \text{or} \quad \theta = \underline{33.4^\circ}$$

## CONCEPT QUIZ


1. Two particles, A and B, are moving in the directions shown. What should be the angle  $\theta$  so that  $v_{B/A}$  is minimum?

A)  $0^\circ$

B)  $180^\circ$

C)  $90^\circ$

D)  $270^\circ$




2. Determine the velocity of plane A with respect to plane B.

A)  $(400 \mathbf{i} + 520 \mathbf{j}) \text{ km/hr}$


B)  $(1220 \mathbf{i} - 300 \mathbf{j}) \text{ km/hr}$

C)  $(-181 \mathbf{i} - 300 \mathbf{j}) \text{ km/hr}$

D)  $(-1220 \mathbf{i} + 300 \mathbf{j}) \text{ km/hr}$



## GROUP PROBLEM SOLVING



**Given:** Car A moves in a straight line while Car B moves along a curve having a radius of curvature of 200 m.

$$v_A = 40 \text{ m/s}$$

$$v_B = 30 \text{ m/s}$$

$$a_A = 4 \text{ m/s}^2$$

$$a_B = -3 \text{ m/s}^2$$

**Find:**  $v_{B/A}$   
 $a_{B/A}$

**Plan:** Write the velocity and acceleration vectors for Cars A and B. Determine  $v_{B/A}$  and  $a_{B/A}$  by using vector relationships.

## GROUP PROBLEM SOLVING (continued)


### Solution:

The velocity of B is:

$$\begin{aligned}\mathbf{v}_B &= -30 \sin(30) \mathbf{i} + 30 \cos(30) \mathbf{j} \\ &= (-15 \mathbf{i} + 25.98 \mathbf{j}) \text{ m/s}\end{aligned}$$

The velocity of A is:

$$\mathbf{v}_A = 40 \mathbf{j} \text{ (m/s)}$$




The relative velocity of B with respect to A is ( $\mathbf{v}_{B/A}$ ):

$$\mathbf{v}_{B/A} = \mathbf{v}_B - \mathbf{v}_A = (-15 \mathbf{i} + 25.98 \mathbf{j}) - (40 \mathbf{j}) = (-15 \mathbf{i} - 14.02 \mathbf{j}) \text{ m/s}$$

or  $v_{B/A} = \sqrt{(-15)^2 + (-14.02)^2} = \underline{20.5 \text{ m/s}}$

$$\theta = \tan^{-1}\left(\frac{14.02}{15}\right) = \underline{43.1^\circ}$$




## GROUP PROBLEM SOLVING (continued)

Since car  $B$  is traveling along a curve, the acceleration of  $B$  is:

$$\begin{aligned}\mathbf{a}_B &= (\mathbf{a}_t)_B + (\mathbf{a}_n)_B = [ -(-3) \sin(30) \mathbf{i} + (-3) \cos(30) \mathbf{j} ] \\ &\quad + \left[ -\left(\frac{30^2}{200}\right) \cos(30) \mathbf{i} - \left(\frac{30^2}{200}\right) \sin(30) \mathbf{j} \right]\end{aligned}$$

$$\mathbf{a}_B = ( -2.397 \mathbf{i} - 4.848 \mathbf{j} ) \text{ m/s}^2$$

The acceleration of  $A$  is:  $\mathbf{a}_A = (4 \mathbf{j}) \text{ m/s}^2$




The relative acceleration of  $B$  with respect to  $A$  is:

$$\mathbf{a}_{B/A} = \mathbf{a}_B - \mathbf{a}_A = ( -2.397 \mathbf{i} - 8.848 \mathbf{j} ) \text{ m/s}^2$$

$$a_{B/A} = \sqrt{(-2.397)^2 + (-8.848)^2} = \underline{9.17 \text{ m/s}^2}$$

$$\beta = \tan^{-1}(8.848 / 2.397) = \underline{74.8^\circ}$$






## ATTENTION QUIZ


1. Determine the relative velocity of particle B with respect to particle A.

- A)  $(48\mathbf{i} + 30\mathbf{j})$  km/h
- B)  $(-48\mathbf{i} + 30\mathbf{j})$  km/h
- C)  $(48\mathbf{i} - 30\mathbf{j})$  km/h
- D)  $(-48\mathbf{i} - 30\mathbf{j})$  km/h



2. If theta equals  $90^\circ$  and A and B start moving **from the same point**, what is the magnitude of  $\mathbf{r}_{B/A}$  at  $t = 5$  s?

- A) 20 ft
- B) 15 ft
- C) 18 ft
- D) 25 ft



*End of the Lecture*

*Let Learning Continue*