




[0293] Synthetic Example E: Synthesis of Example 71

[0294] Aniline (22 mg, 0.241 mmol) and pyridine (0.097 mL, 1.20 mmol) are suspended in CH₂Cl₂ (5 mL) and cooled to 0 °C and stirred for 5 minutes. To the reaction mixture **R-9** (50 mg, 0.241 mmol) is added and the reaction mixture is stirred for 10 minutes. The reaction mixture is concentrated in vacuo. The crude product is purified by preparative HPLC to afford **Example 71** (22 mg, 35%).

[0295] Synthetic Example F: Synthesis of Example 11

[0296] **R-10** (50 mg, 0.359 mmol), **R-11** (69 μ L, 0.539 mmol) are dissolved in CH₂Cl₂ (2 mL). Triethylamine (0.15 mL, 1.08 mmol) is added and the reaction mixture is stirred at room temperature for 17 hrs. The mixture is washed with water (2 mL) then passed through a Telos phase separator. Additional CH₂Cl₂ (2 mL) is passed through the phase separator. The combined organic layers are concentrated in vacuo to give the crude product. The crude material is purified by preparative HPLC to afford **Example 11** (6.0 mg, 5.9%).

[0297] The following examples are prepared in similar fashion from the appropriate aniline and sulphonyl chloride: **Examples 17, 195, and 207-209**.

[0298] Synthetic Example G: Synthesis of Example 23

[0299] **R-12** (5.00 g, 20.6 mmol), triethylamine (8.6 mL, 61.8 mmol) and **R-13** (2.29 g, 20.6 mmol) are suspended in DMF (10 mL). The reaction mixture is sealed under a nitrogen atmosphere and stirred at 80 °C for 1 hour then cooled to room temperature and partitioned