

Understanding the CREB1-miRNA feedback loop in human malignancies

Ya-Wen Wang^{1,2} · Xu Chen¹ · Rong Ma² · Peng Gao¹

Received: 19 November 2015 / Accepted: 1 April 2016 / Published online: 9 April 2016
© International Society of Oncology and BioMarkers (ISOBM) 2016

Abstract cAMP response element binding protein 1 (CREB1, CREB) is a key transcription factor that mediates transcriptional responses to a variety of growth factors and stress signals. CREB1 has been shown to play a critical role in development and progression of tumors. MicroRNAs (miRNAs) are a class of non-coding RNAs. They post-transcriptionally regulate gene expression through pairing with the 3'-UTR of their target mRNAs and thus regulate initiation and progression of various types of human cancers. Recent studies have demonstrated that a number of miRNAs can be transcriptionally regulated by CREB1. Interestingly, CREB1 expression can also be modulated by miRNAs, thus forming a feedback loop. This review outlines the functional roles of CREB1, miRNA, and their interactions in human malignancies. This will help to define a relationship between CREB1 and miRNA in human cancer and develop novel therapeutic strategies.

Keywords CREB1 · miRNA · Transcription factor · Feedback loop · Cancer

Electronic supplementary material The online version of this article (doi:10.1007/s13277-016-5050-x) contains supplementary material, which is available to authorized users.

✉ Peng Gao
gaopeng@sdu.edu.cn

¹ Department of Pathology, School of Medicine, Shandong University, Jinan 250012, People's Republic of China

² Department of Breast Surgery, Qilu Hospital of Shandong University, Jinan 250012, People's Republic of China

Introduction

Cancer is a complex disease that arises through a multi-step process. Acquisition of cancer cell features involves changes in the wiring of signaling pathways that are tightly regulated to control processes such as cell proliferation, apoptosis, and metastasis [1]. Many of the signal transductions assemble to modulate gene expression at transcriptional level via activation or repression of transcription factors. Under different pathological conditions, the transcription factors aberrantly initiate the transcription of cancer-related genes, thereby increasing cell proliferation, differentiation, angiogenesis, and suppressing apoptosis or cell-cell adhesion, which in turn lead to cancer development [2, 3].

cAMP response element binding protein 1 (CREB1, CREB) is a key transcription factor belonging to the basic leucine zipper (bZIP) family [4]. CREB1 is activated through phosphorylation at Ser133 and/or by nuclear translocation of CREB1 coactivators [5]. It has been found to mediate transcriptional responses to a variety of growth factors and stress signals [6] (Fig. 1). Recently, the roles of CREB1 in tumorigenesis have also been established in many types of solid tumors and hematological malignancies [6, 7]. Genome-wide studies have shown that the number of putative CREB1 target genes was approximately 4000 or nearly one-quarter of the human genome [8, 9]. Researches on the targets of CREB1 used to be limited to protein-coding genes [10, 11]; however, the non-coding targets deserve further investigation.

MicroRNAs (miRNAs) are a class of non-coding RNAs that post-transcriptionally regulate gene expression through imperfect pairing with the 3'-UTR of their target mRNAs, by inducing translational repression or degradation of mRNAs [12]. MiRNAs have critical functions across various biological processes, including cell development, differentiation, apoptosis, and proliferation. First identified in B cell




Fig. 1 A schematic model of CREB1 regulation in human cancers. CREB1 binds to the conserved cAMP-responsive element (CRE) on the promoter and mediates transcriptional responses to a variety of stimuli including neurotransmitters, hormones, membrane depolarization, and growth and neurotrophic factors, thereby acting as a mediator between different signal pathways and the downstream target genes transcription. The targets of CREB1 including both protein-coding (e.g., Bcl-2, cyclin D1, MMPs) and non-coding genes (e.g., miR-9, miR-23a, miR-373).

Interestingly, CREB1 could also be negatively regulated by miRNAs (e.g., miR-9, miR-34b, miR-200b), forming a negative feedback loop. MiR-27b directly inhibits CREB1 expression and contains putative CREB1 binding sites. Whether miR-27b is transcriptionally induced by CREB1 deserves further experimental validation. Illustration of the upstream regulators of CREB1 was partially modified from pathway information provided by BioCarta (http://cgap.nci.nih.gov/Pathways/BioCarta_Pathways)

chronic lymphocytic leukemia (CLL) [13], miRNA alterations have been involved in the initiation and progression of many types of human cancers [14–21]. MiRNA has also been associated with diagnosis, prognosis, and response to treatment in human malignancies [15, 18]. MiRNAs may function as oncogenes or tumor suppressors in the majority of human cancers [19]. For example, miR-15a and miR-16-1 generally function as tumor suppressors and downregulate BCL2 in chronic lymphocytic leukemias and cyclin D1 in prostate cancer and mantle cell lymphoma [22]. Of particular interest is miR-21 which tends to function as an oncogene. This miRNA has been shown to be consistently upregulated in various cancers including lung cancer, stomach cancer, prostate cancer, colon cancer, ovarian carcinoma, B cell lymphoma, hepatocellular carcinoma (HCC), cervical cancer, chronic lymphocytic leukemia, and breast cancer [23]. All in all, alterations of miRNA expression have been reported in a variety of cancers and are implicated in various processes during tumor initiation and progression.

Although many reasons, such as genetic abnormalities, epigenetic regulation, post-transcriptional modulation, as well as transcription factor control have been described, the causes of miRNA aberrant expression in tumors are not well understood

[19]. The promoter regions of miRNA genes are highly similar to those of protein-coding genes, which contain CpG islands, TATA box sequences, initiation elements, and certain histone modifications [24, 25]. A myriad of transcription factors are involved in miRNA genes transcription in cells [26, 27]. The transcription factors regulate miRNA expression positively or negatively in a tissue-specific or development-specific manner. For example, MYC stimulates expression of the miR-17-92 oncogenic cluster in lymphoma cells [27] but inhibits expression of several tumor suppressor miRNAs (for example, miR-15a), which promote MYC-mediated tumorigenesis [28]. Recently, CREB1 has been reported to modulate the expression of several miRNAs in human cancers, including miR-9 in glioma [29] and miR-373 in pancreatic cancer [30]. Interestingly, miRNAs frequently act with CREB1 in regulatory networks via feedback loops. For instance, miR-9, which is under CREB1's control, directly targets CREB1 to inhibit the proliferation of glioma cells [29]. In light of the rapid pace of research into CREB1 and miRNAs in human cancers, we perform this review for a better understanding of the CREB1-miRNA feedback loop.

Expression and function of CREB1 in human cancers

Accumulating evidence has shown that CREB1 is over-expressed in numerous human cancers, including astrocytoma, breast cancer, gastric cancer, glioma, HCC, mesothelioma, non-small cell lung cancer (NSCLC), ovarian cancer, and prostate cancer (Table 1). CREB1 is a transcription factor which functions as an oncogene that involves in the proliferation, survival, and metastasis of tumor cells [47]. For instance, CREB1 is elevated in breast cancer tissues and metastatic cancer cells and associated with poor prognosis, metastatic potential, and nodal involvement [32, 33]. CREB1 is further shown to positively regulate the proliferation, metastasis, and bone destruction of breast cancer [32]. Seo et al. shows that CREB1 is highly expressed in most of the NSCLC cell lines and associates with decreased survival in patients with NSCLC [43]. CREB1 inactivation suppresses the growth of NSCLC cells and induces apoptotic cell death [42]. Tan and colleagues find that CREB1 is over-expressed in gliomas, and promotes gliomagenesis by inducing the expression of oncogenic miR-23a, which represses the tumor suppressor PTEN [37]. More recently, Barresi et al. find the correlation between high expression of phosphorylated CREB1 (p-CREB1) and angiogenesis, recurrence risk, and poor disease-free survival in patients with meningiomas [40]. They also demonstrate that astrocytomas and oligodendrogliomas are characterized by distinctive patterns of p-CREB1 expression, which provide the evidence that CREB1 may function as a useful tool in the differential diagnosis of these two main types of gliomas [48]. Previous studies from our laboratory and others indicate that CREB1 is positively correlated with tumor metastasis, tumor stage, poor prognosis, and cancer cell growth in gastric cancer [36, 49]. In addition to its dysregulation in solid tumors, CREB1 over-expression is also associated with poor outcome of acute myelocytic leukemia patients and promotes abnormal proliferation and survival of myeloid cells in vitro and in vivo [7].

In spite of all the above findings, there is also evidence indicating that CREB1 could function as a tumor suppressor. In line with this idea, CREB1 has been shown to suppress AChE-R-induced, PKA-mediated proliferation of glioblastoma tumors [50]. It is interesting that CREB1 plays a pro-proliferative and anti-migratory role in glioma cells [29]. These data suggest that CREB1's role in human cancers is tissue- and/or cell type-specific, though it typically serves as a proto-oncogenic transcription factor.

CREB1 acts as a miRNA transcription activator

A precise control of miRNA levels is essential for maintaining normal cellular homeostasis and uncontrolled deregulation of

miRNA may contribute to tumorigenesis. Interestingly, recent ChIP-seq (chromatin immunoprecipitation sequencing) data has provided evidence that CREB1 can bind to the promoter of several miRNAs in glioma cells and regulate their transcription [37]. In the following section, we will describe some of the verified or predicted CREB1-miRNA interactions in human cancers (Table 2, Fig. 1).

let-7i

Let-7i belongs to the let-7 family, which include 12 human homologues and are considered as tumor suppressors because they are frequently located at fragile sites and genomic regions involved in cancers [14]. Downregulation of let-7i has been correlated with tumor progression and anti-cancer drug resistance in many human malignancies [55–58]. Several groups have identified let-7i as a metastasis-specific miRNA biomarker associated with prognosis in colorectal cancer (CRC) and let-7i is validated to target TRIM41, SOX13, SLC25A4, SEMA4F, RPUSD2, PLEKHG6, CCND2, and BTBD3 in CRC cell lines [59, 60]. Recent evidence has also established let-7i as a transcriptional target of mutant p53 and repression of let-7i exhibits a key role in enhancing migration, invasion, and metastasis of human cancer [61]. Although let-7i plays an important role in various cancers, little is known about how let-7i expression is regulated. By means of bioinformatics-based analysis, Dai et al. find that let-7i is downregulated in apurinic/apyrimidinic endonuclease1 (APE1) knockdown human osteogenic sarcoma cells and contains two putative CREB1 binding sites in its promoter. This may indicate the potential function of CREB1 mediated by APE1 in the regulation of let-7i expression [51].

miR-9

MiR-9 has been reported to exert distinct and even opposite functions in different types of tumors through targeting different cellular genes. MiR-9 is upregulated in bladder cancer and promotes cell proliferation, cell cycle progression, invasion, and chemo-resistance, partly through directly downregulating LASS2 and CBX7 [62, 63]. Over-expression of miR-9 targets PPARA and CDH1 and induces HCC cell growth and invasiveness [64]. Cai et al. shows that high expression of miR-9 is correlated with aggressive clinic-pathological features and poor prognosis in HCC patients [65]. In contrast, Higashi and colleagues identify miR-9 as a tumor suppressor, which inhibits cell proliferation via targeting TAZ in HCC cells [66]. We suspect that different detection methods and specimens from different ethnic groups may account for the discrepancy. Recently, miR-9 is proved to inhibit the proliferation of glioma cells by inhibiting CREB1 expression [29]. Moreover, miR-9 is transcriptionally activated by CREB1, forming a negative feedback minicircuitry [29].

Table 1 Expression and function of CREB1 in human cancers

Cancer/cell types	CREB1 level	Sample size	Detection methods	Functions	Correlation with metastasis or prognosis	CREB1's targets	References
Astrocytoma	Upregulated	Astrocytomas ($n = 122$) v.s. non-neoplastic specimens ($n = 30$)	qRT-PCR	N.A.	Poor prognosis	N.A.	[31]
Breast cancer cells	Upregulated	Metastatic MDA-MB-231 v.s. non-metastatic MCF-7	WB	Increases proliferation, migration, invasion, and bone destruction	Promotes metastasis	PTHrP, MMPs, and OPG	[32]
Breast cancer tissues	Upregulated	Tumor ($n = 120$) v.s. non-neoplastic ($n = 33$)	IHC and qRT-PCR	N.A.	Metastasis and poor prognosis	N.A.	[33]
Breast cancer	N.A.	N.A.	N.A.	Transcriptional regulation of LAPTM4B	N.A.	LAPTM4B	[34]
Gastric cancer	N.A.	Cancer tissues ($n = 66$) v.s. normal gastric mucosa	IHC on TMA	Increases cell proliferation	N.A.	C/EBPbeta	[35]
Gastric cancer	Upregulated	Normal ($n = 50$) v.s. primary tumor ($n = 185$) v.s. metastatic foci ($n = 50$)	IHC	N.A.	Metastasis and poor prognosis	N.A.	[36]
Glioma cells	Upregulated	Glioma cell lines ($n = 4$) v.s. normal human glial cell line ($n = 1$)	qRT-PCR	Inhibits migration	N.A.	miR-9 and NF1	[29]
Glioma tissues	Upregulated	Glioma ($n = 15$) v.s. normal ($n = 2$)	WB and qRT-PCR	Increases growth and survival	N.A.	miR-23a	[37]
HCC	Upregulated	HCC ($n = 130$) v.s. non-neoplastic tissues ($n = 130$)	IHC and WB	N.A.	Poor prognosis	N.A.	[38]
Melanoma	N.A.	N.A.	N.A.	Promotes growth and metastasis	N.A.	CYR61	[39]
Meningioma	Upregulated	Meningiomas ($n = 70$) v.s. normal leptomeninges ($n = 10$)	IHC	N.A.	Angiogenesis, recurrence risk, and poor prognosis	N.A.	[40]
Mesothelioma	Upregulated	Mesothelioma sections ($n = 33$) v.s. normal lung sections ($n = 7$) v.s. reactive mesothelial hyperplasias ($n = 3$)	IHC on TMA	Increases migration and decreases apoptosis	N.A.	Bcl-2, c-FOS and MMP13, etc.	[41]
NSCLC cells	Upregulated	Four NSCLC cell lines v.s. NHTBE (normal) cells	WB	Increases growth and survival	N.A.	Bcl-2 and Bcl-xL	[42]
NSCLC tissues	Upregulated	Tumor ($n = 51$) v.s. normal ($n = 51$)	WB, PCR, and IHC	N.A.	Decreased overall survival	N.A.	[43]
Ovarian cancer	Upregulated	Serous adenocarcinoma ($n = 124$) v.s. mucous adenocarcinoma ($n = 25$) v.s. normal tissues ($n = 8$)	IHC on TMA	Promotes proliferation	Associated with tumor stage	N.A.	[44]
Prostate cancer	Upregulated	Normal/benign glands ($n = 5$) v.s. primary cancers ($n = 10$) v.s. bone metastatic tissues ($n = 7$)	IHC	Induces VEGF transcription	Bone metastasis	HIF-1a and VEGF	[45]
Prostate cancer	Upregulated	Cancer specimens ($n = 15$) v.s. normal prostate epithelium ($n = 15$)	IHC on TMA	Transcriptional regulation of cyclin D1	N.A.	Cyclin D1	[46]

HCC hepatocellular carcinoma, NSCLC non-small cell lung cancer, qRT-PCR quantitative real-time polymerase chain reaction, IHC immunohistochemistry, WB Western blot, N.A. not available or not associated, TMA tissue microarray

Table 2 CREB1-regulated miRNAs in human cancers

miRNA	Cancer/cell types	Validation methods of CREB1's regulation on miRNA	Regulation manner	miRNA alterations	Cell functions driven by miRNAs	Target genes of miRNA	References
let-7i	Osteogenic sarcoma	Bioinformatics analysis	Activation	Downregulated in APE1 knockdown cells	N.A.	APE1 and TLR4	[51]
miR-9	Glioma	Bioinformatics analysis, ChIP-qPCR, and qRT-PCR	Feedback loop	Upregulated in cancer	Inhibits growth but stimulates migration	CREB1 and NF1	[29]
miR-21	Breast and ovarian	Bioinformatics analysis, luciferase assay, and ChIP	Activation	N.A.	Promotes resistance to hypoxia and cell survival	PTEN, PDCD4, and Spry1	[52]
miR-23a	Glioma	MicroRNA microarray, ChIP-chip, and ChIP-qPCR	Activation	Upregulated in cancer	Increases growth and survival	FOXO3a and PTEN	[37]
miR-320a	Cervical	Bioinformatics analysis, qRT-PCR, luciferase assay, and ChIP	Activation	Upregulated in cancer	Promotes mitophagy	VDAC1	[53]
miR-335	HeLa cells	qRT-PCR and ChIP	Activation	Downregulated post irradiation	DNA repair, intra S-phase checkpoints, and radiosensitivity	CtIP	[54]
miR-373	Pancreatic	Luciferase assay, ChIP, and qRT-PCR	Activation	N.A.	Enhancement of cell proliferation, invasion, and tumor growth	TP53INP1, LATS2, and CD44	[30]

ChIP chromatin immunoprecipitation

miR-21

MiR-21 is over-expressed in most of the solid tumors. It regulates various cellular processes, such as cell cycle, proliferation, apoptosis, invasion, and drug resistance, by targeting multiple tumor suppressor genes [17, 67]. Recent studies have shown that miR-21 might be a biomarker for diagnosis and prognosis in cancer patients [21, 68]. One of the most important proof of miR-21's oncogenic activity may come from the Slack laboratory who showed that over-expression of miR-21 resulted in a pre-B cell lymphoma by a genetically engineered miR-21 mouse model. This experiment illustrates the significant impact of a single miRNA at all stages of in vivo tumor development: initiation, maintenance and survival [69]. As a mechanical study for upstream pathways of miR-21, Polyarchou et al. find that the miR-21 induction by Akt2 during hypoxia partially depends upon the binding of CREB1 to the miR-21 promoter [52]. Interestingly, miR-21 seems to be involved in a number of positive and negative feedback loops forming with transcription factors, and these complex regulations may explain why miR-21 is probably one of the most dynamic miRNAs responsive to various stimuli [70]. One evolutionary conserved double-negative feedback module is miR-21 and its direct target NFIB, a transcription factor involved in carcinogenesis [71].

miR-23a

MiR-23a belongs to the miR-23a-27a-24-2 cluster which locates in chromosome 19p13.12. It is among the most dysregulated miRNAs implicated in cancer. miR-23a, over-expressed in gastric cancer, has been reported to participate in the FasL-induced epithelial-mesenchymal transition (EMT) process [72] and confer poor prognosis in patients with gastric cancer [73]. Moreover, miR-23a is also demonstrated to stimulate TGF-beta-induced EMT by targeting E-cadherin [74], and promote the migration and invasion of A549 cells via its target IRS-1 [75] in lung cancer. Recently, miR-23a has been identified to repress the tumor suppressor PTEN and confirmed as a functional downstream target of CREB1 in controlling glioma cell growth and cell survival [37]. However, Cai et al. find that miR-23a is decreased in prostate cancer and inhibits cancer metastasis though the PAK6-LIMK1 pathway [76], suggesting a tissue- or cell-specific role of miR-23a in cancers.

miR-320a

MiR-320a, belonging to the miR-320 family (miR-320a~e), has been shown to be dysregulated in many human malignancies. MiR-320a typically functions as a tumor suppressive miRNA and has emerged as a key regulator of tumor initiation, progression, metastasis, as well as drug resistance.

Human miR-320a gene is localized at chromosome 8p21.3, which is frequently deleted during CRC progression and identified as a liver metastatic susceptibility locus [77, 78], thus miR-320a has been extensively investigated in CRC [60, 79–83]. Notably, miR-320a is found to be down-regulated in liver metastases tissues compared to matched primary CRC tissues [60, 81] and high expression of miR-320a is associated with better disease-free survival in CRC patients [84]. Functional studies have revealed that miR-320a decreases cell proliferation, migration, and invasion, induces G0/G1 growth arrest, and sensitizes chemoradiotherapy in CRC by targeting neuropilin 1, beta-catenin, and Rac1 [80–83]. Interestingly, miR-320a is identified as a direct target of transcriptional factor E2A and is involved in colon cancer cell growth [79]. In addition, miR-320a exerts its suppressive function by targeting BCR/ABL oncogene in chronic myeloid leukemia [85], BMI-1 in nasopharyngeal carcinoma [86], GNAI1 in hepatocellular carcinoma [87], IGF-1R in glioblastoma [88], ITGB3 in salivary adenoid cystic carcinoma [89], and ITGB3 in bladder cancer [90]. Intriguingly, CREB1 is found to activate miR-320a expression and induce mitophagy during serum starvation in cervical cancer cells and the CREB1-miR-320a-VDAC1 axis may provide new insights into cancer cell survival in response to environmental stresses [53].

miR-335

MiR-335 has been identified to be involved in tumorigenesis of numerous cancers such as ovarian cancer, lung cancer, HCC, prostate cancer, breast cancer, cervical cancer, neuroblastoma, pancreatic cancer, clear cell renal cell carcinoma, meningioma, glioma, esophageal squamous cell carcinoma, and gastric cancer [91–110]. The expression and function of miR-335 in different human cancers are highly dependent on the context or types of tumors. For example, Zhou [91] and Yang et al. [96] have demonstrated that the expression of miR-335 is significantly reduced in gastric cancer, and low expression of miR-335 is associated with aggressive clinical features and invasion and metastasis. However, miR-335 seems to be a tumor promoter in brain cancers including meningioma [105] and glioma [109]. Over-expressed miR-335 promotes cell proliferation and invasion by targeting Rb1 and PAX6 in meningioma and glioma, respectively. As to the regulation of CREB1 on miR-335, Martin et al. find that CREB1 could bind to the promoter region of miR-335 and induce miR-335 expression in HeLa cells [54]. They finally propose an ATM-dependent CREB-miR-335-CtIP axis that influences homologous recombination repair (HRR) of certain double-strand break (DSB) lesions.

miR-373

MiR-373 is located in the chromosomal band 19q13.4 and belongs to the miR-371-3 gene cluster (miR-371, miR-372, miR-373, and miR-373*) [111, 112]. MiR-373 was firstly identified as one of the human embryonic stem cell (ESC)-specific miRNAs [112]. Subsequently, it is proved to be an oncogene involving in the proliferation and tumorigenesis of human testicular germ cell tumors by numbing the p53 pathway [111]. Later, numerous studies have shown the pivotal roles of miR-373 in regulation of cell proliferation, apoptosis, senescence, migration, invasion, chemo-sensitivity, as well as DNA damage repair following hypoxia stress [113]. A recent study from Adi Harel et al. have revealed that miR-373 increases cisplatin-induced apoptosis and inhibits cell migration by targeting RelA and PIK3CA in lung cancer cells [114]. Intriguingly, miR-373 enhances cell proliferation, invasion, and tumor growth in pancreatic cancer and is transcriptionally activated by CREB1 through the regulation of miR-373's promoter [30].

CREB1 acts as a direct target of miRNA

MiRNAs regulate their targets by direct cleavage of the mRNA or by inhibition of protein synthesis, according to the degree of complementarities with their targets' 3'-UTR regions [20] and are thereby involved in the initiation and progression of human malignancies. As a key cancer-related pro-oncogenic gene, CREB1 has been predicted and validated to be direct targets of many miRNAs (Table 3, Fig. 1).

miR-17-92 cluster

MiR-17-92, a miRNA polycistron also known as oncomir-1, is among the most potent oncogenic miRNAs [124]. The miR-17-92 cluster is composed of seven miRNAs (miR-17-5p, miR-17-3p, miR-18a, miR-19a, miR-20a, miR-19b-1, and miR-92a-1) [115]. He et al. show that the miR-17-92 polycistron locating at 13q31-32 is upregulated in 65 % of the B cell lymphoma [125]. They also find miR-17-92 cluster acts together with MYC to accelerate tumor development [125]. The oncogenic role of the miR-17-92 cluster is further supported by profiling studies showing upregulation of this cluster in cancers and evidences indicating that miR-17-92 promotes proliferation, inhibits differentiation, increases angiogenesis, and sustains cell survival [124, 126]. Attar et al. show that miR-17-92 cluster over-expression in HEK-293 cells enhances cell proliferation and cell cycle-related genes, including CREB1, CDK2, cyclin-D2, and c-Myc, which were putative targets of miR-17-92.

Table 3 miRNAs directly target CREB1 in human cancers

miRNA	Cancer types	miRNA alterations in cancer	Validation methods of miRNA-CREB1 interaction	Regulation manner	Functions mediated by CREB1	Downstream effectors of CREB1	References
miR-9	Glioma	Upregulated	Luciferase assay and WB	Feedback loop	Enhances proliferation and inhibits migration	NF1	[29]
miR-17-92 cluster	HEK-293 T cells	N.A.	qRT-PCR and WB	Increase	Increases cell proliferation	N.A.	[115]
miR-27b/miR-200b	Gastric	N.A.	Prediction algorithms, luciferase assay, qRT-PCR, and WB	Inhibition	N.A.	N.A.	[36]
miR-34b	Acute myeloid leukemia	Downregulated	Luciferase assay and WB	Inhibition	Increases cell proliferation and clonogenicity	Cyclin A1, cyclin B1, cyclin D1, BCL-2, STAT3, JAK1, NF- κ B, etc.	[116, 117]
miR-181b	Gastric	Downregulated	Fluorescent reporter assay, WB, qRT-PCR, and rescue experiments	Inhibition	Increases cell growth	N.A.	[118]
miR-182	Gastric	Downregulated	Fluorescent reporter assay, WB, qRT-PCR, and rescue experiments	Inhibition	Increases cell growth	N.A.	[49]
miR-200b	Glioma	Downregulated	Prediction algorithms, luciferase assay, qRT-PCR, WB, and rescue experiments	Inhibition	Increases cell growth	N.A.	[31, 119]
miR-203	Melanoma	Hypermethylation/downregulated	Bioinformatics algorithms, luciferase reporter assay, and WB	Inhibition	Increases cell proliferation	MITF and RAB27a	[120–122]
miR-433	HCC	Hypermethylated	Bioinformatics algorithms, luciferase reporter assay, WB, and rescue experiments	Inhibition	Increases cell proliferation, migration, and invasion	N.A.	[123]

miR-27b

As a member of the miR-23b/27b cluster, miR-27b is frequently downregulated in various types of human cancers [127]. Expression of miR-27b is significantly reduced in prostate cancer, especially in metastatic, castration-resistant tumors [128, 129]. MiR-27b significantly inhibits cell proliferation, migration, and invasion in prostate cancer cells by targeting GOLM1 [128] and Rac1 [129]. Similarly, miR-27b-mediated suppression of Sp1 [130] and LIMK1 [131] attenuate the growth and invasion of NSCLC cells. miR-27b has also been reported to enhance drug response by activating p53-dependent apoptosis and reducing CYP1B1-mediated drug detoxification in liver and kidney cancers [132]. However, miR-27b may act as a pro-oncogenic player in several tumors, such as HPV-associated cervical cancer [133], glioma [134], and breast cancer [126, 135]. In addition, we and other groups have identified CREB1 as direct target of miR-27b in gastric carcinogenesis [36] and adipogenesis [136]. In our previous study, we identified CREB1 as a novel target of miR-27b in gastric cancer [36]. However, our recent work suggested a positive correlation between CREB1 and miR-27b in gastric cancer tissues (Online Resource Figure S1A), which were counter to the proposed inverse relationship between miR-27b and its target CREB1. One possible explanation is that CREB1 may act as a transcriptional activator to induce miR-27b expression. To support this hypothesis, we utilized miRStart [137], a source of human microRNA TSSs (transcription start sites, <http://mirstart.mbc.nctu.edu.tw/home.php>), to get the putative TSSs and promoter sequences of miR-27b. Then, we used RegRNA2.0 [138] and PROMO database [139] to predict potential binding sites for CREB1. The analysis revealed the presence of several CREB1-binding sites in the putative promoter of miR-27b (Figure S1B, C). However, further experiments are needed to validate the relationship between CREB1 and miR-27b.

miR-34b

MiR-34b belongs to the evolutionary conserved miRNA family miR-34s [140], which are known to be the transcriptional targets of tumor suppressor p53 [141] and are regulated by DNA hypermethylation [142]. MiR-34b has been implicated in the oncogenesis of colon [143], ovarian [144], prostate [145, 146], hepatocellular [147], lung [148], pancreatic [149], breast [150], and oral cancers [151]. MiR-34b is prevalently identified as an important component of the tumor suppressor network during carcinogenesis; however, increased miR-34b is observed by Hiyoshi et al. in advanced colon tumors and is associated with poor survival of cancer patients [142]. Pigazzi et al. find that miR-34b directly target CREB1 in acute myeloid leukemia (AML) [116]. MiR-34b

expression can cause cell cycle abnormalities, reduce anchorage-independent growth, and alter CREB1 target gene expression in AML [116]. Interestingly, FOXO3a is transcriptionally responsible for miR-34b expression in breast cancer [150], further supporting the transcription factor-miR-34b interaction in human cancers.

miR-181b

MiR-181 family members contain four highly conserved mature miRNAs (miR-181a, miR-181b, miR-181c, and miR-181d) [152], which have been reported to contribute to tumor initiation and progression. Liu et al. briefly summarize the functional significance and clinical implications of miR-181b in cancer [153]. They propose that miR-181b is a critical link between inflammation and cancer, supported by the evidence that miR-181b inhibits CYLD to increase NF-kappaB activity, which is required for maintaining the transformed state [154] and regulating NF-kappaB-mediated vascular inflammation [155]. MiR-181b is over-expressed in ovarian [156], pancreatic [157], head and neck [158], and bladder cancers [159]. By contrast, miR-181b is downregulated in thyroid papillary cancer [160], glioma [161, 162], astrocytoma [163], acute myeloid leukemia [164], and prostate cancer [165]. MiR-181b is a key regulator of the oncogenic process involving cellular growth, apoptosis, invasion, metastasis, and drug sensitivity. Controversial evidence is observed by two research group reporting inconsistent expression and function of miR-181b in gastric cancers [118, 166]. Tang et al. indicates that miR-181b may function as a tumor suppressor in gastric cancer cells through directly targeting CREB1 [118]. However, Guo et al. suggests that miR-181b is over-expressed in gastric cancer cells and tissues and targets TIMP3 to increase cell proliferation, migration, and invasion [166]. Both of the two studies include relatively small sample size (10 to 12 pairs of gastric tissues) and meanwhile lack in vivo confirmation. We thus suggest that further studies are needed to validate the roles of miR-181b in gastric cancer.

miR-182

MiR-182, a member of the evolutionarily conserved miR-183-96-182 cluster, has been demonstrated to play important roles in a variety of cancers [167]. MiR-182 is upregulated in most types of tumors [168] and participate in apoptosis, growth, differentiation, metastasis and chemo-sensitivity of cancer cells [169–176]. For example, CEBPA and RASA1 targeting miR-182 has been shown to be upregulated and promote angiogenesis in HCC [171, 177]. By comparing the expression of miRNAs in metastatic and non-metastatic primary mouse sarcomas, Sachdeva et al. find that miR-182 is over-expressed in tumors metastasized to the lungs [172]. MiR-182 is identified as a metastasis driver of primary

sarcomas by targeting multiple genes [172]. However, reports on the expression and function of miR-182 in gastric cancer are controversial [167]. Kong [49] and Tang et al. [178] find that miR-182 is downregulated in gastric carcinoma and inhibits proliferation through targeting oncogenic CREB1 and ANUBL1. Li et al. report that miR-182 is upregulated in intestinal-type gastric cancers and increases along with tumor stage [179]. Similar to miR-181b, larger samples and in vivo assays may be useful to further elucidate the exact roles of miR-182 in gastric cancer.

miR-200b

Members of the miR-200 miRNA family (miR-200a, miR-200b, miR-200c, miR-141, and miR-429) are mainly downregulated in human cancer cells and tissues and play a crucial role in the suppression of EMT, tumor cell adhesion, migration, invasion, metastasis, and cancer chemo-sensitivity [19, 180, 181]. MiR-200b is found to be lost in invasive breast cancer cell lines with mesenchymal phenotype and prevent TGF-beta-induced EMT and tumor metastasis, by targeting the E-cadherin transcriptional repressors ZEB1 and ZEB2 [182]. Additionally, ZEB1 and ZEB2 have been found to bind directly to an E-box proximal minimal promoter element and repress miR-200b expression in mesenchymal human breast cancer cells, demonstrating a potential double-negative feedback loop between ZEB1/ZEB2 and miR-200b during EMT and tumorigenesis [183]. More recently, Wu et al. have also reported a negative feedback loop between miR-200b and the NF-kappaB pathway via IKBKB/IKK-beta in breast cancer cells [184]. They demonstrate that miR-200b, a transcriptional target of NF-kappaB, suppresses breast cancer cell growth and migration as well as NF-kappaB activation through the downregulation of IKBKB. Interestingly, the transcription factor CREB1 is proved to be a direct target of miR-200b in glioma [119] and gastric cancer [36]. Whether there exists a feedback loop between transcription factor CREB1 and miR-200b warrants further investigation.

miR-203

MiR-203 has been reported to be a tumor suppressor and EMT-associated miRNA which is silenced in different malignancies. It is reported that miR-203 inhibits cell proliferation, adhesion, migration, invasion, metastasis as well as enhances chemo-sensitivity in cancer cells [120, 185–188]. MiR-203 suppression in gastric cancer promotes Slug-mediated cancer metastasis [189]. Similarly, miR-203 acts as a growth-suppressive miRNA in *Helicobacter pylori*-related gastric cancer by repressing CASK expression [186]. A retrospective study by Imaoka et al. shows that low serum miR-203 expression is an independent predictive marker for lymph node,

peritoneal, and distant metastases and associates with poor prognosis in patients with gastric cancer [185]. MiR-203 suppresses the proliferation, migration, and invasion and promotes the apoptosis of lung cancer cells by targeting Bmi1, FZD2, and SRC [190–192]. MiR-203 enhances 5-FU chemo-sensitivity via downregulating TYMS in CRC [193] and attenuates cell proliferation, invasion, and migration by suppressing ZNF217 [194]. MiR-203 is a tumor suppressor miRNA inhibiting cellular proliferation by directly targeting CREB1 [121] and regulating its downstream targets, MITF and RAB27a [120, 122] in melanoma cells.

miR-433

For the last decade, extensive research has identified the pivotal role of miR-433 in human cancers, including ovarian cancer [195, 196], gastric cancer [197], myeloproliferative neoplasms [198], lung cancer [199], and HCC [123]. Expression of miR-433 is downregulated in gastric carcinoma [200–202], which is associated with unfavorable overall survival [197]. MiR-433 inhibits gastric cancer cell proliferation, cell cycle progression, cell migration and invasion by directly interacting with oncogenic KRAS [202] and tumor-associated protein GRB2 [200]. MiR-433 is aberrantly expressed in myeloproliferative neoplasms and suppresses hematopoietic cell growth and differentiation by negatively regulating GBP2 [198]. High expression of miR-433 is associated with poor progression-free survival in patients with high-grade serous ovarian cancer and promotes resistance to paclitaxel through the induction of cellular senescence in ovarian cancer cells [195]. Yang et al. demonstrates that miR-433 inhibits HCC cell migration by repressing the protein expression and function of CREB1 [123].

Conclusions and perspectives

In this review, we briefly summarize the bidirectional interactions between CREB1 and miRNA in human cancers (Fig. 1). We believe that, except for miR-9, other single miRNA may also form a negative feedback loop with CREB1, which deserves further investigation. Both CREB1 and miRNA have been shown to regulate important cancer hallmarks, including proliferation, apoptosis, metastasis, and drug resistance. A recent integrative analysis of gene and miRNA expression profiles with transcription factor-miRNA feed-forward loops identifies CREB1 and its miRNA partners as master regulators across multiple cancer types [203]. Given its critical roles in cancer development and progression, the CREB1-miRNA network (Fig. 1) may represent novel therapeutic targets in human cancers. This network includes the important node—CREB1 itself, the miRNAs (e.g., miR-34b, miR-200b)

severing as upstream regulators of CREB1, and the CREB1 target coding genes (e.g., cyclin D1, BCL-2) and miRNAs (e.g., miR-23a, miR-373). On one hand, miRNA target prediction and validation, including the bioinformatic databases [204], have gain remarkable advances in recent years. We comprehensively searched four databases of experimentally validated miRNA target interactions (miRWalk 2.0, miRTarBase, miRecords, and TarBase 7.0) and found that nearly 100 miRNAs could potentially target CREB1 in human species. Although not all information in these databases are up-to-date or accurate, which need further improvement (see comments of Dweep H and Kenneth Witwer that appeared in PubMed Commons of [205]), they are still useful tools to find putative miRNA targets in the discovery stage of investigations. On the other hand, as more and more miRNAs are proved to be targets of transcription factor CREB1, we comment that in the future, CREB1 ChIP assays in combination with miRNA sequencing should be helpful to identify novel target miRNAs of CREB1. By fully understanding the regulatory networks of CREB1 and miRNA, we could be able to evaluate the potential of CREB1 and miRNA as therapeutic targets in human cancers.

Acknowledgments This work was supported by the National Natural Science Foundation of China (No. 81372856) and Taishan Scholars Program of Shandong Province (No. ts201511096). We thank Dr. Hsin-Sheng Yang (Graduate Center for Toxicology, University of Kentucky) and Dr. Tiantian Liu (Department of Pathology, School of Medicine, Shandong University) for their critical reading and suggestions for the manuscript.

Compliance with ethical standards

Conflicts of interest None

References

- Weinstein IB. Growth factors, oncogenes, and multistage carcinogenesis. *J Cell Biochem.* 1987;33(3):213–24. doi:10.1002/jcb.240330308.
- Ito Y. Signals and transcription factors. *Gan To Kagaku Ryoho Cancer Chemother.* 1989;16(3 Pt 2):509–15.
- Hsieh WT, Tzeng KR, Ciou JS, Tsai JJ, Kurubanjerdjit N, Huang CH, et al. Transcription factor and microRNA-regulated network motifs for cancer and signal transduction networks. *BMC Syst Biol.* 2015;9 Suppl 1:S5. doi:10.1186/1752-0509-9-s1-s5.
- Shaywitz AJ, Greenberg ME. CREB: a stimulus-induced transcription factor activated by a diverse array of extracellular signals. *Annu Rev Biochem.* 1999;68:821–61. doi:10.1146/annurev.biochem.68.1.821.
- Conkright MD, Canettieri G, Sreaton R, Guzman E, Miraglia L, Hogenesch JB, et al. TORCs: transducers of regulated CREB activity. *Mol Cell.* 2003;12(2):413–23.
- Siu YT, Jin DY. CREB—a real culprit in oncogenesis. *FEBS J.* 2007;274(13):3224–32. doi:10.1111/j.1742-4658.2007.05884.x.
- Shankar DB, Cheng JC, Kinjo K, Federman N, Moore TB, Gill A, et al. The role of CREB as a proto-oncogene in hematopoiesis and in acute myeloid leukemia. *Cancer Cell.* 2005;7(4):351–62. doi:10.1016/j.ccr.2005.02.018.
- Zhang X, Odom DT, Koo SH, Conkright MD, Canettieri G, Best J, et al. Genome-wide analysis of cAMP-response element binding protein occupancy, phosphorylation, and target gene activation in human tissues. *Proc Natl Acad Sci U S A.* 2005;102(12):4459–64. doi:10.1073/pnas.0501076102.
- Impey S, McCorkle SR, Cha-Molstad H, Dwyer JM, Yochum GS, Boss JM, et al. Defining the CREB regulon: a genome-wide analysis of transcription factor regulatory regions. *Cell.* 2004;119(7):1041–54. doi:10.1016/j.cell.2004.10.032.
- Maureira A, Sanchez R, Valenzuela N, Torrejon M, Hinrichs MV, Olate J, et al. The CREB transcription factor controls transcriptional activity of the human RIC8B gene. *J Cell Biochem.* 2016. doi:10.1002/jcb.25479.
- Mylroie H, Dumont O, Bauer A, Thornton CC, Mackey J, Calay D, et al. PKCepsilon-CREB-Nrf2 signalling induces HO-1 in the vascular endothelium and enhances resistance to inflammation and apoptosis. *Cardiovasc Res.* 2015;106(3):509–19. doi:10.1093/cvr/cvv131.
- Krol J, Loedige I, Filipowicz W. The widespread regulation of microRNA biogenesis, function and decay. *Nat Rev Genet.* 2010;11(9):597–610. doi:10.1038/nrg2843.
- Calin GA, Dumitru CD, Shimizu M, Bichi R, Zupo S, Noch E, et al. Frequent deletions and down-regulation of micro-RNA genes miR15 and miR16 at 13q14 in chronic lymphocytic leukemia. *Proc Natl Acad Sci U S A.* 2002;99(24):15524–9. doi:10.1073/pnas.242606799.
- Calin GA, Sevignani C, Dumitru CD, Hyslop T, Noch E, Yendamuri S, et al. Human microRNA genes are frequently located at fragile sites and genomic regions involved in cancers. *Proc Natl Acad Sci U S A.* 2004;101(9):2999–3004. doi:10.1073/pnas.0307323101.
- Garzon R, Fabbri M, Cimmino A, Calin GA, Croce CM. MicroRNA expression and function in cancer. *Trends Mol Med.* 2006;12(12):580–7. doi:10.1016/j.molmed.2006.10.006.
- Lu J, Getz G, Miska EA, Alvarez-Saavedra E, Lamb J, Peck D, et al. MicroRNA expression profiles classify human cancers. *Nature.* 2005;435(7043):834–8. doi:10.1038/nature03702.
- Volinia S, Calin GA, Liu CG, Ambs S, Cimmino A, Petrocca F, et al. A microRNA expression signature of human solid tumors defines cancer gene targets. *Proc Natl Acad Sci U S A.* 2006;103(7):2257–61. doi:10.1073/pnas.0510565103.
- Calin GA, Croce CM. MicroRNA signatures in human cancers. *Nat Rev Cancer.* 2006;6(11):857–66. doi:10.1038/nrc1997.
- Di Leva G, Garofalo M, Croce CM. MicroRNAs in cancer. *Annu Rev Pathol.* 2014;9:287–314. doi:10.1146/annurev-pathol-012513-104715.
- Bartel DP. MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell.* 2004;116(2):281–97.
- Bertoli G, Cava C, Castiglioni I. MicroRNAs: new biomarkers for diagnosis, prognosis, therapy prediction and therapeutic tools for breast cancer. *Theranostics.* 2015;5(10):1122–43. doi:10.7150/thno.11543.
- Chivukula RR, Mendell JT. Circular reasoning: microRNAs and cell-cycle control. *Trends Biochem Sci.* 2008;33(10):474–81. doi:10.1016/j.tibs.2008.06.008.
- Selcuklu SD, Donoghue MT, Spillane C. miR-21 as a key regulator of oncogenic processes. *Biochem Soc Trans.* 2009;37(Pt 4):918–25. doi:10.1042/bst0370918.
- Ozsolak F, Poling LL, Wang Z, Liu H, Liu XS, Roeder RG, et al. Chromatin structure analyses identify miRNA promoters. *Genes Dev.* 2008;22(22):3172–83. doi:10.1101/gad.1706508.

25. Corcoran DL, Pandit KV, Gordon B, Bhattacharjee A, Kaminski N, Benos PV. Features of mammalian microRNA promoters emerge from polymerase II chromatin immunoprecipitation data. *PLoS ONE*. 2009;4(4):e5279. doi:10.1371/journal.pone.0005279.
26. He L, He X, Lim LP, de Stanchina E, Xuan Z, Liang Y, et al. A microRNA component of the p53 tumour suppressor network. *Nature*. 2007;447(7148):1130–4. doi:10.1038/nature05939.
27. O'Donnell KA, Wentzel EA, Zeller KI, Dang CV, Mendell JT. c-Myc-regulated microRNAs modulate E2F1 expression. *Nature*. 2005;435(7043):839–43. doi:10.1038/nature03677.
28. Chang TC, Yu D, Lee YS, Wentzel EA, Arking DE, West KM, et al. Widespread microRNA repression by Myc contributes to tumorigenesis. *Nat Genet*. 2008;40(1):43–50. doi:10.1038/ng.2007.30.
29. Tan X, Wang S, Yang B, Zhu L, Yin B, Chao T, et al. The CREB-miR-9 negative feedback microcircuitry coordinates the migration and proliferation of glioma cells. *PLoS ONE*. 2012;7(11):e49570. doi:10.1371/journal.pone.0049570.
30. Zhang Y, Yang J, Cui X, Chen Y, Zhu VF, Hagan JP, et al. A novel epigenetic CREB-miR-373 axis mediates ZIP4-induced pancreatic cancer growth. *EMBO Mol Med*. 2013;5(9):1322–34. doi:10.1002/emmm.201302507.
31. Zhang JQ, Yao QH, Kuang YQ, Ma Y, Yang LB, Huang HD, et al. Prognostic value of coexistence of abnormal expression of microRNA-200b and cyclic adenosine monophosphate-responsive element-binding protein 1 in human Astrocytoma. *Hum Pathol*. 2014;45(10):2154–61. doi:10.1016/j.humpath.2014.01.025.
32. Son J, Lee JH, Kim HN, Ha H, Lee ZH. cAMP-response-element-binding protein positively regulates breast cancer metastasis and subsequent bone destruction. *Biochem Biophys Res Commun*. 2010;398(2):309–14. doi:10.1016/j.bbrc.2010.06.087.
33. Chhabra A, Fernando H, Watkins G, Mansel RE, Jiang WG. Expression of transcription factor CREB1 in human breast cancer and its correlation with prognosis. *Oncol Rep*. 2007;18(4):953–8.
34. Zhang M, Xu JJ, Zhou RL, Zhang QY. cAMP responsive element binding protein-1 is a transcription factor of lysosomal-associated protein transmembrane-4 beta in human breast cancer cells. *PLoS ONE*. 2013;8(2):e57520. doi:10.1371/journal.pone.0057520.
35. Resende C, Regalo G, Duraes C, Pinto MT, Wen X, Figueiredo C, et al. Interleukin-1B signalling leads to increased survival of gastric carcinoma cells through a CREB-C/EBPbeta-associated mechanism. *Gastric Cancer*. 2016;19(1):74–84. doi:10.1007/s10120-014-0448-x.
36. Wang YW, Chen X, Gao JW, Zhang H, Ma RR, Gao ZH, et al. High expression of cAMP-responsive element-binding protein 1 (CREB1) is associated with metastasis, tumor stage and poor outcome in gastric cancer. *Oncotarget*. 2015;6(12):10646–57.
37. Tan X, Wang S, Zhu L, Wu C, Yin B, Zhao J, et al. cAMP response element-binding protein promotes gliomagenesis by modulating the expression of oncogenic microRNA-23a. *Proc Natl Acad Sci U S A*. 2012;109(39):15805–10. doi:10.1073/pnas.1207787109.
38. Yu L, Guo X, Zhang P, Qi R, Li Z, Zhang S. Cyclic adenosine monophosphate-responsive element-binding protein activation predicts an unfavorable prognosis in patients with hepatocellular carcinoma. *OncoTargets Ther*. 2014;7:873–9. doi:10.2147/ott.s63594.
39. Dobroff AS, Wang H, Melnikova VO, Villares GJ, Zigler M, Huang L, et al. Silencing cAMP-response element-binding protein (CREB) identifies CYR61 as a tumor suppressor gene in melanoma. *J Biol Chem*. 2009;284(38):26194–206. doi:10.1074/jbc.M109.019836.
40. Barresi V, Branca G, Caffo M, Tuccari G. p-CREB expression in human meningiomas: correlation with angiogenesis and recurrence risk. *J Neuro-Oncol*. 2015;122(1):87–95. doi:10.1007/s11060-014-1706-9.
41. Shukla A, Bosenberg MW, MacPherson MB, Butnor KJ, Heintz NH, Pass HI, et al. Activated cAMP response element binding protein is overexpressed in human mesotheliomas and inhibits apoptosis. *Am J Pathol*. 2009;175(5):2197–206. doi:10.2353/ajpath.2009.090400.
42. Aggarwal S, Kim SW, Ryu SH, Chung WC, Koo JS. Growth suppression of lung cancer cells by targeting cyclic AMP response element-binding protein. *Cancer Res*. 2008;68(4):981–8. doi:10.1158/0008-5472.can-06-0249.
43. Seo HS, Liu DD, Bekele BN, Kim MK, Pisters K, Lippman SM, et al. Cyclic AMP response element-binding protein overexpression: a feature associated with negative prognosis in never smokers with non-small cell lung cancer. *Cancer Res*. 2008;68(15):6065–73. doi:10.1158/0008-5472.can-07-5376.
44. Linnerth NM, Greenaway JB, Petrik JJ, Moorehead RA. cAMP response element-binding protein is expressed at high levels in human ovarian adenocarcinoma and regulates ovarian tumor cell proliferation. *Int J Gynecol Cancer*. 2008;18(6):1248–57. doi:10.1111/j.1525-1438.2007.01177.x.
45. Wu D, Zhau HE, Huang WC, Iqbal S, Habib FK, Sartor O, et al. cAMP-responsive element-binding protein regulates vascular endothelial growth factor expression: implication in human prostate cancer bone metastasis. *Oncogene*. 2007;26(35):5070–7. doi:10.1038/sj.onc.1210316.
46. Kumar AP, Bhaskaran S, Ganapathy M, Crosby K, Davis MD, Kochunov P, et al. Akt/cAMP-responsive element binding protein/cyclin D1 network: a novel target for prostate cancer inhibition in transgenic adenocarcinoma of mouse prostate model mediated by nexrutine, a phellodendron amurense bark extract. *Clin Cancer Res*. 2007;13(9):2784–94. doi:10.1158/1078-0432.ccr-06-2974.
47. Sakamoto KM, Frank DA. CREB in the pathophysiology of cancer: implications for targeting transcription factors for cancer therapy. *Clin Cancer Res*. 2009;15(8):2583–7. doi:10.1158/1078-0432.ccr-08-1137.
48. Barresi V, Mondello S, Branca G, Rajan TS, Vitarelli E, Tuccari G. p-CREB expression in human gliomas: potential use in the differential diagnosis between astrocytoma and oligodendroglioma. *Hum Pathol*. 2015;46(2):231–8. doi:10.1016/j.humpath.2014.10.011.
49. Kong WQ, Bai R, Liu T, Cai CL, Liu M, Li X, et al. MicroRNA-182 targets cAMP-responsive element-binding protein 1 and suppresses cell growth in human gastric adenocarcinoma. *FEBS J*. 2012;279(7):1252–60. doi:10.1111/j.1742-4658.2012.08519.x.
50. Perry C, Sklan EH, Soreq H. CREB regulates AChE-R-induced proliferation of human glioblastoma cells. *Neoplasia (New York, NY)*. 2004;6(3):279–86. doi:10.1593/neo.3424.
51. Dai N, Zhong ZY, Cun YP, Qing Y, Chen C, Jiang P, et al. Alteration of the microRNA expression profile in human osteosarcoma cells transfected with APE1 siRNA. *Neoplasia*. 2013;60(4):384–94. doi:10.4149/neo_2013_050.
52. Polytaichou C, Iliopoulos D, Hatziaepostolou M, Kottakis F, Maroulakou I, Struhl K, et al. Akt2 regulates all Akt isoforms and promotes resistance to hypoxia through induction of miR-21 upon oxygen deprivation. *Cancer Res*. 2011;71(13):4720–31. doi:10.1158/0008-5472.can-11-0365.
53. Li QQ, Zhang L, Wan HY, Liu M, Li X, Tang H. CREB1-driven expression of miR-320a promotes mitophagy by down-regulating VDAC1 expression during serum starvation in cervical cancer cells. *Oncotarget*. 2015. doi:10.18632/oncotarget.5318.
54. Martin NT, Nakamura K, Davies R, Nahas SA, Brown C, Tunuguntla R, et al. ATM-dependent MiR-335 targets CtIP and modulates the DNA damage response. *PLoS Genet*. 2013;9(5):e1003505. doi:10.1371/journal.pgen.1003505.
55. Wendler A, Keller D, Albrecht C, Peluso JJ, Wehling M. Involvement of let-7/miR-98 microRNAs in the regulation of

- progesterone receptor membrane component 1 expression in ovarian cancer cells. *Oncol Rep.* 2011;25(1):273–9.
56. Zhao Y, Deng C, Lu W, Xiao J, Ma D, Guo M, et al. let-7 microRNAs induce tamoxifen sensitivity by downregulation of estrogen receptor alpha signaling in breast cancer. *Mol Med (Cambridge, Mass.)*. 2011;17(11–12):1233–41. doi:10.2119/molmed.2010.00225.
 57. Yang N, Kaur S, Volinia S, Greshock J, Lassus H, Hasegawa K, et al. MicroRNA microarray identifies Let-7i as a novel biomarker and therapeutic target in human epithelial ovarian cancer. *Cancer Res.* 2008;68(24):10307–14. doi:10.1158/0008-5472.can-08-1954.
 58. Tian Y, Hao S, Ye M, Zhang A, Nan Y, Wang G, et al. MicroRNAs let-7b/i suppress human glioma cell invasion and migration by targeting IKBKE directly. *Biochem Biophys Res Commun.* 2015;458(2):307–12. doi:10.1016/j.bbrc.2015.01.105.
 59. Zhang P, Ma Y, Wang F, Yang J, Liu Z, Peng J, et al. Comprehensive gene and microRNA expression profiling reveals the crucial role of hsa-let-7i and its target genes in colorectal cancer metastasis. *Mol Biol Rep.* 2012;39(2):1471–8. doi:10.1007/s11033-011-0884-1.
 60. Hur K, Toiyama Y, Schetter AJ, Okugawa Y, Harris CC, Boland CR, et al. Identification of a metastasis-specific microRNA signature in human colorectal cancer. *J Natl Cancer Inst.* 2015;107(3):dju492. doi:10.1093/jnci/dju492.
 61. Subramanian M, Francis P, Bilke S, Li XL, Hara T, Lu X, et al. A mutant p53/let-7i-axis-regulated gene network drives cell migration, invasion and metastasis. *Oncogene.* 2015;34(9):1094–104. doi:10.1038/ncr.2014.46.
 62. Wang H, Zhang W, Zuo Y, Ding M, Ke C, Yan R, et al. miR-9 promotes cell proliferation and inhibits apoptosis by targeting LASS2 in bladder cancer. *Tumour Biol.* 2015. doi:10.1007/s13277-015-3713-7.
 63. Xie D, Shang C, Zhang H, Guo Y, Tong X. Up-regulation of miR-9 target CBX7 to regulate invasion ability of bladder transitional cell carcinoma. *Med Sci Monit.* 2015;21:225–30. doi:10.12659/msm.893232.
 64. Drakaki A, Hatziaepostolou M, Polytarchou C, Vorvis C, Poultsides GA, Souglakos J, et al. Functional microRNA high throughput screening reveals miR-9 as a central regulator of liver oncogenesis by affecting the PPARA-CDH1 pathway. *BMC Cancer.* 2015;15:542. doi:10.1186/s12885-015-1562-9.
 65. Cai L, Cai X. Up-regulation of miR-9 expression predicate advanced clinicopathological features and poor prognosis in patients with hepatocellular carcinoma. *Diagn Pathol.* 2014;9:1000. doi:10.1186/s13000-014-0228-2.
 66. Higashi T, Hayashi H, Ishimoto T, Takeyama H, Kaida T, Arima K, et al. miR-9-3p plays a tumour-suppressor role by targeting TAZ (WWTR1) in hepatocellular carcinoma cells. *Br J Cancer.* 2015;113(2):252–8. doi:10.1038/bjc.2015.170.
 67. Pan X, Wang ZX, Wang R. MicroRNA-21: a novel therapeutic target in human cancer. *Cancer Biol Ther.* 2010;10(12):1224–32.
 68. Wang W, Li J, Zhu W, Gao C, Jiang R, Li W, et al. MicroRNA-21 and the clinical outcomes of various carcinomas: a systematic review and meta-analysis. *BMC Cancer.* 2014;14:819. doi:10.1186/1471-2407-14-819.
 69. Medina PP, Nolde M, Slack FJ. OncomiR addiction in an in vivo model of microRNA-21-induced pre-B-cell lymphoma. *Nature.* 2010;467(7311):86–90. doi: <http://www.nature.com/nature/journal/v467/n7311/abs/nature09284.html#supplementary-information>.
 70. Krichevsky AM, Gabriely G. miR-21: a small multi-faceted RNA. *J Cell Mol Med.* 2009;13(1):39–53. doi:10.1111/j.1582-4934.2008.00556.x.
 71. Fujita S, Ito T, Mizutani T, Minoguchi S, Yamamichi N, Sakurai K, et al. miR-21 gene expression triggered by AP-1 is sustained through a double-negative feedback mechanism. *J Mol Biol.* 2008;378(3):492–504. doi:10.1016/j.jmb.2008.03.015.
 72. Zheng H, Li W, Wang Y, Xie T, Cai Y, Wang Z, et al. miR-23a inhibits E-cadherin expression and is regulated by AP-1 and NFAT4 complex during Fas-induced EMT in gastrointestinal cancer. *Carcinogenesis.* 2014;35(1):173–83. doi:10.1093/carcin/bgt274.
 73. Ma G, Dai W, Sang A, Yang X, Gao C. Upregulation of microRNA-23a/b promotes tumor progression and confers poor prognosis in patients with gastric cancer. *Int J Clin Exp Pathol.* 2014;7(12):8833–40.
 74. Cao M, Seike M, Soeno C, Mizutani H, Kitamura K, Minegishi Y, et al. MiR-23a regulates TGF-beta-induced epithelial-mesenchymal transition by targeting E-cadherin in lung cancer cells. *Int J Oncol.* 2012;41(3):869–75. doi:10.3892/ijo.2012.1535.
 75. Cao M, Li Y, Lu H, Meng Q, Wang L, Cai L, et al. MiR-23a-mediated migration/invasion is rescued by its target, IRS-1, in non-small cell lung cancer cells. *J Cancer Res Clin Oncol.* 2014;140(10):1661–70. doi:10.1007/s00432-014-1725-0.
 76. Cai S, Chen R, Li X, Cai Y, Ye Z, Li S, et al. Downregulation of microRNA-23a suppresses prostate cancer metastasis by targeting the PAK6-LIMK1 signaling pathway. *Oncotarget.* 2015;6(6):3904–17.
 77. Macartney-Coxson DP, Hood KA, Shi HJ, Ward T, Wiles A, O'Connor R, et al. Metastatic susceptibility locus, an 8p hot-spot for tumour progression disrupted in colorectal liver metastases: 13 candidate genes examined at the DNA, mRNA and protein level. *BMC Cancer.* 2008;8:187. doi:10.1186/1471-2407-8-187.
 78. Knosel T, Schluns K, Stein U, Schwabe H, Schlag PM, Dietel M, et al. Chromosomal alterations during lymphatic and liver metastasis formation of colorectal cancer. *Neoplasia (New York, NY).* 2004;6(1):23–8.
 79. Huang A, Zhao H, Quan Y, Jin R, Feng B, Zheng M. E2A predicts prognosis of colorectal cancer patients and regulates cancer cell growth by targeting miR-320a. *PLoS ONE.* 2014;9(1):e85201. doi:10.1371/journal.pone.0085201.
 80. Salendo J, Spitzner M, Kramer F, Zhang X, Jo P, Wolff HA, et al. Identification of a microRNA expression signature for chemoradiosensitivity of colorectal cancer cells, involving miRNAs-320a, -224, -132 and let7g. *Radiother Oncol.* 2013;108(3):451–7. doi:10.1016/j.radonc.2013.06.032.
 81. Zhang Y, He X, Liu Y, Ye Y, Zhang H, He P, et al. MicroRNA-320a inhibits tumor invasion by targeting neuropilin 1 and is associated with liver metastasis in colorectal cancer. *Oncol Rep.* 2012;27(3):685–94. doi:10.3892/or.2011.1561.
 82. Sun JY, Huang Y, Li JP, Zhang X, Wang L, Meng YL, et al. MicroRNA-320a suppresses human colon cancer cell proliferation by directly targeting beta-catenin. *Biochem Biophys Res Commun.* 2012;420(4):787–92. doi:10.1016/j.bbrc.2012.03.075.
 83. Zhao H, Dong T, Zhou H, Wang L, Huang A, Feng B, et al. miR-320a suppresses colorectal cancer progression by targeting Rac1. *Carcinogenesis.* 2014;35(4):886–95. doi:10.1093/carcin/bgt378.
 84. Schepeler T, Reinert JT, Ostenfeld MS, Christensen LL, Silahtaroglu AN, Dyrskjot L, et al. Diagnostic and prognostic microRNAs in stage II colon cancer. *Cancer Res.* 2008;68(15):6416–24. doi:10.1158/0008-5472.can-07-6110.
 85. Kishan Z, Ziying L, Jing D, Gang L. MicroRNA-320a acts as a tumor suppressor by targeting BCR/ABL oncogene in chronic myeloid leukemia. *Sci Rep.* 2015;5:12460. doi:10.1038/srep12460.
 86. Qi X, Li J, Zhou C, Lv C, Tian M. MicroRNA-320a inhibits cell proliferation, migration and invasion by targeting BMI-1 in nasopharyngeal carcinoma. *FEBS Lett.* 2014;588(20):3732–8. doi:10.1016/j.febslet.2014.08.021.
 87. Yao J, Liang LH, Zhang Y, Ding J, Tian Q, Li JJ, et al. GNAI1 suppresses tumor cell migration and invasion and is post-

- transcriptionally regulated by Mir-320a/c/d in hepatocellular carcinoma. *Cancer Biol Med*. 2012;9(4):234–41. doi:10.7497/j.issn.2095-3941.2012.04.003.
88. Guo T, Feng Y, Liu Q, Yang X, Jiang T, Chen Y, et al. MicroRNA-320a suppresses in GBM patients and modulates glioma cell functions by targeting IGF-1R. *Tumour Biol*. 2014;35(11):11269–75. doi:10.1007/s13277-014-2283-4.
 89. Sun L, Liu B, Lin Z, Yao Y, Chen Y, Li Y, et al. MiR-320a acts as a prognostic factor and inhibits metastasis of salivary adenoid cystic carcinoma by targeting ITGB3. *Mol Cancer*. 2015;14:96. doi:10.1186/s12943-015-0344-y.
 90. Shang C, Zhang H, Guo Y, Hong Y, Liu Y, Xue Y. MiR-320a down-regulation mediates bladder carcinoma invasion by targeting ITGB3. *Mol Biol Rep*. 2014;41(4):2521–7. doi:10.1007/s11033-014-3110-0.
 91. Li H, Xie S, Liu M, Chen Z, Liu X, Wang L, et al. The clinical significance of downregulation of mir-124-3p, mir-146a-5p, mir-155-5p and mir-335-5p in gastric cancer tumorigenesis. *Int J Oncol*. 2014;45(1):197–208. doi:10.3892/ijo.2014.2415.
 92. Wang H, Li M, Zhang R, Wang Y, Zang W, Ma Y, et al. Effect of miR-335 upregulation on the apoptosis and invasion of lung cancer cell A549 and H1299. *Tumour Biol*. 2013;34(5):3101–9. doi:10.1007/s13277-013-0878-9.
 93. Dohi O, Yasui K, Gen Y, Takada H, Endo M, Tsuji K, et al. Epigenetic silencing of miR-335 and its host gene MEST in hepatocellular carcinoma. *Int J Oncol*. 2013;42(2):411–8. doi:10.3892/ijo.2012.1724.
 94. Yan Z, Xiong Y, Xu W, Gao J, Cheng Y, Wang Z, et al. Identification of hsa-miR-335 as a prognostic signature in gastric cancer. *PLoS ONE*. 2012;7(7):e40037. doi:10.1371/journal.pone.0040037.
 95. Lynch J, Meehan MH, Crean J, Copeland J, Stallings RL, Bray IM. Metastasis suppressor microRNA-335 targets the formin family of actin nucleators. *PLoS ONE*. 2013;8(11):e78428. doi:10.1371/journal.pone.0078428.
 96. Li Z, Li D, Zhang G, Xiong J, Jie Z, Cheng H, et al. Methylation-associated silencing of MicroRNA-335 contributes tumor cell invasion and migration by interacting with RASA1 in gastric cancer. *Am J Cancer Res*. 2014;4(6):648–62.
 97. Xiong SW, Lin TX, Xu KW, Dong W, Ling XH, Jiang FN, et al. MicroRNA-335 acts as a candidate tumor suppressor in prostate cancer. *Pathol Oncol Res*. 2013;19(3):529–37. doi:10.1007/s12253-013-9613-5.
 98. Meng Y, Zou Q, Liu T, Cai X, Huang Y, Pan J. MicroRNA-335 inhibits proliferation, cell-cycle progression, colony formation, and invasion via targeting PAX6 in breast cancer cells. *Molecular Med Rep*. 2015;11(1):379–85. doi:10.3892/mmr.2014.2684.
 99. Wang C, Jiang T. MicroRNA-335 represents an independent prognostic marker in cervical cancer. *Tumour Biol*. 2015. doi:10.1007/s13277-015-3252-2.
 100. Qiao J, Lee S, Paul P, Theiss L, Tiao J, Qiao L, et al. miR-335 and miR-363 regulation of neuroblastoma tumorigenesis and metastasis. *Surgery*. 2013;154(2):226–33. doi:10.1016/j.surg.2013.04.005.
 101. Gao L, Yang Y, Xu H, Liu R, Li D, Hong H, et al. MiR-335 functions as a tumor suppressor in pancreatic cancer by targeting OCT4. *Tumour Biol*. 2014;35(8):8309–18. doi:10.1007/s13277-014-2092-9.
 102. Gao Y, Zeng F, Wu JY, Li HY, Fan JJ, Mai L, et al. MiR-335 inhibits migration of breast cancer cells through targeting oncoprotein c-Met. *Tumour Biol*. 2015;36(4):2875–83. doi:10.1007/s13277-014-2917-6.
 103. Gong M, Ma J, Guillemette R, Zhou M, Yang Y, Yang Y, et al. miR-335 inhibits small cell lung cancer bone metastases via IGF-IR and RANKL pathways. *Mol Cancer Res*. 2014;12(1):101–10. doi:10.1158/1541-7786.mcr-13-0136.
 104. Wang K, Chen X, Zhan Y, Jiang W, Liu X, Wang X, et al. miR-335 inhibits the proliferation and invasion of clear cell renal cell carcinoma cells through direct suppression of BCL-W. *Tumour Biol*. 2015. doi:10.1007/s13277-015-3382-6.
 105. Shi L, Jiang D, Sun G, Wan Y, Zhang S, Zeng Y, et al. miR-335 promotes cell proliferation by directly targeting Rb1 in meningiomas. *J Neuro-Oncol*. 2012;110(2):155–62. doi:10.1007/s11060-012-0951-z.
 106. Cao J, Cai J, Huang D, Han Q, Yang Q, Li T, et al. miR-335 represents an invasion suppressor gene in ovarian cancer by targeting Bcl-w. *Oncol Rep*. 2013;30(2):701–6. doi:10.3892/or.2013.2482.
 107. Wang Y, Zhao W, Fu Q. miR-335 suppresses migration and invasion by targeting ROCK1 in osteosarcoma cells. *Mol Cell Biochem*. 2013;384(1–2):105–11. doi:10.1007/s11010-013-1786-4.
 108. Heidary MF, Mahmoodzadeh Hosseini H, Mehdizadeh Aghdam E, Nourani MR, Ranjbar R, Mirnejad R, et al. Overexpression of metastatic related microRNAs, Mir-335 and Mir-10b, by staphylococcal enterotoxin B in the metastatic breast cancer cell line. *Adv Pharma Bull*. 2015;5(2):255–9. doi:10.15171/apb.2015.035.
 109. Cheng Q, Cao H, Chen Z, Ma Z, Wan X, Peng R, et al. PAX6, a novel target of miR-335, inhibits cell proliferation and invasion in glioma cells. *Mol Med Rep*. 2014;10(1):399–404. doi:10.3892/mmr.2014.2150.
 110. Zhang BJ, Gong HY, Zheng F, Liu DJ, Liu HX. Up-regulation of miR-335 predicts a favorable prognosis in esophageal squamous cell carcinoma. *Int J Clin Exp Pathol*. 2014;7(9):6213–8.
 111. Voorhoeve PM, le Sage C, Schrier M, Gillis AJ, Stoop H, Nagel R, et al. A genetic screen implicates miRNA-372 and miRNA-373 as oncogenes in testicular germ cell tumors. *Cell*. 2006;124(6):1169–81. doi:10.1016/j.cell.2006.02.037.
 112. Suh MR, Lee Y, Kim JY, Kim SK, Moon SH, Lee JY, et al. Human embryonic stem cells express a unique set of microRNAs. *Dev Biol*. 2004;270(2):488–98. doi:10.1016/j.ydbio.2004.02.019.
 113. Wei F, Cao C, Xu X, Wang J. Diverse functions of miR-373 in cancer. *J Transl Med*. 2015;13:162. doi:10.1186/s12967-015-0523-z.
 114. Adi Harel S, Bossel Ben-Moshe N, Aylon Y, Bublik DR, Moskovits N, Toperoff G, et al. Reactivation of epigenetically silenced miR-512 and miR-373 sensitizes lung cancer cells to cisplatin and restricts tumor growth. *Cell Death Differ*. 2015;22(8):1328–40. doi:10.1038/cdd.2014.221.
 115. Attar M, Arefian E, Nabiuni M, Adegani FJ, Bakhtiari SH, Karimi Z, et al. MicroRNA 17–92 expressed by a transposon-based vector changes expression level of cell-cycle-related genes. *Cell Biol Int*. 2012;36(11):1005–12. doi:10.1042/cbi20110089.
 116. Pigazzi M, Manara E, Baron E, Basso G. miR-34b targets cyclic AMP-responsive element binding protein in acute myeloid leukemia. *Cancer Res*. 2009;69(6):2471–8. doi:10.1158/0008-5472.can-08-3404.
 117. Pigazzi M, Manara E, Bresolin S, Tregnago C, Beghin A, Baron E, et al. MicroRNA-34b promoter hypermethylation induces CREB overexpression and contributes to myeloid transformation. *Haematologica*. 2013;98(4):602–10. doi:10.3324/haematol.2012.070664.
 118. Chen L, Yang Q, Kong WQ, Liu T, Liu M, Li X, et al. MicroRNA-181b targets cAMP responsive element binding protein 1 in gastric adenocarcinomas. *IUBMB life*. 2012;64(7):628–35. doi:10.1002/iub.1030.
 119. Peng B, Hu S, Jun Q, Luo D, Zhang X, Zhao H, et al. MicroRNA-200b targets CREB1 and suppresses cell growth in human malignant glioma. *Mol Cell Biochem*. 2013;379(1–2):51–8. doi:10.1007/s11010-013-1626-6.

120. Noguchi S, Kumazaki M, Yasui Y, Mori T, Yamada N, Akao Y. MicroRNA-203 regulates melanosome transport and tyrosinase expression in melanoma cells by targeting kinesin superfamily protein 5b. *J Invest Dermatol*. 2014;134(2):461–9. doi:10.1038/jid.2013.310.
121. Wong KY, Liang R, So CC, Jin DY, Costello JF, Chim CS. Epigenetic silencing of MIR203 in multiple myeloma. *Br J Haematol*. 2011;154(5):569–78. doi:10.1111/j.1365-2141.2011.08782.x.
122. Noguchi S, Kumazaki M, Mori T, Baba K, Okuda M, Mizuno T, et al. Analysis of microRNA-203 function in CREB/MITF/RAB27a pathway: comparison between canine and human melanoma cells. *Vet Comp Oncol*. 2014. doi:10.1111/vco.12118.
123. Yang Z, Tsuchiya H, Zhang Y, Hartnett ME, Wang L. MicroRNA-433 inhibits liver cancer cell migration by repressing the protein expression and function of cAMP response element-binding protein. *J Biol Chem*. 2013;288(40):28893–9. doi:10.1074/jbc.M113.502682.
124. Olive V, Jiang I, He L. mir-17-92, a cluster of miRNAs in the midst of the cancer network. *Int J Biochem Cell Biol*. 2010;42(8):1348–54. doi:10.1016/j.biocel.2010.03.004.
125. He L, Thomson JM, Hemann MT, Hernando-Monge E, Mu D, Goodson S, et al. A microRNA polycistron as a potential human oncogene. *Nature*. 2005;435(7043):828–33. doi:10.1038/nature03552.
126. Jin L, Wessely O, Marcusson EG, Ivan C, Calin GA, Alahari SK. Prooncogenic factors miR-23b and miR-27b are regulated by Her2/Neu, EGF, and TNF-alpha in breast cancer. *Cancer Res*. 2013;73(9):2884–96. doi:10.1158/0008-5472.can-12-2162.
127. Chiyomaru T, Seki N, Inoguchi S, Ishihara T, Mataka H, Matsushita R, et al. Dual regulation of receptor tyrosine kinase genes EGFR and c-Met by the tumor-suppressive microRNA-23b/27b cluster in bladder cancer. *Int J Oncol*. 2015;46(2):487–96. doi:10.3892/ijo.2014.2752.
128. Goto Y, Kojima S, Nishikawa R, Enokida H, Chiyomaru T, Kinoshita T, et al. The microRNA-23b/27b/24-1 cluster is a disease progression marker and tumor suppressor in prostate cancer. *Oncotarget*. 2014;5(17):7748–59.
129. Ishteiyw RA, Ward TM, Dykxhoorn DM, Burnstein KL. The microRNA -23b/-27b cluster suppresses the metastatic phenotype of castration-resistant prostate cancer cells. *PLoS ONE*. 2012;7(12):e52106. doi:10.1371/journal.pone.0052106.
130. Jiang J, Lv X, Fan L, Huang G, Zhan Y, Wang M, et al. MicroRNA-27b suppresses growth and invasion of NSCLC cells by targeting Sp1. *Tumour Biol*. 2014;35(10):10019–23. doi:10.1007/s13277-014-2294-1.
131. Wan L, Zhang L, Fan K, Wang J. MiR-27b targets LIMK1 to inhibit growth and invasion of NSCLC cells. *Mol Cell Biochem*. 2014;390(1–2):85–91. doi:10.1007/s11010-013-1959-1.
132. Mu W, Hu C, Zhang H, Qu Z, Cen J, Qiu Z, et al. miR-27b synergizes with anticancer drugs via p53 activation and CYP1B1 suppression. *Cell Res*. 2015;25(4):477–95. doi:10.1038/cr.2015.23.
133. Zhang S, Liu F, Mao X, Huang J, Yang J, Yin X, et al. Elevation of miR-27b by HPV16 E7 inhibits PPARgamma expression and promotes proliferation and invasion in cervical carcinoma cells. *Int J Oncol*. 2015. doi:10.3892/ijo.2015.3162.
134. Chen L, Li H, Han L, Zhang K, Wang G, Wang Y, et al. Expression and function of miR-27b in human glioma. *Oncol Rep*. 2011;26(6):1617–21. doi:10.3892/or.2011.1458.
135. Wang Y, Rathinam R, Walch A, Alahari SK. ST14 (suppression of tumorigenicity 14) gene is a target for miR-27b, and the inhibitory effect of ST14 on cell growth is independent of miR-27b regulation. *J Biol Chem*. 2009;284(34):23094–106. doi:10.1074/jbc.M109.012617.
136. Zhu Y, Zhang X, Ding X, Wang H, Chen X, Zhao H, et al. miR-27 inhibits adipocyte differentiation via suppressing CREB expression. *Acta Biochim Biophys Sin*. 2014;46(7):590–6. doi:10.1093/abbs/gmu036.
137. Chien CH, Sun YM, Chang WC, Chiang-Hsieh PY, Lee TY, Tsai WC, et al. Identifying transcriptional start sites of human microRNAs based on high-throughput sequencing data. *Nucleic Acids Res*. 2011;39(21):9345–56. doi:10.1093/nar/gkr604.
138. Chang TH, Huang HY, Hsu JB, Weng SL, Horng JT, Huang HD. An enhanced computational platform for investigating the roles of regulatory RNA and for identifying functional RNA motifs. *BMC Bioinforma*. 2013;14 Suppl 2:S4. doi:10.1186/1471-2105-14-s2-s4.
139. Messegue X, Escudero R, Farre D, Nunez O, Martinez J, Alba MM. PROMO: detection of known transcription regulatory elements using species-tailored searches. *Bioinformatics (Oxford, Engl)*. 2002;18(2):333–4.
140. Fabbri M, Croce CM, Calin GA. MicroRNAs. *Cancer J (Sudbury, Mass)*. 2008;14(1):1–6. doi:10.1097/PPO.0b013e318164145e.
141. He L, He X, Lowe SW, Hannon GJ. MicroRNAs join the p53 network—another piece in the tumour-suppression puzzle. *Nat Rev Cancer*. 2007;7(11):819–22. doi:10.1038/nrc2232.
142. Hiyoshi Y, Schetter AJ, Okayama H, Inamura K, Anami K, Nguyen GH, et al. Increased microRNA-34b and -34c predominantly expressed in stromal tissues is associated with poor prognosis in human colon cancer. *PLoS ONE*. 2015;10(4):e0124899. doi:10.1371/journal.pone.0124899.
143. Toyota M, Suzuki H, Sasaki Y, Maruyama R, Imai K, Shinomura Y, et al. Epigenetic silencing of microRNA-34b/c and B-cell translocation gene 4 is associated with CpG island methylation in colorectal cancer. *Cancer Res*. 2008;68(11):4123–32. doi:10.1158/0008-5472.can-08-0325.
144. Suzuki H, Yamamoto E, Nojima M, Kai M, Yamano HO, Yoshikawa K, et al. Methylation-associated silencing of microRNA-34b/c in gastric cancer and its involvement in an epigenetic field defect. *Carcinogenesis*. 2010;31(12):2066–73. doi:10.1093/carcin/bgq203.
145. Forno I, Ferrero S, Russo MV, Gazzano G, Giangiobbe S, Montanari E, et al. Deregulation of MiR-34b/Sox2 predicts prostate cancer progression. *PLoS ONE*. 2015;10(6):e0130060. doi:10.1371/journal.pone.0130060.
146. Majid S, Dar AA, Saini S, Shahryari V, Arora S, Zaman MS, et al. miRNA-34b inhibits prostate cancer through demethylation, active chromatin modifications, and AKT pathways. *Clin Cancer Res*. 2013;19(1):73–84. doi:10.1158/1078-0432.ccr-12-2952.
147. Xie K, Liu J, Chen J, Dong J, Ma H, Liu Y, et al. Methylation-associated silencing of microRNA-34b in hepatocellular carcinoma cancer. *Gene*. 2014;543(1):101–7. doi:10.1016/j.gene.2014.03.059.
148. Wang LG, Ni Y, Su BH, Mu XR, Shen HC, Du JJ. MicroRNA-34b functions as a tumor suppressor and acts as a nodal point in the feedback loop with Met. *Int J Oncol*. 2013;42(3):957–62. doi:10.3892/ijo.2013.1767.
149. Liu C, Cheng H, Shi S, Cui X, Yang J, Chen L, et al. MicroRNA-34b inhibits pancreatic cancer metastasis through repressing Smad3. *Curr Mol Med*. 2013;13(4):467–78.
150. Liu X, Feng J, Tang L, Liao L, Xu Q, Zhu S. The regulation and function of miR-21-FOXO3a-miR-34b/c signaling in breast cancer. *Int J Mol Sci*. 2015;16(2):3148–62. doi:10.3390/ijms16023148.
151. Corney DC, Hwang CI, Matoso A, Vogt M, Flesken-Nikitin A, Godwin AK, et al. Frequent downregulation of miR-34 family in human ovarian cancers. *Clin Cancer Res*. 2010;16(4):1119–28. doi:10.1158/1078-0432.ccr-09-2642.
152. Liu G, Min H, Yue S, Chen CZ. Pre-miRNA loop nucleotides control the distinct activities of mir-181a-1 and mir-181c in early

- T cell development. PLoS ONE. 2008;3(10):e3592. doi:[10.1371/journal.pone.0003592](https://doi.org/10.1371/journal.pone.0003592).
153. Liu J, Shi W, Wu C, Ju J, Jiang J. miR-181b as a key regulator of the oncogenic process and its clinical implications in cancer (review). *BioMed Rep*. 2014;2(1):7–11. doi:[10.3892/br.2013.199](https://doi.org/10.3892/br.2013.199).
 154. Iliopoulos D, Jaeger SA, Hirsch HA, Bulyk ML, Struhl K. STAT3 activation of miR-21 and miR-181b-1 via PTEN and CYLD are part of the epigenetic switch linking inflammation to cancer. *Mol Cell*. 2010;39(4):493–506. doi:[10.1016/j.molcel.2010.07.023](https://doi.org/10.1016/j.molcel.2010.07.023).
 155. Sun X, Icli B, Wara AK, Belkin N, He S, Kobzik L, et al. MicroRNA-181b regulates NF-kappaB-mediated vascular inflammation. *J Clin Invest*. 2012;122(6):1973–90. doi:[10.1172/jci61495](https://doi.org/10.1172/jci61495).
 156. Xia Y, Gao Y. MicroRNA-181b promotes ovarian cancer cell growth and invasion by targeting LATS2. *Biochem Biophys Res Commun*. 2014;447(3):446–51. doi:[10.1016/j.bbrc.2014.04.027](https://doi.org/10.1016/j.bbrc.2014.04.027).
 157. Panarelli NC, Chen YT, Zhou XK, Kitabayashi N, Yantiss RK. MicroRNA expression aids the preoperative diagnosis of pancreatic ductal adenocarcinoma. *Pancreas*. 2012;41(5):685–90. doi:[10.1097/MPA.0b013e318243a905](https://doi.org/10.1097/MPA.0b013e318243a905).
 158. Nurul-Syakima AM, Yoke-Kqueen C, Sabariah AR, Shiran MS, Singh A, Learn-Han L. Differential microRNA expression and identification of putative miRNA targets and pathways in head and neck cancers. *Int J Mol Med*. 2011;28(3):327–36. doi:[10.3892/ijmm.2011.714](https://doi.org/10.3892/ijmm.2011.714).
 159. Ratert N, Meyer HA, Jung M, Mollenkopf HJ, Wagner I, Miller K, et al. Reference miRNAs for miRNAome analysis of urothelial carcinomas. PLoS ONE. 2012;7(6):e39309. doi:[10.1371/journal.pone.0039309](https://doi.org/10.1371/journal.pone.0039309).
 160. Li D, Jian W, Wei C, Song H, Gu Y, Luo Y, et al. Down-regulation of miR-181b promotes apoptosis by targeting CYLD in thyroid papillary cancer. *Int J Clin Exp Pathol*. 2014;7(11):7672–80.
 161. Sun YC, Wang J, Guo CC, Sai K, Wang J, Chen FR, et al. MiR-181b sensitizes glioma cells to teniposide by targeting MDM2. *BMC Cancer*. 2014;14:611. doi:[10.1186/1471-2407-14-611](https://doi.org/10.1186/1471-2407-14-611).
 162. Shi ZM, Wang XF, Qian X, Tao T, Wang L, Chen QD, et al. MiRNA-181b suppresses IGF-1R and functions as a tumor suppressor gene in gliomas. *RNA (New York, NY)*. 2013;19(4):552–60. doi:[10.1261/ma.035972.112](https://doi.org/10.1261/ma.035972.112).
 163. Zhi F, Wang Q, Deng D, Shao N, Wang R, Xue L, et al. MiR-181b-5p downregulates NOVA1 to suppress proliferation, migration and invasion and promote apoptosis in astrocytoma. PLoS ONE. 2014;9(10):e109124. doi:[10.1371/journal.pone.0109124](https://doi.org/10.1371/journal.pone.0109124).
 164. Lu F, Zhang J, Ji M, Li P, Du Y, Wang H, et al. miR-181b increases drug sensitivity in acute myeloid leukemia via targeting HMGB1 and Mcl-1. *Int J Oncol*. 2014;45(1):383–92. doi:[10.3892/ijo.2014.2390](https://doi.org/10.3892/ijo.2014.2390).
 165. Schaefer A, Jung M, Mollenkopf HJ, Wagner I, Stephan C, Jentzmik F, et al. Diagnostic and prognostic implications of microRNA profiling in prostate carcinoma. *Int J Cancer*. 2010;126(5):1166–76. doi:[10.1002/ijc.24827](https://doi.org/10.1002/ijc.24827).
 166. Guo JX, Tao QS, Lou PR, Chen XC, Chen J, Yuan GB. miR-181b as a potential molecular target for anticancer therapy of gastric neoplasms. *Asian Pac J Cancer Prev*. 2012;13(5):2263–7.
 167. Tang X, Zheng D, Hu P, Zeng Z, Li M, Tucker L, et al. Glycogen synthase kinase 3 beta inhibits microRNA-183-96-182 cluster via the beta-catenin/TCF/LEF-1 pathway in gastric cancer cells. *Nucleic Acids Res*. 2014;42(5):2988–98. doi:[10.1093/nar/gkt1275](https://doi.org/10.1093/nar/gkt1275).
 168. Zhang QH, Sun HM, Zheng RZ, Li YC, Zhang Q, Cheng P, et al. Meta-analysis of microRNA-183 family expression in human cancer studies comparing cancer tissues with noncancerous tissues. *Gene*. 2013;527(1):26–32. doi:[10.1016/j.gene.2013.06.006](https://doi.org/10.1016/j.gene.2013.06.006).
 169. Zhu YJ, Xu B, Xia W. Hsa-mir-182 downregulates RASA1 and suppresses lung squamous cell carcinoma cell proliferation. *Clin Lab*. 2014;60(1):155–9.
 170. Li Y, Zhang D, Wang X, Yao X, Ye C, Zhang S, et al. Hypoxia-inducible miR-182 enhances HIF1alpha signaling via targeting PHD2 and FIH1 in prostate cancer. *Sci Rep*. 2015;5:12495. doi:[10.1038/srep12495](https://doi.org/10.1038/srep12495).
 171. Du C, Weng X, Hu W, Lv Z, Xiao H, Ding C, et al. Hypoxia-inducible MiR-182 promotes angiogenesis by targeting RASA1 in hepatocellular carcinoma. *J Exp Clin Cancer Res*. 2015;34(1):67. doi:[10.1186/s13046-015-0182-1](https://doi.org/10.1186/s13046-015-0182-1).
 172. Sachdeva M, Mito JK, Lee CL, Zhang M, Li Z, Dodd RD, et al. MicroRNA-182 drives metastasis of primary sarcomas by targeting multiple genes. *J Clin Invest*. 2014;124(10):4305–19. doi:[10.1172/jci77116](https://doi.org/10.1172/jci77116).
 173. Ning FL, Wang F, Li ML, Yu ZS, Hao YZ, Chen SS. MicroRNA-182 modulates chemosensitivity of human non-small cell lung cancer to cisplatin by targeting PDCD4. *Diagn Pathol*. 2014;9:143. doi:[10.1186/1746-1596-9-143](https://doi.org/10.1186/1746-1596-9-143).
 174. Kouri FM, Hurley LA, Daniel WL, Day ES, Hua Y, Hao L, et al. miR-182 integrates apoptosis, growth, and differentiation programs in glioblastoma. *Genes Dev*. 2015;29(7):732–45. doi:[10.1101/gad.257394.114](https://doi.org/10.1101/gad.257394.114).
 175. Zhang Y, Wang X, Wang Z, Tang H, Fan H, Guo Q. miR-182 promotes cell growth and invasion by targeting forkhead box F2 transcription factor in colorectal cancer. *Oncol Rep*. 2015;33(5):2592–8. doi:[10.3892/or.2015.3833](https://doi.org/10.3892/or.2015.3833).
 176. Zhu H, Fang J, Zhang J, Zhao Z, Liu L, Wang J, et al. miR-182 targets CHL1 and controls tumor growth and invasion in papillary thyroid carcinoma. *Biochem Biophys Res Commun*. 2014;450(1):857–62. doi:[10.1016/j.bbrc.2014.06.073](https://doi.org/10.1016/j.bbrc.2014.06.073).
 177. Wang C, Ren R, Hu H, Tan C, Han M, Wang X, et al. MiR-182 is up-regulated and targeting Cebpa in hepatocellular carcinoma. *Chin J Cancer Res*. 2014;26(1):17–29. doi:[10.3978/j.issn.1000-9604.2014.01.01](https://doi.org/10.3978/j.issn.1000-9604.2014.01.01).
 178. Tang L, Chen F, Pang EJ, Zhang ZQ, Jin BW, Dong WF. MicroRNA-182 inhibits proliferation through targeting oncogenic ANUBL1 in gastric cancer. *Oncol Rep*. 2015;33(4):1707–16. doi:[10.3892/or.2015.3798](https://doi.org/10.3892/or.2015.3798).
 179. Li X, Luo F, Li Q, Xu M, Feng D, Zhang G, et al. Identification of new aberrantly expressed miRNAs in intestinal-type gastric cancer and its clinical significance. *Oncol Rep*. 2011;26(6):1431–9. doi:[10.3892/or.2011.1437](https://doi.org/10.3892/or.2011.1437).
 180. Feng B, Wang R, Chen LB. Review of miR-200b and cancer chemosensitivity. *Biomed Pharmacother*. 2012;66(6):397–402. doi:[10.1016/j.biopha.2012.06.002](https://doi.org/10.1016/j.biopha.2012.06.002).
 181. Mongroo PS, Rustgi AK. The role of the miR-200 family in epithelial-mesenchymal transition. *Cancer Biol Ther*. 2010;10(3):219–22.
 182. Gregory PA, Bert AG, Paterson EL, Barry SC, Tsykin A, Farshid G, et al. The miR-200 family and miR-205 regulate epithelial to mesenchymal transition by targeting ZEB1 and SIP1. *Nat Cell Biol*. 2008;10(5):593–601. doi:[10.1038/ncb1722](https://doi.org/10.1038/ncb1722).
 183. Bracken CP, Gregory PA, Kolesnikoff N, Bert AG, Wang J, Shannon MF, et al. A double-negative feedback loop between ZEB1-SIP1 and the microRNA-200 family regulates epithelial-mesenchymal transition. *Cancer Res*. 2008;68(19):7846–54. doi:[10.1158/0008-5472.can-08-1942](https://doi.org/10.1158/0008-5472.can-08-1942).
 184. Wu H, Wang G, Wang Z, An S, Ye P, Luo S. A negative feedback loop between miR-200b and the NF-kappaB pathway via IKBKB/IKK-beta in breast cancer cells. *FEBS J*. 2015. doi:[10.1111/febs.13543](https://doi.org/10.1111/febs.13543).
 185. Imaoka H, Toiyama Y, Okigami M, Yasuda H, Saigusa S, Ohi M, et al. Circulating microRNA-203 predicts metastases, early recurrence, and poor prognosis in human gastric cancer. *Gastric Cancer*. 2015. doi:[10.1007/s10120-015-0521-0](https://doi.org/10.1007/s10120-015-0521-0).
 186. Zhou X, Xu G, Yin C, Jin W, Zhang G. Down-regulation of miR-203 induced by helicobacter pylori infection promotes the

- proliferation and invasion of gastric cancer by targeting CASK. *Oncotarget*. 2014;5(22):11631–40.
187. Siu MK, Abou-Kheir W, Yin JJ, Chang YS, Barrett B, Suau F, et al. Loss of EGFR signaling regulated miR-203 promotes prostate cancer bone metastasis and tyrosine kinase inhibitors resistance. *Oncotarget*. 2014;5(11):3770–84.
 188. Liao H, Bai Y, Qiu S, Zheng L, Huang L, Liu T, et al. MiR-203 downregulation is responsible for chemoresistance in human glioblastoma by promoting epithelial-mesenchymal transition via SNAI2. *Oncotarget*. 2015;6(11):8914–28.
 189. Shi Y, Tan YJ, Zeng DZ, Qian F, Yu PW. miR-203 suppression in gastric carcinoma promotes slug-mediated cancer metastasis. *Tumour Biol*. 2015. doi:10.1007/s13277-015-3765-8.
 190. Chen T, Xu C, Chen J, Ding C, Xu Z, Li C, et al. MicroRNA-203 inhibits cellular proliferation and invasion by targeting Bmi1 in non-small cell lung cancer. *Oncol Lett*. 2015;9(6):2639–46. doi:10.3892/ol.2015.3080.
 191. Mine M, Yamaguchi K, Sugiura T, Chigita S, Yoshihama N, Yoshihama R, et al. miR-203 inhibits frizzled-2 expression via CD82/KAI1 expression in human lung carcinoma cells. *PLoS ONE*. 2015;10(7):e0131350. doi:10.1371/journal.pone.0131350.
 192. Wang N, Liang H, Zhou Y, Wang C, Zhang S, Pan Y, et al. miR-203 suppresses the proliferation and migration and promotes the apoptosis of lung cancer cells by targeting SRC. *PLoS ONE*. 2014;9(8):e105570. doi:10.1371/journal.pone.0105570.
 193. Li T, Gao F, Zhang XP. miR-203 enhances chemosensitivity to 5-fluorouracil by targeting thymidylate synthase in colorectal cancer. *Oncol Rep*. 2015;33(2):607–14. doi:10.3892/or.2014.3646.
 194. Li Z, Du L, Dong Z, Yang Y, Zhang X, Wang L, et al. MiR-203 suppresses ZNF217 upregulation in colorectal cancer and its oncogenicity. *PLoS ONE*. 2015;10(1):e0116170. doi:10.1371/journal.pone.0116170.
 195. Weiner-Gorzel K, Dempsey E, Milewska M, McGoldrick A, Toh V, Walsh A, et al. Overexpression of the microRNA miR-433 promotes resistance to paclitaxel through the induction of cellular senescence in ovarian cancer cells. *Cancer Med*. 2015;4(5):745–58. doi:10.1002/cam4.409.
 196. Furlong F, Fitzpatrick P, O'Toole S, Phelan S, McGrogan B, Maguire A, et al. Low MAD2 expression levels associate with reduced progression-free survival in patients with high-grade serous epithelial ovarian cancer. *J Pathol*. 2012;226(5):746–55. doi:10.1002/path.3035.
 197. Ueda T, Volinia S, Okumura H, Shimizu M, Taccioli C, Rossi S, et al. Relation between microRNA expression and progression and prognosis of gastric cancer: a microRNA expression analysis. *Lancet Oncol*. 2010;11(2):136–46. doi:10.1016/s1470-2045(09)70343-2.
 198. Lin X, Rice KL, Buzzai M, Hexner E, Costa FF, Kilpivaara O, et al. miR-433 is aberrantly expressed in myeloproliferative neoplasms and suppresses hematopoietic cell growth and differentiation. *Leukemia*. 2013;27(2):344–52. doi:10.1038/leu.2012.224.
 199. Del Vescovo V, Meier T, Inga A, Denti MA, Borlak J. A cross-platform comparison of affymetrix and agilent microarrays reveals discordant miRNA expression in lung tumors of c-Raf transgenic mice. *PLoS ONE*. 2013;8(11):e78870. doi:10.1371/journal.pone.0078870.
 200. Luo H, Zhang H, Zhang Z, Zhang X, Ning B, Guo J, et al. Down-regulated miR-9 and miR-433 in human gastric carcinoma. *J Exp Clin Cancer Res*. 2009;28:82. doi:10.1186/1756-9966-28-82.
 201. Yang O, Huang J, Lin S. Regulatory effects of miRNA on gastric cancer cells. *Oncol Lett*. 2014;8(2):651–6. doi:10.3892/ol.2014.2232.
 202. Guo LH, Li H, Wang F, Yu J, He JS. The tumor suppressor roles of miR-433 and miR-127 in gastric cancer. *Int J Mol Sci*. 2013;14(7):14171–84. doi:10.3390/ijms140714171.
 203. Yan Z, Shah PK, Amin SB, Samur MK, Huang N, Wang X, et al. Integrative analysis of gene and miRNA expression profiles with transcription factor-miRNA feed-forward loops identifies regulators in human cancers. *Nucleic Acids Res*. 2012;40(17):e135. doi:10.1093/nar/gks395.
 204. Lee YJ, Kim V, Muth DC, Witwer KW. Validated microRNA target databases: an evaluation. *Drug Dev Res*. 2015;76(7):389–96. doi:10.1002/ddr.21278.
 205. Dweep H, Gretz N. miRWalk2.0: a comprehensive atlas of microRNA-target interactions. *Nat Methods*. 2015;12(8):697. doi:10.1038/nmeth.3485.