



**SCIENTIFIC COMMITTEE
TWENTY-FIRST REGULAR SESSION**

Nuku'alofa, Tonga
13-21 August 2025

2025 Stock Assessment of Striped Marlin in the Southwest Pacific Ocean: Part II Data-moderate approach

**WCPFC-SC21-2025/SA-WP-07
21 July 2025**

N. Ducharme-Barth¹, C. Castillo-Jordán², F. Carvalho¹, P. Hamer²

¹NOAA Fisheries, Pacific Islands Fisheries Science Center, 1845 Wasp Boulevard, Bldg. 176, Honolulu HI 96818

²Oceanic Fisheries Programme of the Pacific Community

Table of contents

1	Executive summary	3
2	Introduction	4
3	Methods	5
3.1	Modelling framework	5
3.2	Input data	6
3.3	Population dynamics	6
3.4	Observation model	7
3.4.1	Catch treatment	7
3.5	Prior development	8
3.5.1	Biological simulation framework	8
3.5.2	Fishing data simulation framework	10
3.5.3	Other priors	11
3.5.4	Prior pushforward	11
3.6	Description of one-off sensitivities	12
3.6.1	Index	12
3.6.2	Catch observation error σ_C	12
3.6.3	Early data	13
3.6.4	Scaling priors	13
3.6.5	Catch observation model	13
3.6.6	Temporal resolution of catchability deviates	13
3.6.7	Catch reporting	14
3.6.8	Process error σ_P	14
3.6.9	Start year	14
3.6.10	Inclusion of effort deviates	15
3.6.11	Direct estimation of fishing mortality	15
3.6.12	Alternative R_{Max} prior	15
3.7	Model Implementation	15
3.8	Model diagnostics	16
3.8.1	Convergence diagnostics	16
3.8.2	Leave-one-out cross-validation diagnostics	16
3.8.3	Data fits	17
3.8.4	Posterior predictive checks	17
3.8.5	Retrospective analysis	17
3.8.6	Hindcast cross-validation	18
3.8.7	Model-free hindcasts	18
3.9	Forecasts	18
4	Results	18
4.1	Diagnostic case	18
4.1.1	Model convergence and diagnostics	18
4.1.2	Model fit to data	19
4.1.3	Model validation	19
4.1.4	Parameter estimates	19

4.1.5	Model-free hindcasts	20
4.2	One-off sensitivities	21
4.2.1	Index	21
4.2.2	Catch observation error σ_C	22
4.2.3	Early data	22
4.2.4	Scaling priors	22
4.2.5	Catch observation model	23
4.2.6	Temporal resolution of catchability deviates	23
4.2.7	Catch reporting	23
4.2.8	Process error σ_P	24
4.2.9	Start year	24
4.2.10	Inclusion of effort deviates	24
4.2.11	Direct estimation of fishing mortality	24
4.2.12	Alternative R_{Max} prior	25
4.3	Model ensemble	25
4.3.1	Ensemble diagnostics and fit	25
4.3.2	Population dynamics and trends	26
4.3.3	Stock status	27
4.3.4	Future projections	27
5	Discussion	27
5.1	General remarks	27
5.2	Challenges, limitations & key uncertainties	27
5.3	Future stock assessment modeling considerations	28
6	Acknowledgements	28
7	Declaration of generative AI use	28
8	References	29
9	Tables	32
10	Figures	42
11	Technical Annex	89
11.1	Equilibrium Age-Structured Population Dynamics	89
11.1.1	Survival	89
11.1.2	Fecundity	89
11.1.3	Stock-Recruitment Relationship	89

1 Executive summary

2 Introduction

Assessments of Southwest Pacific Ocean (SWPO) striped marlin (*Kajikia audax*) have been challenging. As documented in the joint NOAA-SPC modeling meeting report (Ducharme-Barth et al., 2025), key issues identified by WCPFC SC20 included poor fits to size composition data and relative abundance indices, conflicts between different data sources, and difficulties in estimating model initial conditions.

Discussions at the 2025 SPC Pre-Assessment Workshop highlighted additional challenges related to the estimation of total population scale. The low biomass scale indicated by previous versions of the assessment appeared largely driven by the size composition from multiple fisheries. While estimation of low biomass scale is not inherently problematic, the resulting high ratio of fishing mortality (F) to natural mortality (M) that is maintained over multiple decades in the face of relative stability in catches and catch-per-unit-effort (CPUE) was noted as suspicious and flagged for further investigation.

The complexity of integrated age-structured models, while providing detailed population dynamics representation, can become problematic when fundamental data conflicts exist or when key biological parameters are uncertain. For SWPO striped marlin, these challenges are compounded by spatiotemporal heterogeneity in fleet coverage, potential non-representativeness in mixed-fleet composition data, and limitations in age data from opportunistic sampling. Changes to productivity assumptions (growth, natural mortality, maturity, and/or steepness) or selectivity patterns will impact biomass scaling through predictions of expected size composition data.

Scaling back model complexity to a data-moderate approach, like Bayesian surplus production models (BSPMs), offers analysts a simplified yet robust alternative for stock assessment when data limitations or conflicts present challenges for more complex models. These models have proven effective in recent pelagic fish stock assessments (Winker et al., 2018), and are routinely applied for WCPFC shark assessments (ISC, 2024; Neubauer et al., 2019, 2024). They facilitate a more tractable exploration of model assumptions by distilling productivity and fishing assumptions into a restricted parameter subset. The BSPM framework allows for explicit incorporation of parameter uncertainty through informative priors while maintaining computational efficiency and interpretability.

One advantage of the Bayesian approach is the use of prior information to propagate uncertainty in model assumptions. Biological simulation approaches can provide robust methods for parameterizing key population dynamics parameters (ISC, 2024; Pardo et al., 2016). For species with complex life histories like striped marlin, these simulation-based priors can incorporate uncertainty in growth, maturity, natural mortality, and reproductive parameters to derive realistic distributions for maximum intrinsic rate of increase and carrying capacity. Additionally, prior pushforward checks (Kim & Neubauer, 2025; Monnahan, 2024) can be used to further refine parameter distributions by identifying parameter combinations that produce clearly improbable outcomes.

This analysis presents a data-moderate approach for SWPO striped marlin that addresses some of the key technical challenges identified in previous assessments while providing a robust framework for stock status evaluation. The model incorporates biological uncertainty through simulation-based priors and includes approaches to address catch uncertainty and population depletion. It is important to note that this analysis complements the 2025 revised stock assessment report (Castillo-Jordan et al., 2025). While the relative simplicity of data-moderate approaches offers some advantages relative to more data-intensive approaches

like fully-integrated age-structured stock assessment models, particularly when uncertainty in data or key assumptions is high, this comes at the cost of simplifying the model fishery and population dynamics. Over-simplification of key age-structured processes, or lack of spatiotemporal specificity could result in biases in data-moderate approaches. Taking multiple modeling approaches into account can give managers a holistic view on the likely status and trajectory of the stock.

3 Methods

3.1 Modelling framework

To address some of the issues raised in Section 2, a series of Bayesian state-space surplus production models (BSPMs) spanning the period 1952-2022 were developed following the Fletcher-Schaefer production model framework (Edwards, 2024; Winker et al., 2019). Development of the BSPM followed the approach of (ISC, 2024; Neubauer et al., 2019) and incorporated recent best practices for surplus production models (Kokkalis et al., 2024) and Bayesian workflows for stock assessment (Monnahan, 2024) as guides for model development, analysis, and presentation.

The models were implemented in the Stan probabilistic programming language (Team, 2024b) to take advantage of enhanced convergence diagnostics, greater efficiency in posterior sampling through the no-U-turn (NUTS) Hamiltonian Monte Carlo (HMC) algorithm (Betancourt & Girolami, 2013), and greater flexibility with model configuration and prior specification. Implementation in R using the *rstan* package (Team, 2024a) provides connection to an ecosystem of R packages for model diagnostics and validation: *bayesplot* (Gabry & Mahr, 2024) for posterior visualization and *loo* (Vehtari et al., 2024).

In their baseline configuration (diagnostic case), the models begin from an assumed unfished state in 1952 and incorporate process error in population dynamics, observation error in the relative abundance index and catch time series, and uncertainty in biological parameters. Population scale or carrying capacity $\log(K)$ is given in the units of the catch which is total numbers. The models fit to a single time series of relative abundance (e.g., standardized CPUE index) at a time. Rather than subtracting the catch directly, multiplicative survival after fishing mortality was applied. Fishing mortality was estimated internally by fitting to a time series of observed catches. Computationally, this was necessary to avoid placing hard constraints on the posterior parameter space which can cause convergence issues for the HMC sampling algorithm. Additionally, rather than directly estimating the independent fishing mortality values required to produce the catch, fishing mortality was derived from an estimated time-varying catchability trend and a time series of observed longline fishing effort. The primary rationale for this modelling choice being that it is easier to develop a defensible prior for catchability rather than an arbitrary prior for fishing mortality, given that previous work (ISC, 2024) showed a strong correlation between estimated population scale and the fishing mortality prior. Unless otherwise mentioned, all models used the same prior parameterizations (Table 2) based on the biological simulation filtering described in Section 3.5.

Using the diagnostic model¹ as a starting point, a number of one-off sensitivity analyses were conducted to explore key model assumptions or data questions. These are described in Section 3.6 and listed in Table 3. Based on the results of the sensitivity analyses a model ensemble was developed for the provision of stock

¹The Stan code for the diagnostic model `bspm_estqsimple_softdep_fullmvprior_x0_sttgamma_flexsigmaC_OPT` is publicly available on [GitHub](#).

status and management reference points. Full model development code and input data are available online at the following GitHub repository: <https://github.com/N-DucharmeBarth-NOAA/2025-swpo-mls-bspm>.

3.2 Input data

The BSPMs were fitted to catch and standardized CPUE data for SWPO striped marlin. Effort was used as an input to drive the time series of estimated fishing mortality via the estimated catchability trend.

Annual catch data (in numbers of individuals) spanning 1952-2022 were compiled from the 2024 assessment (Castillo-Jordan et al., 2024) sources and aggregated into total removals by year. The catch series (Figure 1) shows initially low removals in 1952-1953, followed by a high peak in 1954. Overall, however, catches have been relatively stable though catches in recent decades have shown a slight decline.

Annual longline effort (in numbers of hooks) matching the SWPO assessment domain, as specified in Castillo-Jordan et al. (2024), was extracted from publicly available WCPFC data. Longline effort in the assessment domain (Figure 1) increased gradually from zero in the early 1950s to ~100 million hooks per year in the early 1970s where it stabilized between 100-150 million hooks per year through the 1990s. Beginning in 2000 effort increased substantially to ~350 million hooks per year in 2003, and has fluctuated around that level for the last 20 years.

A number of indices were investigated either as one-off sensitivities or included in the structural uncertainty grid (Figure 2). These included the composite distant water fishing nation (DWFN) longline standardized CPUE index (1988-2022) from the diagnostic case of the 2024 assessment (Castillo-Jordan et al., 2024), an Australian longline index (Tremblay-Boyer & Williams, 2024), a New Zealand recreational sportfish index (Holdsworth, 2023), and three different observer data based indices (Neubauer et al., 2025). The Pacific Islands observer program indices contained information from New Caledonia, Fiji, Tonga, and French Polynesia observer programs. Three alternative indices were constructed either by including data from all programs together, including data from just the western programs (New Caledonia, Fiji, and Tonga) or including data just from French Polynesia. The DWFN, Australia and observer without French Polynesia indices show recent increases; the New Zealand and observer with only French Polynesia indices show recent declines; and the all observer program index shows no trend.

3.3 Population dynamics

The population dynamics follow a hybrid Fletcher-Schaefer surplus production model (Edwards, 2024; ISC, 2024) with state-space formulation where population depletion x_t (relative to carrying capacity K) evolves according to:

$$x_t = (x_{t-1} + \text{surplus production}_{t-1}) \times \epsilon_t \times \exp(-F_{t-1})$$

where surplus production is defined as:

$$\text{surplus production}_{t-1} = \begin{cases} R_{Max} \cdot x_{t-1} \left(1 - \frac{x_{t-1}}{h}\right), & x_{t-1} \leq D_{MSY} \\ \gamma \cdot m \cdot x_{t-1} \left(1 - x_{t-1}^{n-1}\right), & x_{t-1} > D_{MSY} \end{cases}$$

Here, R_{Max} is the maximum intrinsic rate of increase, n is the shape parameter, F_{t-1} is the fishing mortality in year $t - 1$, and ϵ_t represents multiplicative process error with $\epsilon_t = \exp(\delta_t - \sigma_P^2/2)$ and $\delta_t \sim N(0, \sigma_P)$. The intermediate parameters are: $D_{MSY} = (1/n)^{1/(n-1)}$, $h = 2D_{MSY}$, $m = R_{Max} \cdot h/4$, and $\gamma = n^{n/(n-1)}/(n-1)$.

Fishing mortality F_t is in turn defined as a function of fishing effort and time-varying catchability:

$$F_t = q_{eff} \times \exp(\delta_{q,t} - \sigma_{qdev}^2/2) \times E_t$$

where q_{eff} is the base catchability coefficient, E_t is the fishing effort² in year t , and $\delta_{q,t}$ represents temporal variation in catchability effectiveness.

The catchability deviations $\delta_{q,t}$ follow a structured temporal process organized into discrete periods of length n_{step} years. In the diagnostic model $n_{step} = 1$ which allows for annual catchability deviates across 70³ periods p . Catchability remains constant within periods p , and evolves between periods according to an AR(1) process:

$$\delta_{q,p} = \rho \times \delta_{q,p-1} + \epsilon_{q,p} \times \sigma_{qdev} \times \sqrt{1 - \rho^2}$$

where ρ is the autocorrelation parameter, σ_{qdev} is the standard deviation of catchability innovations, and $\epsilon_{q,p} \sim N(0, 1)$. The initial period is constrained such that $\delta_{q,1} = 0$. The term $\sqrt{1 - \rho^2}$ normalizes the variance of $\delta_{q,p}$ to remain stationary at σ_{qdev}^2 regardless of the value of ρ .

For each year t , the corresponding period assignment is:

$$\text{period}(t) = \min \left(\left\lfloor \frac{t-1}{n_{step}} \right\rfloor + 1, n_{periods} \right)$$

and the catchability deviation for year t is assigned as $\delta_{q,t} = \delta_{q,\text{period}(t)}$. This fishing mortality parametrization is a simplification of the *catch-errors* formulation in MULTIFAN-CL (Fournier et al., 1998) and allows catchability to vary smoothly over time while maintaining temporal structure and avoiding overfitting to annual fluctuations in the effort-catch relationship.

Model initial conditions are defined for $t = 1$ as $x_t = x_0 \times \epsilon_t$ where x_0 is the estimated depletion relative to unfished at the start of the model period. The prior specified for x_0 in the diagnostic case assumes an unfished initial condition with some uncertainty (Table 2).

3.4 Observation model

The model fits to a standardized CPUE index using a lognormal likelihood:

$$I_t \sim \text{Lognormal}(\log(q \times x_t) - \frac{\sigma_{O,t}^2}{2}, \sigma_{O,t})$$

²When input in the model the raw input effort, defined as *hundred hooks* in the WCPFC database is divided by [1e6](#) to re-scale catchability so that it is numerically on the same scale as other estimated parameters. This improves convergence of the HMC sampling of the posterior distribution.

³There are 71 total periods in the model period 1952-2022 and fishing mortality is not estimated in the terminal year.

where catchability q is analytically derived assuming an uninformative uniform prior, and total observation error $\sigma_{O,t} = \sigma_{O,input} + \sigma_{O,add}$ combines fixed input uncertainty with an estimated additional error component.

3.4.1 Catch treatment

Population removals (catch) were calculated in each time step as:

$$\text{Removals}_{t-1} = (x_{t-1} + \text{surplus production}_{t-1}) \times \epsilon_t \times (1 - \exp(-F_{t-1})) \times \exp(\log K)$$

As described in Section 3.3, instantaneous fishing mortality F_t was transformed to a discrete proportion of the population that survived fishing, survivorship, through the exponential decay function $\exp(-F_{t-1})$. Removals are simply calculated as the complementary proportion $(1 - \exp(-F_{t-1}))$ of the total production (including surplus production and process error) that are removed due to fishing and converted to absolute units, numbers, through $\exp(\log K)$.

The model fits to the observed total catch in numbers using a Student's t-distribution likelihood:

$$\log(\text{Removals}_{obs,t}) \sim \text{Student-t} \left(\nu_c, \log(\text{Removals}_t) - \frac{\sigma_{C,t}^2}{2}, \sigma_{C,t} \right)$$

where the degrees of freedom parameter ν_c controls the tail behavior⁴, and the observation error $\sigma_{C,t}$ allows for time-varying uncertainty in catch observations. However, in the diagnostic model configuration $\sigma_{C,t} = 0.2$ for all t .

3.5 Prior development

The baseline prior distributions for leading estimated model parameters are specified in Table 2. A description of the derivation of these prior distributions follows.

Informed prior distributions for R_{Max} , and the Pella-Tomlinson shape parameter n were developed using numerical simulation based on biological knowledge. Informed prior distributions for $\log(K)$, and baseline catchability q_{eff} were developed using numerical simulation based on existing fishing data observations (e.g., catch and effort). Priors for other estimated leading model parameters (initial depletion x_0 , AR1 correlation in catchability deviates ρ , variability in catchability deviates σ_{qdev} , process error σ_P , index observation error $\sigma_{O,add}$, and degrees of freedom of the catch likelihood ν_{catch}) were based on expert opinion and values observed in the literature. The naïve or input prior distributions were refined using a prior pushforward analysis.

3.5.1 Biological simulation framework

Informative priors for key population parameters were developed through biological simulation using Monte Carlo sampling from biologically plausible parameter distributions⁵. The simulation framework incorporated uncertainty and correlations in life history parameters:

⁴As $\nu_c \rightarrow \infty$ the likelihood equation converges to the lognormal distribution $\text{Lognormal} \left(\log(\text{Removals}_t) - \frac{\sigma_{C,t}^2}{2}, \sigma_{C,t} \right)$

⁵The R code for the simulation [01-01-define-rmax-prior.r](#) is publicly available on [GitHub](#).

- **Growth parameters:** Francis parameterization with independent sampling
- **Natural mortality:** Reference mortality M_{ref} negatively correlated with maximum age ($r = -0.3$)
- **Maturity:** Length-based logistic function parameters
- **Reproduction:** Steepness parameter for stock-recruit relationship and sex ratio
- **Weight-length relationship:** Correlated allometric parameters ($r = -0.5$) with biological constraints
- **Fishery selectivity:** Length at first capture for depletion prior

Parameter combinations (Table 1) were filtered to ensure biological realism and included correlations between life history parameters reflecting biological trade-offs (e.g., shorter lived individuals having higher natural mortality rates).

3.5.1.1 Maximum intrinsic rate of increase (R_{Max}) prior

R_{Max} was calculated by numerically solving the Euler-Lotka equation using bounded optimization (nlminb function in R) to find the value of R_{Max} that satisfies the equation within convergence criteria. Starting values and bounds were set to ensure biologically realistic solutions. Formulation of the Euler-Lotka equation followed the approach implemented in openMSE (Hordyk et al., 2024; Stanley et al., 2009):

$$\alpha \sum_{a=1}^{A_{Max}} l_a f_a \exp(-R_{Max} \times a) = 1$$

where α is the slope at the origin of the stock-recruitment curve, l_a is the probability of survival to age a , and f_a is the fecundity (reproductive output) at age a . Additional technical detail can be found in Section 11.1 of the Technical Annex. Repeating this calculation across the 500,000 parameter combinations resulted in a prior distribution of R_{Max} conditioned on plausible biology and life-history characteristics. The resulting distribution was first filtered to values of $R_{Max} < 1.5$ representative of what the literature (Gravel et al., 2024; Hutchings et al., 2012) indicates is most likely for teleosts. A lognormal distribution was fit to the filtered distribution to derive input prior parameters for the Stan model.

3.5.1.2 Pella-Tomlinson shape parameter (n) prior

Similarly, a prior distribution for the Pella-Tomlinson shape parameter n was derived from the filtered R_{Max} values and associated biological parameter combinations through a two-step process. First, the inflection point of the production function was estimated using the relationship established by Fowler (1988):

$$\phi = 0.633 - 0.187 \times \log(T_G \times R_{Max})$$

where ϕ is the inflection point (representing the population level at which maximum sustainable yield occurs as a fraction of carrying capacity), T_G is the generation time calculated following Grant & Grant (1992) as $T_G = \sum_{a=1}^{A_{Max}} a \times l_a \times f_a / \sum_{a=1}^{A_{Max}} l_a \times f_a$, and R_{Max} is the maximum intrinsic rate of increase.

The Pella-Tomlinson shape parameter n was then derived from the inflection point through the mathematical relationship $\phi = (1/n)^{1/(n-1)}$, solved numerically using bounded optimization in R as done for R_{Max} . This transformation links the biologically-derived inflection point to the production function curvature, where $n = 2$ corresponds to the symmetric Schaefer model (e.g., MSY achieved at depletion $D = 0.5$). Lower values of n ($n < 2$) indicate more productive stocks that achieves MSY at $D < 0.5$ and higher values of n indicate less productive stocks with MSY occurring at $D > 0.5$. A lognormal distribution was fit to the distribution of

n to derive input prior parameters for the Stan model. Additionally, linking the shape parameter n to R_{Max} necessitated a multivariate prior structure so the correlation between the two parameters was estimated as well.

3.5.2 Fishing data simulation framework

Prior distributions for carrying capacity $\log(K)$ and initial catchability coefficient q_{eff} were derived using catch and effort data combined with the biologically-informed parameter distributions⁶.

3.5.2.1 Carrying capacity prior ($\log(K)$)

The carrying capacity prior was constructed by assuming the maximum observed catch approximates maximum sustainable yield (MSY). This provides a conservative assumption for the upper limit of population scale⁷ as larger MSY correlates positively with larger population scale. To account for uncertainty in the maximum catch $\sim$ MSY relationship (e.g., true MSY was above or below the maximum observed catch), candidate MSY values were sampled from a lognormal distribution centered on the maximum observed catch with moderate levels of uncertainty ($MSY \sim \text{Lognormal}(\log(\text{max catch}), 0.5)$).

For each combination of MSY, biologically-derived R_{Max} and shape parameter n , carrying capacity was calculated as:

$$K = \frac{MSY}{r \times \phi \times (1 - \phi)^{n-1}}$$

where $\phi = (1/n)^{1/(n-1)}$ represents the inflection point of the production function. For the Fox model special case where $n = 1$, the relationship simplifies to $K = e \times MSY/r$. Non-finite values and extreme outliers ($\log(K) > 20$) were replaced through resampling from the valid parameter space to ensure numerical stability. A lognormal distribution was fit to the filtered distribution to derive input prior parameters for the Stan model.

3.5.2.2 Initial effort catchability prior (q_{eff})

Next, the initial catchability coefficient prior was developed using the relationship between CPUE and stock abundance:

$$q_{eff} = \frac{CPUE}{K}$$

where CPUE was calculated from early fishery data (first 10 years of catch and effort records) when the stock was assumed to be closer to carrying capacity. The prior distribution was constructed by randomly pairing CPUE values sampled from the early period with carrying capacity values from the $\log(K)$ prior distribution to generate a distribution of q_{eff} values incorporating both the uncertainty in population scale

⁶This is applied in the R script `03-06-new-logK-prior-pushforward-bspm_estqsimple_softdep_mvprior_x0.r` lines 64-92 which is publicly available on [GitHub](#).

⁷The upper limit of population scale is typically more challenging to estimate than the lower population limit as population scales of infinite sizes can support any observed catches. However, there typically is a hard floor to population scale in that it must be sufficiently large to maintain a population that produces the time series of observed catches.

and the variability observed in early fishery CPUE. Again, a lognormal distribution was fit to the filtered distribution to derive input prior parameters for the Stan model.

3.5.3 Other priors

As mentioned previously, initial prior distributions for the remaining leading model parameters (initial depletion x_0 , AR1 correlation in catchability deviates ρ , variability in catchability deviates σ_{qdev} , process error σ_P , index observation error $\sigma_{O,add}$, and degrees of freedom of the catch likelihood ν_{catch}) were specified either using expert opinion or literature.

Given that the limited large-scale industrial fishing existed in the SWPO prior to the expansion of the Japanese distant-water longline fleet in 1952, the population was assumed to start at an unfished condition with some uncertainty. Initial depletion x_0 was modeled with a lognormal distribution centered on 1. A lognormal prior was used for the standard deviation of the process error where the parameters were converted from the JABBA default prior for process error (Winker et al., 2018). The degrees of freedom ν_{catch} parameter assumed a gamma prior based on suggestions in the literature that it serves as a sensible default choice (Juárez & Steel, 2010). The prior for index observation error $\sigma_{O,add}$ assumed a half-normal distribution following ISC (2024) which provides sufficient support for a broad range of extra estimated observation error. For the autocorrelation coefficient in catchability deviations (ρ) and the standard deviation of catchability deviations (σ_{qdev}), initial priors were derived from data-filtering simulation results and specified as a bivariate distribution on the transformed scale: $\tanh^{-1}(\rho)$ and $\log(\sigma_{qdev})$.

3.5.4 Prior pushforward

Following Kim & Neubauer (2025), a prior pushforward analysis was conducted to refine the naïve prior distributions derived from the biological and fishing data simulation frameworks. Random parameter combinations drawn from the input prior distributions were passed through the diagnostic Stan model used for inference but with the likelihoods turned off (`fit_to_data = 0L`), generating simulated population trajectories and fishing dynamics under the model structure. This approach ensures that simulated trajectories are consistent with the actual model structure used for inference.

Parameter combinations were filtered through sequential criteria based on population viability and fishery realism (Figure 3):

- **Initial survival filter** (`iter_surve`): Terminal year depletion $> 1\%$ (`mindep > 0.01`), minimum population size $> 15,000$ individuals which matches recent observed catch levels (`minN > 15000`), and biologically reasonable shape parameter (`shape < 20`)
- **Fishing mortality filter** (`iter_f`): Maximum fishing mortality < 2.5 (`maxF < 2.5`) and average fishing mortality < 0.8 (`avgF < 0.8`) to ensure realistic exploitation levels
- **Catch consistency filter** (`iter_catch`): Simulated catches within realistic bounds relative to observed data: maximum catch between $0.5\text{--}5.0\times$ observed maximum, minimum catch $> 0.5\times$ observed minimum, and average catch between $0.5\text{--}5.0\times$ observed average
- **CPUE consistency filter** (`iter_cpue`): Early period CPUE (first 10 years) between $0.5\text{--}5.0\times$ observed early CPUE average to ensure realistic abundance levels during the initial fishery development

Parameter combinations passing all filtering stages (`iter_cpue`) were used to fit updated multivariate prior distributions using maximum likelihood estimation (Figure 4). Given that a number of parameters

showed correlations following the prior pushforward, a 7-dimensional multivariate prior was developed for core parameters: $\log(K)$, $\log(R_{Max})$, $\log(n)$, $\log(x_0)$, $\log(q_{eff})$, $\tanh^{-1}(\rho)$ and $\log(\sigma_{qdev})$. Correlations with absolute values < 0.1 were reset to zero to avoid weak spurious correlations, retaining only meaningful parameter relationships. Though $\log(x_0)$ was included in the multivariate prior, for convenience, it was treated as an unchanged univariate prior from the initial naïve prior distribution since the pushforward did not provide an update to this parameter.

Prior distributions for process error σ_P , index observation error $\sigma_{O,add}$, and degrees of freedom of the catch likelihood ν_{catch} were unchanged following the prior pushforward analysis and remained as univariate priors. Final prior distributions are listed in Table 2.

3.6 Description of one-off sensitivities

During the course of model development 38 one-off sensitivity models, organized into 12 groups, were investigated to explore alternative assumptions, model structures and sensitivities to data inputs. These models are listed in Table 3 and are described in further detail below.

3.6.1 Index

The diagnostic case model fits only to the DWFN index. One-off model sensitivities were conducted to explore how model estimates changed as a result of alternative indices:

- 0071 : Fits to the Australian longline index
- 0072 : Fits to the New Zealand recreational sportfish index
- 0073 : Fits to the longline observer data combining the programs from New Caledonia, Fiji, Tonga, and French Polynesia
- 0074 : Fits to the longline observer data excluding the French Polynesia program
- 0075 : Fits only to the longline observer data from the French Polynesia program

The French Polynesia observer program data was treated separately as it showed differing trends from the other 3 observer programs (Neubauer et al., 2025). Additionally, given that the shorter indices (Australia and longline observer) provided little information about population scale, several other models were set-up to simultaneously fit them with the two longer indices (DWFN and New Zealand).

- 0129 : Fits to the DWFN & Australia index
- 0130 : Fits to the New Zealand & Australia index
- 0131 : Fits to the DWFN & Observer (New Caledonia, Fiji & Tonga) index
- 0132 : Fits to the New Zealand & Observer (New Caledonia, Fiji & Tonga) index

3.6.2 Catch observation error σ_C

The diagnostic case model assumes a constant catch observation error of $\sigma_C = 0.2$. Three one-off models explored alternative assumptions about catch uncertainty:

- 0076 : Time-varying σ_C following a power curve declining from 0.5 in early years to 0.2 in recent years, assuming improved catch reporting over time
- 0077 : Constant $\sigma_C = 0.1$, representing lower uncertainty in catch observations throughout the time series

- **0078** : Time-varying σ_C following a power curve declining from 0.5 in early years to 0.1 in recent years, combining both temporal improvement and lower overall uncertainty

3.6.3 Early data

Several models explored sensitivity to problematic early data years, particularly 1952 and 1954 which showed unusual catch or effort patterns. In particular the 1952 catch observation was anomalously low especially given the reported effort, and was identified to have high influence on model estimates. This could be a valid observation, as the Japanese longline fleet expanded into the south Pacific in 1952 they fished closer to the equator which tended to have lower striped marlin catch rates than in subsequent years when they fished between Australia and New Caledonia. Alternatively, this outlier point could be the result of erroneous reporting in either catch or effort. In either case, sensitivity to this data point was investigated:

- **0079** : Recalculates the low 1952 catch observation as the expected catch given the effort and average nominal CPUE from 1953-1957 (1952 catch is increased substantially).
- **0080** : Recalculates 1952 effort given the low 1952 catch and average nominal CPUE from 1953-1957 (1952 effort is decreased substantially)

Additionally, the large catch event in 1954 has been heavily scrutinized. This could be a valid observation as the Japanese longline fleet expended considerable effort during the time of year (3rd and 4th quarters of the year) striped marlin form spawning aggregations, in an area between Australia and New Caledonia where these aggregations form. Regardless, sensitivity to this observation was investigated in case it is a reporting error.

- **0081** : The 1954 catch was reduced by 50%
- **0082** : Combined alternative catch values for both 1952 and 1954
- **0083** : Combined alternative 1952 effort and 1954 catch values

These sensitivities address concerns about sensitivity to potentially anomalous observations in the early years of the fishery when reporting systems were less standardized.

3.6.4 Scaling priors

The diagnostic case model estimates a low population scale relative to the prior. Two models tested the impact of forcing the model toward higher population scale estimates:

- **0093** : Modified priors on $\log(K)$ and catchability (q_{eff}) to favor higher carrying capacity and lower catchability estimates
- **0096** : Same as **0093** but combined with more conservative shape parameter prior (centered on 2)

These models explored whether alternative prior assumptions about population scale could shift model estimates.

3.6.5 Catch observation model

- **0110** : Replaced the default Student-t likelihood for catch observations with a lognormal error structure to test sensitivity to distributional assumptions about catch observation error

3.6.6 Temporal resolution of catchability deviates

The diagnostic case model estimates catchability deviates in single time steps (annual). Four models explored different temporal resolutions:

- 0111 : Catchability deviates fixed across 2-year periods ($n_step = 2$)
- 0112 : Catchability deviates fixed across 3-year periods ($n_step = 3$)
- 0113 : Catchability deviates fixed across 5-year periods ($n_step = 5$)
- 0114 : Catchability deviates fixed across 10-year periods ($n_step = 10$)

These sensitivities tested whether the temporal resolution of catchability changes affected model estimates, with longer periods resulting in a more parsimonious model but potentially diminishing the models ability to fit to data.

3.6.7 Catch reporting

Striped marlin are a commercially valuable non-target species, and as such it is assumed that if they are caught they are retained, and recorded in the logbook. However, given that they are a non-target species it is possible that they are not, or have not always been reported in the logbook and that under-reporting of catches or discarding could occur (e.g., given value of striped marlin and hold space required relative to target species). Three models explored potential systematic under-reporting of catches:

- 0115 : Doubled all observed catch values to test sensitivity to potential 50% under-reporting throughout the time series
- 0116 : Applied a linear multiplier increasing from 1x to 2x observed catch over time, representing improving catch reporting
- 0117 : Applied a linear multiplier decreasing from 2x to 1x observed catch over time, representing deteriorating catch reporting in recent years

3.6.8 Process error σ_P

The diagnostic case model estimated non-zero process error in the early model period, before the CPUE index began, and balanced process error with the catchability deviates in order to fit to the catch.

- 0118 : Restricted process error to be negligible ($\sigma_P \sim 0$) to evaluate the impact of environmental stochasticity on model estimates by forcing the population to follow deterministic surplus production dynamics

3.6.9 Start year

Given potentially anomalous catch observations in 1952 and 1954, along with uncertainty in the accuracy of early catch observations, four models explored sensitivity to the assumed initial conditions by starting the model in different years:

- 0119 : Model starts in 1955 with assumed initial depletion $x_0 = 0.9$, skipping the early data years

Three models were also tested which started the model in the same year as the CPUE index, and varied the level of initial depletion:

- 0120 : Model starts in 1988 with $x_0 = 0.5$
- 0121 : Model starts in 1988 with $x_0 = 0.7$
- 0122 : Model starts in 1988 with $x_0 = 0.9$

3.6.10 Inclusion of effort deviates

Early model development explored a more complex formulation of fishing mortality F_t where effort deviates were also used in addition to the catchability deviates to link effort to F_t :

$$F_t = q_{eff} \times \exp(\delta_{q,t} - \sigma_{qdev}^2/2) \times E_t \times \exp(\delta_{e,t} - \sigma_{edev}^2/2)$$

This formulation of fishing mortality is closer to the full catch-errors with effort deviation approaches available in MULTIFAN-CL (Fournier et al., 1998). Two models tested whether estimating deviations from the effort time series improved model performance:

- 0124 : Estimated effort deviates with 2-year temporal resolution ($n_step = 2$)
- 0126 : Estimated effort deviates with 5-year temporal resolution ($n_step = 5$) for more constrained estimation

3.6.11 Direct estimation of fishing mortality

Rather than estimating fishing mortality indirectly through catchability and effort, two models directly estimated annual fishing mortality rates to see if the simpler formulation for fishing mortality could replicate the results of the diagnostic case model, or if the formulation of fishing mortality significantly impacted model estimates:

- 0127 : Direct F estimation with σ_F calibrated to match the variance in fishing mortality observed in the diagnostic case model
- 0128 : Direct F estimation with σ_F set to 0.25 times the diagnostic case variance, providing more constraint on fishing mortality variability in order to see how estimates of scale are impacted

3.6.12 Alternative R_{Max} prior

The diagnostic case model estimates a fairly productive stock with an R_{Max} that is considerably higher than an analogous species in the Atlantic Ocean, white marlin (*Kajikia albida*). The most recent ICCAT assessment (ICCAT, 2020), implemented using JABBA (Winker et al., 2018), estimated R_{Max} at 0.163 for white marlin. Sensitivity models assuming this lower level of productivity were developed to see the impact it would have on model estimates, particularly population scale.

- 0133 : Apply lower productivity white marlin R_{max} prior
- 0134 : Apply lower productivity white marlin R_{max} prior and apply a more conservative shape parameter prior (centered on 2)

3.7 Model Implementation

The models were implemented using the No-U-Turn Sampler (NUTS) in Stan. Each model was run using 5 independent Markov chains with randomly generated starting values to ensure robust exploration of

the posterior parameter space. Each chain was sampled for 3,000 iterations, with the first 1,000 iterations discarded as warmup to allow for algorithm adaptation. To reduce posterior sample autocorrelation and storage requirements, every 10th samples was retained from the post-warmup iterations, yielding a final sample size of 1,000 draws from the joint posterior distribution across all chains. The NUTS algorithm was configured with strict adaptation parameters to ensure reliable convergence. The adaptation delta was set to 0.99 to reduce the step size and minimize divergent transitions, while the maximum tree depth was increased to 12 to allow for longer trajectories during the dynamic sampling process.

Model convergence and performance were assessed using multiple diagnostic criteria recommended for Bayesian stock assessment workflows (Monnahan, 2024). The potential scale reduction factor ($\hat{R}$) was required to be less than 1.01 for all parameters, indicating convergence across chains. Effective sample sizes were required to exceed 500 to ensure adequate precision in posterior estimates. Additionally, posterior predictive checks were conducted to evaluate model adequacy through comparison of observed versus predicted index and catch values, while parameter identifiability was assessed through visual examination of posterior distributions and correlation structures among estimated parameters (Gabry et al., 2019).

3.8 Model diagnostics

Model performance was comprehensively evaluated using multiple diagnostic criteria recommended for Bayesian stock assessment workflows (Monnahan, 2024). The diagnostic framework included Stan model convergence criteria, data fits, posterior predictive checks, retrospective analysis, and hindcast cross-validation.

3.8.1 Convergence diagnostics

Conventional Stan model convergence diagnostics and thresholds were employed to identify whether posterior distributions were representative and unbiased (Monnahan, 2024). Models were considered to have ‘converged’ to a stable, unbiased posterior distribution if they satisfied the following criteria:

- **Potential scale reduction factor** ($\hat{R}$) less than 1.01 for all leading model parameters, indicating convergence across chains
- **Bulk effective sample size** (N_{eff}) greater than 500 for all leading model parameters, ensuring adequate precision in posterior estimates
- **No divergent transitions** during the NUTS sampling process, which can indicate problematic posterior geometry
- **Maximum tree depth** not exceeded during sampling, ensuring adequate exploration of parameter space

Additionally, trace plots and rank plots were examined to visually assess mixing and convergence behavior across chains (Gabry et al., 2019).

3.8.2 Leave-one-out cross-validation diagnostics

Model predictive performance and influential observations were assessed using leave-one-out cross-validation (LOO-CV) implemented through the *loo* package (Vehtari et al., 2017; Vehtari et al., 2024). LOO-CV provides an estimate of the expected log pointwise predictive density (ELPD) while identifying problematic observations that may unduly influence model predictions.

LOO-CV was also used as a factor to compare performance between models. Following the recommendations of Sivula et al. (2020), models were considered to have meaningfully different performance when the ELPD difference exceeded 4 units. For differences larger than 4 units, statistical significance was assessed by comparing the ELPD difference to twice its standard error, with differences greater than $2 \times$ the standard error considered statistically significant.

The effective number of parameters (p_{loo}) was examined as an additional diagnostic measure, with values substantially larger than the nominal number of model parameters indicating potential model misspecification. Catch and index observations were categorized as having low influence ($\text{Pareto-k} \leq 0.7$), high influence ($0.7 < \text{Pareto-k} \leq 1.0$), or very high influence ($\text{Pareto-k} > 1.0$). Observations with high or very-high influence were investigated to identify potential outliers, data entry errors, or systematic model misspecification that could compromise predictive performance.

3.8.3 Data fits

Model fits to the abundance index and catch time series were each quantified using normalized root-mean-squared error (NRMSE):

$$RMSE = \sqrt{\frac{\sum_{i=1}^n (y_i - \hat{y}_i)^2}{n}}$$

$$NRMSE = \frac{RMSE}{\sum_{i=1}^n y_i / n}$$

where y_i are the observed values and $\hat{y}_i$ are the model predictions. The average NRMSE across all posterior samples was calculated for each data component. Additionally, Probability Integral Transform (PIT) style residuals were calculated for the fit to the index.

3.8.4 Posterior predictive checks

Posterior predictive checks were conducted to evaluate model adequacy by comparing observed data with data simulated from the fitted model. For each posterior sample, synthetic datasets were generated using the estimated parameters, and agreement between observed and predicted datasets was assessed visually.

3.8.5 Retrospective analysis

Retrospective analysis was conducted for each model by sequentially peeling off a year from the terminal end of the fitted index and re-running the model. Data were removed for up to five years, from 2022 back to 2018. Estimates of x in the terminal year of each retrospective peel were compared to the corresponding estimate of x from the full model run to better understand any potential biases or uncertainty in terminal year estimates. The Mohn's ρ statistic (Mohn, 1999) was calculated and presented. This statistic measures the average relative difference between an estimated quantity from an assessment (e.g., depletion in final year) with a reduced time-series of information and the same quantity estimated from an assessment using the full time-series.

Additionally, following Kokkalis et al. (2024) the proportion of retrospective peels (or coverage) where the

relative exploitation rate (U/U_{MSY}) and relative depletion (D/D_{MSY}) were inside the credible intervals of the full model run was calculated. This metric provides an additional perspective on retrospective consistency by evaluating whether the uncertainty estimates from the full model appropriately capture the range of estimates obtained when using reduced datasets.

3.8.6 Hindcast cross-validation

Hindcast cross-validation (Kell et al., 2021) was conducted for each index to determine model performance in predicting the observed CPUE I one-step-ahead into the future relative to a naïve predictor. Briefly, the ‘model-free’ approach to hindcast cross-validation was used, and used the same set of five retrospective peels described in Section 3.8.5.

The ‘model-free’ hindcast calculation is described using the model from the last peel $BSPM_{2017}$ as an example. This model fit to index data through 2017 but included catch through 2022. The model estimates of predicted CPUE in 2018 based on $BSPM_{2017}$ is the ‘model-free’ hindcast for 2018, $\hat{I}_{2018}$. The naïve prediction of CPUE in 2018 is simply the observed CPUE from 2017, $I_{2017} = \check{I}_{2018}$. The absolute scaled error (ASE) of the prediction is:

$$ASE_{2018} = \frac{|I_{2018} - \hat{I}_{2018}|}{|I_{2018} - \check{I}_{2018}|}$$

Repeating this calculation across all retrospective peels for years 2019-2022 and taking the average across ASE values gives the mean ASE or MASE for the model. An MASE value less than one indicates that the model has greater predictive skill than the naïve predictor.

3.8.7 Model-free hindcasts

Additional model-free hindcasts, similar to those described in Section 3.8.5, where the model was run for the full period 1952-2022 but without the last y years of index data. Hindcasts excluding the last 5, 10, 15, and 20 years of index data were conducted in order to evaluate the models ability to recapture the dynamics of the full model in the terminal year. Good hindcast performance over a long time horizon is an indication of a well-determined production function, and can give an idea of the reliability of the predictions from the full model.

3.9 Forecasts

Simple, status quo catch-based stochastic projections over a 10-year window are provided for each model. For each model, projected catch was taken as the average catch over the terminal 5 years (2018-2022). Process error in the projection period was randomly resampled from the estimated process error in the main model period.

4 Results

4.1 Diagnostic case

4.1.1 Model convergence and diagnostics

The diagnostic case model 0100 achieved satisfactory convergence based on standard Hamiltonian Monte Carlo diagnostics (Table 4). $\hat{R}$ values were below the 1.01 threshold and effective sample sizes exceeding the recommended minimum of 500 (100 samples per chain) for all parameters. No divergent transitions were observed across chains, and maximum tree depth was not exceeded during sampling, indicating successful exploration of the posterior space. Additionally, LOO-CV identified only two observations with Pareto-k > 0.7 indicating the model structure and choice of likelihoods adequately matched the fitted data. Lastly, the effective number of estimated parameters p_{loo} was well below both the number of observations and the actual estimated parameter count indicating that model is likely to be reasonably well specified.

4.1.2 Model fit to data

4.1.2.1 Catch data fit

The diagnostic case model adequately captured the trend and magnitude of the observed catches (1952-2022) with the obvious exceptions of the 1952 and 1954 observations (Figure 5). Use of a catch likelihood, student-t, with heavier tails made the model more robust to these outlier observations, and accordingly estimated larger and smaller predicted catches, respectively, for these two observations. However, use of the student-t distribution along with an assumed 20% CV on the observation error did mean that catch observations were not fit exactly though the observed catches did tend to fall within the posterior predicted distribution (Figure 6). Given that SWPO striped marlin is a non-target species, some uncertainty in the predicted catches is considered to be appropriate.

4.1.2.2 CPUE index fit

Diagnostic case model fit to the standardized DWFN CPUE index (1988-2022) was evaluated using NRMSE and showed reasonable fits to the observed index given the high variability and observation error (Table 5). The model also successfully captured the general declining trend in standardized CPUE from 1988-2022 (Figure 7). PIT residuals did not show persistent positive or negative patterns, however they did indicate marginally better fit within the last decade of the index relative to the earlier portion (Figure 8). Posterior predictive checks confirmed that models replicated key features of the observed CPUE data (Figure 9).

4.1.3 Model validation

Overall retrospective bias was within the acceptable range proposed by (Hurtado-Ferro et al., 2015), however it did indicate a slight underestimate to the population recovery indicated by the DWFN index (Table 5; Figure 10). This is likely driven by the large observation in 2021 that is used by the full model to drag estimates up from the lower values observed in 2015-2019. Estimates of fishing mortality (U/U_{MSY}) and depletion (D/D_{MSY}) relative to MSY fell within the credible intervals of the full model run 100% of the time (Table 5). For hindcast cross-validation, the diagnostic model substantially exceeded the performance of the naïve predictor (Table 5; Figure 11). This provides support that models successfully identified and estimated a production function for the stock.

4.1.4 Parameter estimates

Following the guidance from Kim & Neubauer (2025), posterior distributions of parameter estimates were compared to the realized prior in addition to the naïve or input prior. The realized prior represents the effective prior distribution that results from passing naïve parameter combinations through the model structure with likelihoods turned off, revealing how model constraints and structural assumptions modify the input priors before data-based updating occurs (Kim & Neubauer, 2025). In the current analysis, there was little difference between the naïve and realized priors given the approach taken in developing the priors. Based on these comparisons the diagnostic model showed substantial posterior learning across several key leading parameters (Figure 12; Table 6), indicating that the data contained sufficient information to update parameter estimates from their prior distributions.

The carrying capacity $\log(K)$ exhibited substantial posterior learning, with the combination of index and catch data in the diagnostic case model strongly supporting lower population scales than suggested by either the naïve or realized priors. Similarly, the intrinsic rate of increase R_{Max} showed strong posterior updates, with the data favoring lower productivity than the biological simulation framework initially suggested. Inverse transform sampling to match the R_{Max} posterior distribution to the input biological simulation framework prior for R_{Max} indicates that the lower productivity is characterized by pronounced shifts toward lower steepness values h (Figure 13).

The median estimate of R_{Max} from the diagnostic case, 0.396 (Table 6), is considerably higher than estimates for an analogous species, white marlin (*Kajikia albida*), assessed using a JABBA surplus production model (ICCAT, 2020; Winker et al., 2018) in the Atlantic Ocean where $R_{Max} = 0.163$ (95% credible interval: 0.122-0.215). However, it is worth noting that the ICCAT assessment assumed a much more precise prior for R_{Max} , and that the 95% credible intervals of the diagnostic case model almost completely cover the ICCAT posterior distribution. Notably, the Atlantic white marlin JABBA model showed remarkable consistency with stock status and trend estimates from a fully integrated, length-structured model implemented in Stock Synthesis (ICCAT, 2020).

Shape and initial depletion parameters did not demonstrate notable posterior updates (Figure 12). The Pella-Tomlinson shape parameter n showed limited deviation from the prior, suggesting the data provided weak information about production function curvature. Initial depletion x_0 also remained close to the assumed unfished prior, indicating little information in the data to suggest large-scale fishing pressure prior to 1952.

The catchability and temporal variability parameters showed large posterior updates (Figure 12). The base catchability coefficient q_{eff} showed substantial learning, with strong negative correlations with $\log(K)$. This demonstrates a trade-off between population scale and catchability in order to fit the observed catches. If a more precise prior on q_{eff} were to be developed, this would be expected to set population scale and dramatically reduce uncertainty in model estimates. The autocorrelation parameter ρ and catchability variability σ_{qdev} both showed posterior updates, with the data informing a highly correlated temporal structure of catchability deviations.

Lastly, process error σ_P showed moderate posterior learning, while additional observation error $\sigma_{O,add}$ demonstrated substantial updates from the prior (Figure 12). This indicated that the model estimated additional uncertainty beyond the input index observation error, rather than using process error, to reconcile the observed CPUE variability. This results in a smoother year-on-year population trajectory. The degrees of

freedom parameter ν_{catch} for the catch likelihood showed substantial updating (Figure 12), indicating the model preferred a Student-t distribution with very heavy tails in order to adequately account for the catch observations given the level of observation error.

4.1.5 Model-free hindcasts

Model-free hindcasting was conducted to evaluate the model’s ability to predict population dynamics when fitted to progressively truncated index datasets. The diagnostic model was compared against retrospective peels excluding the final 5, 10, 15, and 20 years of index data (terminating in 2017, 2012, 2007, and 2002, respectively), with the model providing estimates through the full model period based on based only on the estimated production function and predicted catch.

The model demonstrated remarkably consistent performance across most hindcast scenarios (Figure 14) providing strong evidence of a well-determined production function. In general, the hindcast models successfully captured the general increasing trend in the standardized index after 2015. Model predictions for the DWFN CPUE index showed good coherence between the diagnostic model and all retrospective peels except the 15 year hindcast model, with overlapping 95% credible intervals throughout most of the time series. Estimates from the 15 year hindcast model appear to be strongly influenced by the series of higher observations (2003-2005) which cause it to underestimate the decline in the observed index.

Population dynamics estimates showed reasonable consistency across hindcast scenarios (Figure 15), with the notable exception of the 15 year hindcast model mentioned in the previous paragraph. Depletion trajectories remained largely stable, with all models estimating similar patterns of decline from unfished conditions to minimum levels around the early 1990s, though predictions begin to diverge slightly in the recovery period post 2015. Population scale estimates also showed good agreement, though with increasing uncertainty for earlier time periods in longer hindcasts, as expected. The stock status metrics (D/D_{MSY} and F/F_{MSY}) showed fairly good consistency across most hindcast scenarios particularly for F/F_{MSY} . Additionally, given the The estimated trajectories for these key management metrics were nearly identical between the diagnostic model and all retrospective peels, suggesting that management advice derived from this model would have been stable regardless of when the assessment was conducted over the past two decades.

This consistency across multiple extended hindcast periods provides support that the underlying population dynamics are well described, given the available data. This gives confidence that projections made from the full model will likely capture the stock trajectory within the near to medium term, assuming stationarity in future productivity and carrying capacity.

4.2 One-off sensitivities

As described in Section 3.6, 38 different one-off sensitivities were conducted to either explore plausible uncertainty in model assumptions and configuration, or to build intuition about the model dynamics and gain confidence in the results.

4.2.1 Index

The first one-off sensitivity group explores alternative abundance indices compared to the diagnostic case model that fits only to the DWFN index. This group includes models testing individual indices (0071-0075) and combinations of indices (0129-0132). In general model fit to individual indices (0071-0075) was acceptable

(Figure 16), though model predictions tended to over-predict the last 5 years of the New Zealand recreational sportfish index (0072). However, observation errors for these observations were fairly large relative to earlier observations meaning that the model was not overly constrained to fit them, and could instead follow the trajectory implied by the production function. Notably, only the two longer indices (DWFN and New Zealand recreational) showed sufficient contrast with the catches to meaningfully move the scale estimate from the prior and models fitting either of those indices estimated similar population scales (Figure 17). The Australia longline index (0071) showed some posterior update but still estimated a larger overall population scale, the observer index with only data from French Polynesia (0075) also showed a noticeable posterior update however had poor convergence diagnostics (Table 4).

For the models fitting to multiple simultaneous indices (0129-0132), estimates of model quantities were largely consistent with the diagnostic case model in terms of scale and trend (Figure 18), even for models fitting to the New Zealand index. This indicated that scale estimates from models fitting to shorter indices may not be reliable. Additionally, fits to the shorter indices were not substantially degraded when also simultaneously fitting to the longer indices.

4.2.2 Catch observation error σ_C

The diagnostic case model assumes a constant catch observation error of $\sigma_C = 0.2$. Three one-off models (0076 - 0078) explored alternative assumptions about catch uncertainty including time-varying error structures and different constant error levels. In general parameter estimates were very consistent regardless of the assumed catch observation error level and temporal pattern (Figure 19). The one exception was with the estimation of ν_{catch} , which showed a preference for a Student-t distribution with lighter tails (e.g., more *Normal* distribution-like) when catch observation error was structured to decline overtime from 0.5 following a power curve (0076 & 0078). This suggests that the earlier observations may warrant a higher level of observation error. Regardless, estimated model quantities, including key management metrics D/D_{MSY} and F/F_{MSY} , were largely unchanged no matter what structure was used with the exceptions that models assuming less catch uncertainty had predictably less estimated uncertainty, and models using a power curve to structure the observation error estimated lower levels of initial nominal CPUE (Figure 20).

4.2.3 Early data

Several models explored sensitivity to problematic early data years, particularly 1952 and 1954 which showed unusual catch or effort patterns. Five one-off models (0079 - 0083) explored alternative assumptions about the catch and or effort in these years. Parameter estimates were very similar across all formulations (Figure 21). However, estimation of ν_{catch} , showed a preference for a Student-t distribution with lighter tails when either alternative catch or effort was assumed for the 1952 observation. This in combination with the results from Section 4.2.2 suggests that the 1952 observation may be a reporting error and could be modified or downweighted with larger observation error. Fortunately, estimated model quantities, including key management metrics, do not appear sensitive to the treatment of the 1952 observation (Figure 20).

4.2.4 Scaling priors

The diagnostic case model estimates a low population scale relative to the prior. Two models (0093 & 0096) tested the impact of forcing the model toward higher population scale estimates through modified priors on carrying capacity and catchability. This was more of an academic sensitivity analysis to see if

the model could be moved through the prior information from the low estimate. While the sensitivity was successful in shifting the estimated scale higher (Figure 23), it showed a clear trade-off with the ability to fit the index (Figure 24) and showed an incompatibility between the data and larger estimates of scale. This provides additional evidence that the smaller estimates of population scale are driven by the data.

While not investigated in the current analysis, it would be interesting to see how sensitive estimates of population scale are to just changes in the precision of the catchability prior as suggested in Section 4.1.4.

4.2.5 Catch observation model

A single model (0110) replaced the default Student-t likelihood for catch observations with a lognormal error structure which is less statistically robust and more impacted by outlier observations. This tested sensitivity to distributional assumptions about catch observation error. Parameter estimates, particularly those related to population scale, showed some sensitivity to choice of catch observation model (Figure 25), and this did translate into some differences in estimated management quantities (Figure 26). However, as previous sensitivity analyses have identified (Section 4.2.2 & Section 4.2.3), some of these early observations appear less reliable and should potentially be made less influential in the likelihood (e.g., use of a Student-t distribution). This finding, combined with the 0110 model showing substantially more observations with large Pareto-k values (Table 4) did not support further use of the lognormal error structure.

4.2.6 Temporal resolution of catchability deviates

The diagnostic case model estimates catchability deviates in single time steps (annual). Four models explored different temporal resolutions with catchability deviates fixed across 2, 3, 5, and 10-year periods (0111 - 0114). Reducing the temporal resolution of the catchability deviates (increasing n_{step}) predictably degraded fit to the catch as n_{step} increased (Table 5; Figure 27). It also changed estimates of σ_P and q_{eff} (Figure 28) which in turn impacted estimates of management quantities particularly depletion and D/D_{MSY} (Figure 29), again particularly for larger values of n_{step} . Reducing the temporal resolution of the catchability deviates did improve model parsimony, and the model with $n_{step} = 2$ (0111) was the best performing according to LOO-ELPD. However, the LOO-ELPD for this model was not statistically different to the diagnostic case model (ELPD difference = 0.9074112; Table 4) and model estimates were nearly identical. In general, the increase in process error estimation as n_{step} increased, the reduced ability to fit the catch were undesirable, and the model equivalence according to LOO-ELPD supported maintaining the diagnostic case model with $n_{step} = 1$.

4.2.7 Catch reporting

Striped marlin are a commercially valuable non-target species, and systematic under-reporting of catches or discarding could occur. Three models (0115 - 0117) explored potential systematic under-reporting scenarios including constant and time-varying multipliers. As expected, increasing the magnitude of all the catches by $2\times$ (0115) shifted the population scale of the population up by ~ 2 (Figure 30) though relative quantities were unchanged (Figure 31). However, increasing either the terminal (0116) or initial (0117) catches by $2\times$ resulted in an increase in population scale, change in the depletion trend, and change in the management advice (Figure 31). The 0116 model represented a scenario where potential non-reported catch increased over time, but resulted in a less depleted stock with a removals time series that increased. The 0117 model

represented a scenario where non-reported catch decreased over time producing a more depleted stock with a decreasing removals time series.

Though there is no evidence at this stage that this level of systematic under-reporting of catches is occurring for SWPO striped marlin, it is useful to be aware of the impacts that changes to our perception of population removals can have on our understanding of stock status. Additionally, for future analyses that may consider changes to the level of catch reporting, it may be useful to do so on a *per hook* basis to take into account the impact of the change in the number of hooks fished over time on total population removals.

4.2.8 Process error σ_P

The diagnostic case model estimated non-zero process error in the early model period before the CPUE index began. Process error was restricted $\sigma_P \sim 0$ as a one-off sensitivity (0118) in order to evaluate if larger initial process error deviates impacted scale estimates or other management quantities. Turning-off the process error did degrade fit to the index slightly (Figure 32), but overall parameter estimates (Figure 33) and management quantities (Figure 34) were largely insensitive to a lack of process error. This confirms that the early process error deviates are not impacting the estimation of key management quantities, and also provides further evidence of a well-determined production function.

4.2.9 Start year

Given potentially anomalous catch observations in 1952 and 1954, four models (0119 - 0122) explored sensitivity to the assumed initial conditions. Models started in either 1955 or 1988 with varying levels of initial depletion. In every case, model results (Figure 35) appeared to be fairly sensitive to the choice of prior, however all models except perhaps model 0122 (1988 start with $x_0 = 0.9$) still estimated substantially lower population carrying capacity $\log(K)$ than the prior (Figure 36). However, these models all assumed fairly restrictive priors on x_0 , and future analyses could evaluate sensitivity to less informative priors or alternatively develop more specific priors for given years based on a prior pushforward analysis.

4.2.10 Inclusion of effort deviates

Early model development explored a more complex formulation of fishing mortality where effort deviates were used in addition to catchability deviates. Two models (0124 - 0126) tested whether returning to a full MULTIFAN-CL style *catch-errors* formulation and estimating effort deviates improved model performance. Estimating effort deviates as expected, did produce an improved fit to the catch (Figure 37) however parameter estimates (Figure 38) and model quantities (Figure 39) showed negligible differences from the diagnostic case model.

4.2.11 Direct estimation of fishing mortality

Rather than estimating fishing mortality indirectly through catchability and effort, two models (0127 - 0128) directly estimated annual fishing mortality rates. This tested whether the simpler formulation for fishing mortality could replicate the results of the diagnostic case model, or if the highly parametrized structure of the fishing mortality calculation determined model outcomes. Removing the relationship with effort, and directly estimating fishing mortality allowed the catch to be fit exactly (Figure 40) though this resulted in high levels of catch observations showing large Pareto-k values (Table 4) which is undesirable. Matching the prior

variability on fishing mortality σ_F to the level of variability in fishing mortality seen in the diagnostic case model (0127) more or less replicated the estimates from the diagnostic case model (Figure 41). Assuming a smaller, more restrictive prior on σ_F (0128) resulted in a large increase to population scale to accomodate a time-series of fishing mortality with smaller variability (Figure 41). These results showed that model outcomes were insensitive to how fishing mortality was parametrized, and reaffirmed the sensitivity of model results, particularly population scale, to the priors placed on fishing mortality.

4.2.12 Alternative R_{Max} prior

The diagnostic case model estimated a stock with higher productivity than an analogous species, Atlantic white marlin. Two models (0133 - 0134) tested sensitivity to a more conservative productivity assumption based on the white marlin R_{Max} estimates from ICCAT (ICCAT, 2020). As hypothesized, forcing productivity R_{Max} to be lower did result in the model estimating a larger population scale (Figure 42). However, this did not necessarily translate to differences in estimated relative management metrics (depletion, D/D_{MSY} , or F/F_{MSY}), particularly in recent years.

4.3 Model ensemble

Following extensive sensitivity analysis, and iterative rounds of model development, a four model ensemble is proposed to be considered for the provision of management advice:

- 0100 : DWFN index & life history based shape prior
- 0102 : New Zealand index & life history based shape prior
- 0105 : DWFN index & alternative shape prior
- 0107 : New Zealand index & alternative shape prior

Both candidate indices were the only indices of sufficient length and contrast to inform estimates of population scale. Additionally, the shape parameter is very difficult to estimate from data, and as such model estimates are dependent on the choice of prior. While the baseline shape parameter prior (Table 2) may be sensibly derived from striped marlin life history characteristics, this derivation is sensitive to underlying assumptions and relationships (Fowler, 1988; Grant & Grant, 1992). In the absence of other information, a conservative assumption for the shape parameter is to assume it is centered on $n \sim 2$, with uncertainty from Table 2. This reduces the production model to a Schaefer-type production model where $D_{MSY} = 0.5$.

4.3.1 Ensemble diagnostics and fit

4.3.1.1 Convergence

All ensemble models achieved satisfactory convergence based on standard Hamiltonian Monte Carlo diagnostics (Table 4). All models met convergence criteria with $\hat{R}$ values well below the 1.01 threshold and effective sample sizes exceeding the recommended minimum of 500 (100 samples per chain). No divergent transitions were observed across chains, and maximum tree depth was not exceeded during sampling, indicating successful exploration of the posterior space. The proportion of observations with Pareto-k > 0.7 was very low ($< 5\%$) across all models and the effective number of estimated parameters p_{loo} was well below both the number of observations and the actual estimated parameter count.

4.3.1.2 Data fit

Model fits to the catch data were equivalent between models and captured the trend and magnitude of the catches adequately (Figure 43). For models fitting to the same index (e.g., 0100 & 0105), the alternative assumption for shape n had no influence on the estimated fit. However, comparing relative fits between models fitting to different indices, the models fitting to the DWFN index did a noticeably better job (lower RMSE) capturing the trend than models fitting to the New Zealand index (Table 5; Figure 44). As mentioned in Section 4.2.1, models fitting to the New Zealand index failed to capture the trend over the last five years.

4.3.1.3 Model validation

Overall retrospective bias was within the acceptable range proposed by (Hurtado-Ferro et al., 2015) for all models, with the models fitting to the New Zealand index showing positive retrospective bias rather than negative for models fitting to the DWFN index (Table 5). Furthermore, all quantities were within the 95% credible interval of the full time-series model for all peels (Table 5; Figure 45).

For hindcast cross-validation, both models fitting to the DWFN index substantially exceeded the performance of the naïve predictor (Table 5; Figure 46). Models fitting to the New Zealand recreational index showed poor performance ($MASE > 1$), as the models were unable to follow the declining trend over the last five years.

4.3.2 Population dynamics and trends

The estimated depletion⁸ time series from the model ensemble shows a decline from the assumed unfished state in 1952 to minimum levels around the mid-2010s (Figure 47). The decline was most pronounced during the 1950s and 1960s, followed by relatively stable depletion levels from the 1970s through the early 2000s which fluctuated round $\sim 0.4 - 0.6$. The stock has been estimated to be recovering from the mid-2010s. Credible intervals around depletion estimates are relatively wide, reflecting inherent uncertainty in the population scale.

The estimated population time series mirrors these depletion patterns. It is important to note that while the uncertainty in absolute population size is substantial, as reflected in wide credible intervals, this uncertainty is asymmetric. The model is fairly confident that the model can not be much smaller than what it currently is estimated to be due to the hard constraint of at least being large enough to support the 70 year time-series of catches. As seen in the one-off sensitivities (Section 4.2.4 & Section 4.2.12), uncertainty to the high-side is driven in part by the prior assumptions for $\log(K)$, q_{eff} , and R_{Max} , as well as by model structural choices with regards to treatment of catch uncertainty (Section 4.2.5) and structure of fishing mortality (Section 4.2.10).

Process error estimates showed consistent temporal variability across ensemble models, with the largest deviations occurring in the 1970s and 1980s. This indicates that removals alone could not explain the observed changes in the standardized indices in these time periods, suggesting additional factors influencing population dynamics during these periods or insufficient observation error.

[Fishing mortality estimates display high variability throughout the time series, with substantial uncertainty reflected in wide credible intervals. The F time series shows episodic peaks during the 1970s and 1980s,

⁸Note that depletion for the surplus production model is a *static* depletion defined relative to total unfished numbers $D_t = P_t/P_0$; where P_t is the total population level in numbers at time t , and $P_0 = K$ the unfished population numbers or carrying capacity. This is different from the *dynamic* depletion of spawning biomass $SB_t/SB_{t,F=0}$ that is typically reported from integrated age-structured models such as MULTIFAN-CL or Stock Synthesis (Methot Jr. & Wetzel, 2013).

corresponding to periods of intensive fishing pressure, followed by more variable but generally declining trends in recent decades. This pattern is consistent with the observed recovery in population depletion during the same period.

Catchability deviations demonstrate substantial temporal variation, particularly during the early period of the fishery. The estimates show generally declining trends from elevated levels in the 1950s and 1960s toward more stable values in recent decades. This temporal pattern likely reflects changes in fishing technology, fleet behavior, and spatial distribution of fishing effort over the model period. Nominal CPUE estimates from the model ensemble track the general declining trend observed in the standardized index, showing elevated values during the early unfished period followed by substantial declines through the 1980s and 1990s. Recent nominal CPUE estimates show stabilization at lower levels consistent with the moderate depletion levels estimated for the contemporary period, though with continued high uncertainty around point estimates.]

4.3.3 Stock status

4.3.4 Future projections

5 Discussion

5.1 General remarks

The three BSPM variants provide a consistent picture of stock trajectory and current status. All models estimated similar depletion patterns: decline from unfished conditions in 1952 to minimum levels around the early 1990s, followed by gradual recovery through recent decades. This trajectory suggests the stock is currently recovering. While median estimates indicate the stock was not overfished or undergoing overfishing during the assessment period, the 0003-2024cpueFPrior model suggests some recent risk of falling below these thresholds.

The BSPM approach addresses key limitations encountered in previous integrated assessments while providing robust stock status estimates. The primary advantage lies in condensing complex biological and fishery uncertainties into a manageable parameter set. Integrated assessments can be influenced by size composition data, which are sensitive to assumptions about growth, mortality, selectivity, and reproduction parameters. When these assumptions are uncertain or mis-specified, population scale estimates informed by size composition data can be biased. Production models circumvent this issue by aggregating these uncertainties into productivity parameters, allowing more efficient exploration of plausible population dynamics while maintaining biological realism through informed priors.

5.2 Challenges, limitations & key uncertainties

Several important limitations affect this analysis. The assumption of no age-specific population dynamics may be problematic when mortality or reproductive potential varies significantly by age. This simplification prevents incorporation of biological information beyond production function parameters and excludes explicit use of size composition data.

The aggregated nature of surplus production models can potentially bias MSY estimates when age-specific processes differ from production function assumptions. For species with complex life histories like striped

marlin, managers should consider these limitations when applying MSY-based reference points derived from surplus production models.

Data limitations also constrain the analysis. Fitting to a single standardized abundance index limits evaluation of uncertainty in abundance trends. Multiple indices would enable model ensemble approaches to better characterize this uncertainty.

The incorporation of depletion information from New Zealand recreational size data through a 1988 depletion prior, while novel, represents a simplified approach. More sophisticated modeling frameworks like delay-difference or age-structured models could explicitly fit size composition time series in a likelihood-based framework, providing more elegant integration of this information.

Modeling assumptions introduce additional uncertainties. Current modeling assumes the same level of catch uncertainty across all time periods. Future versions could allow time-varying catch uncertainty (e.g., greater uncertainty in historical versus recent observations) or accommodate effort-driven removals similar to catch-errors with effort deviation approaches available in MULTIFAN-CL (Fournier et al., 1998).

Population scale estimates were highly uncertain and sensitive to model configuration choices, while depletion estimates were more precisely estimated. The choice of prior on σ_F was particularly influential. Although this prior was developed using prior pushforward analysis (Kim & Neubauer, 2025), it represents a key uncertainty. Similarly, uncertainty in biological parameters (growth, maturity, natural mortality) due to limited sample sizes contributed to broad initial priors for R_{Max} . While the posterior distribution for R_{Max} was substantially updated from the prior, reducing uncertainty in key biological productivity parameters could help reduce model uncertainty estimates and facilitate development of age-structured models.

5.3 Future stock assessment modeling considerations

Future assessment efforts should progressively build toward full age-structure through a systematic approach: delay-difference models, age-structured production models, and then fully integrated age-structured models. This development should occur within a Bayesian framework to inform parameter estimates where prior information exists or can be derived using prior pushforward approaches (Kim & Neubauer, 2025).

Developing Bayesian age-structured modeling approaches can provide more stable estimation of increased parameters due to the use of priors, explicitly account for parameter uncertainty rather than making fixed assumptions, potentially enable informed estimation of difficult-to-observe biological relationships (e.g., stock-recruit relationships), address BSPM limitations by explicitly considering age-structured processes and fishery selectivity, and estimate leading parameters with likelihood components for indices and size composition.

While simplified models offer advantages when data conflicts exist, over-simplification of age-structured processes could result in biases. Future modeling must balance complexity with data quality and conflicts. The transition should be guided by careful evaluation of whether increased model complexity is supported by available data and whether it improves the reliability of management advice.

6 Acknowledgements

The authors appreciate feedback received on earlier versions of this analysis, and are thankful to the following for their insights and thoughtful discussion which greatly improved the assessment: K. Kim (SPC), N. Davies (Te Takina Ltd.), S. Hoyle (Hoyle Consulting), R. Ahrens (PIFSC), and M. Nadon (PIFSC).

7 Declaration of generative AI use

A generative artificial intelligence (AI) assistant, Anthropic Claude Sonnet 4.0, was used to parse model code in the preparation of an initial draft of this report.

8 References

- Betancourt, M. J., & Girolami, M. (2013). *Hamiltonian Monte Carlo for Hierarchical Models*. <https://doi.org/10.48550/ARXIV.1312.0906>
- Castillo-Jordan, C., Day, J., Teears, T., Davies, N., Hampton, J., McKechnie, S., Magnusson, A., Peatman, T., Vidal, T., Williams, P., & Hamer, P. (2024). *Stock Assessment of Striped Marlin in the Southwest Pacific Ocean: 2024* (WCPFC-SC20-2024/SA-WP-03 (Ver. 2.01)). <https://meetings.wcpfc.int/node/23120>
- Castillo-Jordan, C., Day, J., Teears, T., Davies, N., Hampton, J., McKechnie, S., Magnusson, A., Peatman, T., Vidal, T., Williams, P., & Hamer, P. (2025). *Revised Stock Assessment of Striped Marlin in the Southwest Pacific Ocean: 2025* (WCPFC-SC21-2025/SA-WP-XX).
- Ducharme-Barth, N., Castillo-Jordan, C., Sculley, M., Ahrens, R., & Carvalho, J. (2025). *Summary report from the NOAA-SPC assessment model meeting on SWPO striped marlin* (WCPFC-SC21-2025/SA-IP-XX).
- Edwards, C. (2024). *Bdm: Bayesian biomass dynamics model* (R package version 0.0.0.9036).
- Farley, J., Eveson, P., Krusic-Golub, K., & Kopf, K. (2021). *Southwest Pacific Striped Marlin Population Biology (Project 99)* (WCPFC-SC-17/SA-IP-11). <https://meetings.wcpfc.int/node/12569>
- Fournier, D. A., Hampton, J., & Sibert, J. R. (1998). MULTIFAN-CL: A length-based, age-structured model for fisheries stock assessment, with application to South Pacific albacore, *Thunnus alalunga*. *Canadian Journal of Fisheries and Aquatic Sciences*, 55(9), 2105–2116. <https://doi.org/10.1139/f98-100>
- Fowler, C. W. (1988). Population dynamics as related to rate of increase per generation. *Evolutionary Ecology*, 2(3), 197–204. <https://doi.org/10.1007/BF02214283>
- Gabry, J., & Mahr, T. (2024). *Bayesplot: Plotting for Bayesian Models* (R package version 1.11.1). <https://mc-stan.org/bayesplot/>
- Gabry, J., Simpson, D., Vehtari, A., Betancourt, M., & Gelman, A. (2019). Visualization in Bayesian Workflow. *Journal of the Royal Statistical Society Series A: Statistics in Society*, 182(2), 389–402. <https://doi.org/10.1111/rssa.12378>
- Grant, P. R., & Grant, B. R. (1992). Demography and the Genetically Effective sizes of Two Populations of Darwin’s Finches. *Ecology*, 73(3), 766–784. <https://doi.org/10.2307/1940156>
- Gravel, S., Bigman, J. S., Pardo, S. A., Wong, S., & Dulvy, N. K. (2024). Metabolism, population growth, and the fast-slow life history continuum of marine fishes. *Fish and Fisheries*, 25(2), 349–361. <https://doi.org/10.1111/faf.12811>
- Hamel, O. S., & Cope, J. M. (2022). Development and considerations for application of a longevity-based prior for the natural mortality rate. *Fisheries Research*, 256, 106477. <https://doi.org/10.1016/j.fishres>

2022.106477

- Holdsworth, J. C. (2023). *Striped marlin catch and CPUE in the New Zealand sport fishery, 2019-20 to 2021-22* (New Zealand Fisheries Assessment Report 2023/05).
- Hordyk, A., Huynh, Q., & Carruthers, T. (2024). *openMSE: Easily Install and Load the 'openMSE' Packages* (R package version 1.0.1). <https://openmse.com/>
- Hurtado-Ferro, F., Szuwalski, C. S., Valero, J. L., Anderson, S. C., Cunningham, C. J., Johnson, K. F., Licandeo, R., McGilliard, C. R., Monnahan, C. C., Muradian, M. L., Ono, K., Vert-Pre, K. A., Whitten, A. R., & Punt, A. E. (2015). Looking in the rear-view mirror: Bias and retrospective patterns in integrated, age-structured stock assessment models. *ICES Journal of Marine Science*, 72(1), 99–110. <https://doi.org/10.1093/icesjms/fsu198>
- Hutchings, J. A., Myers, R. A., García, V. B., Lucifora, L. O., & Kuparinen, A. (2012). Life-history correlates of extinction risk and recovery potential. *Ecological Applications*, 22(4), 1061–1067. <https://doi.org/10.1890/11-1313.1>
- ICCAT. (2020). Report of the 2019 ICCAT white marlin stock assessment meeting. *Collect. Vol. Sci. Pap. ICCAT*, 76(4), 97–181. https://www.iccat.int/Documents/CVSP/CV076_2019/colvol76.html#
- ISC. (2024). *Stock Assessment of Shortfin Mako Shark in the North Pacific Ocean through 2022* (WCPFC-SC20-2024/SA-WP-14). <https://meetings.wcpfc.int/node/22828>
- Juárez, M. A., & Steel, M. F. J. (2010). Model-Based Clustering of Non-Gaussian Panel Data Based on Skew-*t*Distributions. *Journal of Business & Economic Statistics*, 28(1), 52–66. <https://doi.org/10.1198/jbes.2009.07145>
- Kell, L. T., Sharma, R., Kitakado, T., Winker, H., Mosqueira, I., Cardinale, M., & Fu, D. (2021). Validation of stock assessment methods: Is it me or my model talking? *ICES Journal of Marine Science*, 78(6), 2244–2255. <https://doi.org/10.1093/icesjms/fsab104>
- Kim, K., & Neubauer, P. (2025). Examining potential biases through prior predictive checks: Prior misspecifications and their impact on Bayesian stock assessments. *Fisheries Research*, 288, 107405. <https://doi.org/10.1016/j.fishres.2025.107405>
- Kokkalis, A., Berg, C. W., Kapur, M. S., Winker, H., Jacobsen, N. S., Taylor, M. H., Ichinokawa, M., Miyagawa, M., Medeiros-Leal, W., Nielsen, J. R., & Mildenerberger, T. K. (2024). Good practices for surplus production models. *Fisheries Research*, 275, 107010. <https://doi.org/10.1016/j.fishres.2024.107010>
- Methot Jr., R. D., & Wetzell, C. R. (2013). Stock synthesis: A biological and statistical framework for fish stock assessment and fishery management. *Fisheries Research*, 142, 86–99. <https://doi.org/10.1016/j.fishres.2012.10.012>
- Mohn, R. (1999). The retrospective problem in sequential population analysis: An investigation using cod fishery and simulated data. *ICES Journal of Marine Science*, 56(4), 473–488. <https://doi.org/10.1006/jmsc.1999.0481>
- Monnahan, C. C. (2024). Toward good practices for Bayesian data-rich fisheries stock assessments using a modern statistical workflow. *Fisheries Research*, 275, 107024. <https://doi.org/10.1016/j.fishres.2024.107024>
- Neubauer, P., Castillo-Jordan, C., Day, J., & Hamer, P. (2025). *Exploring the potential for observer CPUE for southwest Pacific swordfish (Xiphias gladius) and striped marlin (Kajikia audax)* (WCPFC-SC21-2025/SA-IP-13). <https://meetings.wcpfc.int/node/26539>
- Neubauer, P., Kim, K., Large, K., & Brouwer, S. (2024). *Stock Assessment of Silky Shark in the Western and Central Pacific Ocean 2024* (WCPFC-SC20-2024/SA-WP-04-Rev2). <https://meetings.wcpfc.int/node/23088>

- Neubauer, P., Richard, Y., & Tremblay-Boyer, L. (2019). *Alternative assessment methods for oceanic whitetip shark* (WCPFC-SC15-2019/SA-WP-13).
- Pardo, S. A., Kindsvater, H. K., Reynolds, J. D., & Dulvy, N. K. (2016). Maximum intrinsic rate of population increase in sharks, rays, and chimaeras: The importance of survival to maturity. *Canadian Journal of Fisheries and Aquatic Sciences*, 73(8), 1159–1163. <https://doi.org/10.1139/cjfas-2016-0069>
- Sivula, T., Magnusson, M., Matamoros, A. A., & Vehtari, A. (2020). *Uncertainty in Bayesian Leave-One-Out Cross-Validation Based Model Comparison*. arXiv. <https://doi.org/10.48550/ARXIV.2008.10296>
- Stanley, R. D., McAllister, M., Starr, P., & Olsen, N. (2009). *Stock assessment for bocaccio (Sebastes paucispinis) in British Columbia waters* (Canadian {Science} {Advisory} {Secretariat} Research Document 2009/055). Fisheries; Oceans Canada. https://www.dfo-mpo.gc.ca/csas-sccs/publications/resdocs-docrech/2009/2009_055-eng.htm
- Team, S. D. (2024a). *RStan: The R interface to Stan* (R package version 2.32.6). <https://mc-stan.org/>
- Team, S. D. (2024b). *Stan Modeling Language Users Guide and Reference Manual, v 2.32.6*.
- Tremblay-Boyer, L., & Williams, A. (2024). *Standardised cpue indices for the target species in the eastern tuna and billfish fishery–1998 to 2023* (Technical Report Working Paper presented to the 41st meeting of the Tropical Tuna Resource Assessment Group held 16-17 July 2024, Brisbane, CSIRO, Hobart).
- Vehtari, A., Gabry, J., Magnusson, M., Yao, Y., Burkner, P., Paananen, T., & Gelman, A. (2024). *Loo: Efficient leave-one-out cross-validation and WAIC for Bayesian models* (R package version 2.7.0). <https://mc-stan.org/loo/>
- Vehtari, A., Gelman, A., & Gabry, J. (2017). Practical Bayesian model evaluation using leave-one-out cross-validation and WAIC. *Statistics and Computing*, 27(5), 1413–1432. <https://doi.org/10.1007/s11222-016-9696-4>
- Winker, H., Carvalho, F., & Kapur, M. (2018). JABBA: Just Another Bayesian Biomass Assessment. *Fisheries Research*, 204, 275–288. <https://doi.org/10.1016/j.fishres.2018.03.010>
- Winker, H., Mourato, B., & Chang, Y. (2019). Unifying parameterizations between age-structured and surplus production models: An application to atlantic white marlin (*Kajiki albidia*) with simulation testing. *Collect. Vol. Sci. Pap. ICCAT*, 76(4), 219–234.

9 Tables

Table 1: Biological parameter distributions and sources used in simulation framework

Parameter	Symbol	Distribution	Mean/Mode	CV/Range	Correlation	Source/Notes
Length at age 1	L_1	Lognormal	60 cm	0.2	Independent	(Ducharme-Barth et al., 2025)
Length at age 10	L_2	Lognormal	210 cm	0.2	Independent	(Ducharme-Barth et al., 2025)
Growth coefficient	k	Beta(6.5, 3.5)	~0.65	-	Independent	(Ducharme-Barth et al., 2025)
Length CV	cv_{len}	Uniform	-	[0.05, 0.25]	-	Growth variability
Maximum age	A_{Max}	Lognormal	15 years	0.2	$\rho = -0.3$ with M_{ref}	(Farley et al., 2021)
Reference mortality	M_{ref}	Lognormal	0.36 yr^{-1}	0.44	$\rho = -0.3$ with A_{Max}	(Hamel & Cope, 2022)
Maturity slope	a_{mat}	Normal	-20	0.2	-	(5.4/ A_{Max}) Logistic steepness
Length at 50% maturity	L_{50}	Lognormal	184 cm	0.2	-	(Farley et al., 2021)
Weight coefficient	a_w	Lognormal	5.4×10^{-7}	0.05	$\rho = -0.5$ with b_w	(Castillo-Jordan et al., 2024)
Weight exponent	b_w	Normal	3.58	0.05	$\rho = -0.5$ with a_w	(Castillo-Jordan et al., 2024) Constrained: 150-350 kg at 300 cm
Steepness	h	Beta(3, 1.5)	~0.73	-	-	Censored to [0.2, 1.0]
Sex ratio (prop. female)	s	Normal	0.5	0.05	-	Constrained [0.01, 0.99]
Reproductive cycle	-	Fixed	1 year	-	-	Annual spawning
Reference ages	age_1, age_2	Fixed	0, 10 years	-	-	Francis parameterization

Table 2: Prior distributions for diagnostic model leading parameters.

Parameter	Distribution	Mean	SD	Description	Key Correlations
Core Population Parameters					
K	MV Lognormal	$\log(1,110,533)$	0.59	Carrying capacity (N)	$\rho(K, R_{Max}) = -0.22$; $\rho(K, q_{eff}) = -0.241$
R_{Max}	MV Lognormal	$\log(0.6206)$	0.58	Max rate of increase	$\rho(R_{Max}, n) = -0.881$
n	MV Lognormal	$\log(1.16)$	0.54	Pella-Tomlinson shape parameter	
x_0	MV Lognormal	$\log(1)$	0.025	Initial depletion	(Independent)
q_{eff}	MV Lognormal	$\log(0.5019)$	0.95	Initial catchability	$\rho(q_{eff}, \sigma_{qdev}) = 0.559$; $\rho(q_{eff}, \rho) = 0.243$
ρ	MV Transformed Normal	$\tanh^{-1}(0.981)$	0.62	Autocorrelation in catchability deviations	$\rho(\rho, \sigma_{qdev}) = 0.148$
σ_{qdev}	MV Lognormal	$\log(1.619)$	0.3	Catchability variability	
Process and Observation Error					
σ_P	Lognormal	$\log(0.0533)$	0.27	Process error	(Independent)
$\sigma_{O,add}$	Half-Normal	0	0.2	Additional estimated observation error	(Independent)
Catch Likelihood					
ν_{catch}	Gamma	Shape = 2	Rate = 0.1	Student-t degrees of freedom for catch	(Independent)

Notes:

Distributions with *MV* indicate that the parameter is part of the 7-dimensional multivariate prior.

Table 3: Model descriptions

Model	Group	Description
0100	Final ensemble	Diagnostic case model: DWFN index & baseline priors
0102	Final ensemble	New Zealand index & baseline priors
0105	Final ensemble	DWFN index & alternative shape prior
0107	Final ensemble	New Zealand index & alternative shape prior
0071	One-off: Index	Australia index
0072	One-off: Index	New Zealand index
0073	One-off: Index	Observer (all) index
0074	One-off: Index	Observer (New Caledonia, Fiji & Tonga) index
0075	One-off: Index	Observer (French Polynesia) index
0129	One-off: Index	DWFN & Australia index
0130	One-off: Index	New Zealand & Australia index
0131	One-off: Index	DWFN & Observer (New Caledonia, Fiji & Tonga) index
0132	One-off: Index	New Zealand & Observer (New Caledonia, Fiji & Tonga) index
0076	One-off: sigma_c	Time-varying sigma_c: Power curve from 0.5-0.2
0077	One-off: sigma_c	Constant sigma_c: 0.1
0078	One-off: sigma_c	Time-varying sigma_c: Power curve from 0.5-0.1
0079	One-off: Early data	Alternative 1952 catch
0080	One-off: Early data	Alternative 1952 effort
0081	One-off: Early data	Alternative 1954 catch
0082	One-off: Early data	Alternative 1952 & 1954 catch
0083	One-off: Early data	Alternative 1952 effort & 1954 catch
0093	One-off: Scaling priors	Force higher scale (logK & qeff)
0096	One-off: Scaling priors	Force higher scale (logK & qeff), alternative shape prior
0110	One-off: Catch obs model	Lognormal error structure
0111	One-off: qdev steps	n_step = 2
0112	One-off: qdev steps	n_step = 3
0113	One-off: qdev steps	n_step = 5
0114	One-off: qdev steps	n_step = 10
0115	One-off: Catch reporting	2x observed catch
0116	One-off: Catch reporting	Linear increase to 2x observed catch
0117	One-off: Catch reporting	Linear decrease from 2x observed catch
0118	One-off: Process error	Process error restricted to 0
0119	One-off: Start year	Start in 1955: x0 = 0.9
0120	One-off: Start year	Start in 1988: x0 = 0.5
0121	One-off: Start year	Start in 1988: x0 = 0.7
0122	One-off: Start year	Start in 1988: x0 = 0.9
0124	One-off: Est. effort devs	Estimate effort deviates & n_step = 2
0126	One-off: Est. effort devs	Estimate effort deviates & n_step = 5
0127	One-off: Direct F estimation	sigmaF to match diagnostic case
0128	One-off: Direct F estimation	0.25x diagnostic case sigmaF

Table 3: Model descriptions

Model	Group	Description
0133	One-off: Rmax prior	Apply lower productivity Rmax prior
0134	One-off: Rmax prior	Apply lower productivity Rmax prior and alternative shape prior

Table 4: Model diagnostics and fit statistics

	Model	n par	n obs	Max $\hat{R}$	Min ESS	Divergent	Tree Depth	BFMI Issues	LOOIC	p	ELPD	Prop Pareto k
										LOO	LOO	
37	0100	150	114	1.008	788	0	0	0	45.4	32.7	-22.7	0.982
	0102	150	118	1.008	688	0	0	0	64.9	32.9	-32.4	0.983
	0105	150	114	1.008	677	0	0	0	46.1	33.0	-23.0	0.982
	0107	150	118	1.007	523	0	0	0	66.1	33.3	-33.1	0.966
	0071	150	95	1.007	792	0	0	0	-1.6	35.7	0.8	0.937
	0072	150	118	1.014	610	0	0	0	65.3	33.7	-32.7	0.983
	0073	150	92	1.008	738	0	0	0	-0.7	27.5	0.4	1.000
	0074	150	92	1.008	760	0	0	0	30.2	28.9	-15.1	0.978
	0075	150	90	1.036	230	0	0	0	58.3	29.7	-29.2	0.989
	0129	150	139	1.014	666	0	0	0	24.7	36.5	-12.3	0.971
	0130	150	143	1.012	760	0	0	0	56.2	39.3	-28.1	0.986
	0131	150	136	1.005	710	0	0	0	35.9	35.5	-17.9	0.985
	0132	150	140	1.011	783	0	0	0	61.7	36.7	-30.8	0.979
	0076	150	114	1.008	644	0	0	0	49.8	19.2	-24.9	0.991
	0077	150	114	1.013	693	0	0	0	32.0	57.3	-16.0	0.991
	0078	150	114	1.010	779	0	0	0	28.6	26.9	-14.3	0.991
	0079	150	114	1.010	754	0	0	0	19.5	33.6	-9.7	0.939
	0080	150	114	1.010	773	0	0	0	20.9	34.3	-10.5	0.930
	0081	150	114	1.006	748	0	0	0	45.8	32.1	-22.9	0.982
	0082	150	114	1.009	748	0	0	0	21.3	33.4	-10.7	0.974
	0083	150	114	1.008	537	0	0	0	21.5	33.3	-10.7	0.956
	0093	150	114	1.006	749	0	0	0	52.4	35.3	-26.2	0.982
	0096	150	114	1.007	746	1	0	0	57.0	35.1	-28.5	0.974
	0110	149	114	1.008	729	0	0	0	1457.1	49.6	-728.5	0.711
	0111	115	114	1.009	747	0	0	0	43.6	30.3	-21.8	0.974
	0112	104	114	1.007	764	0	0	0	55.1	28.3	-27.5	0.974
	0113	94	114	1.007	815	0	0	0	43.6	26.9	-21.8	0.982

Table 4: Model diagnostics and fit statistics

Model	n par	n obs	Max $\hat{R}$	Min ESS	Divergent	Tree Depth	BFMI Issues	LOOIC	p	ELPD	Prop Pareto k
									LOO	LOO	
0114	87	114	1.007	745	0	0	0	53.8	27.9	-26.9	0.991
0115	150	114	1.010	764	0	0	0	44.8	32.2	-22.4	0.991
0116	150	114	1.010	730	0	0	0	45.8	32.6	-22.9	0.982
0117	150	114	1.011	688	0	0	0	42.3	31.4	-21.2	0.982
0118	150	114	1.009	581	0	0	0	52.0	29.3	-26.0	0.991
0119	147	111	1.007	701	0	0	0	5.3	7.5	-2.6	0.977
0120	114	69	1.007	771	1	0	0	3.9	5.8	-2.0	0.971
0121	114	69	1.008	612	0	0	0	5.1	6.5	-2.6	0.971
0122	114	69	1.008	737	0	0	0	6.3	6.2	-3.1	0.971
0124	185	114	1.007	658	0	0	0	65.2	53.4	-32.6	0.860
0126	164	114	1.006	771	0	0	0	61.0	48.6	-30.5	0.904
0127	149	114	1.006	700	0	0	0	60.8	60.2	-30.4	0.535
0128	149	114	1.011	702	0	0	0	62.8	57.1	-31.4	0.544
0133	150	114	1.007	742	0	0	0	45.5	31.7	-22.8	0.982
0134	150	114	1.015	762	0	0	0	44.7	31.5	-22.4	0.982

Notes:

- n par: Number of parameters in the model
- n obs: Number of observations used in model fitting
- Max $\hat{R}$: Maximum potential scale reduction factor across all parameters (should be < 1.01)
- Min ESS: Minimum effective sample size across all parameters (should be > 500)
- Divergent: Number of divergent transitions during sampling (should be 0)
- Tree Depth: Whether maximum tree depth was exceeded during sampling

- BFMI Issues: Whether low Bayesian Fraction of Missing Information was detected
- LOOIC: Leave-one-out information criterion (lower is better)
- p LOO: Effective number of parameters estimated by LOO (should be $< n_{\text{par}} \& n_{\text{obs}}$)
- ELPD LOO: Expected log pointwise predictive density from LOO (higher is better)
- Prop Pareto k: Proportion of Pareto k values ≤ 0.7 (should be high for reliable LOO estimates)

Table 5: Model performance and validation statistics

Model	Index RMSE	Catch RMSE	Mohn's ρ	MASE	Coverage D/D_{MSY}	Coverage U/U_{MSY}
0100	0.257	0.250	-0.087	0.587	100%	100%
0102	0.317	0.251	0.069	2.577	100%	100%
0105	0.258	0.250	-0.099	0.566	100%	100%
0107	0.317	0.256	0.087	2.579	100%	100%
0071	0.086	0.246	NA	NA	NA%	NA%
0072	0.316	0.250	NA	NA	NA%	NA%
0073	0.064	0.245	NA	NA	NA%	NA%
0074	0.188	0.244	NA	NA	NA%	NA%
0075	0.390	0.251	NA	NA	NA%	NA%
0129	0.189	0.248	NA	NA	NA%	NA%
0130	0.231	0.247	NA	NA	NA%	NA%
0131	0.230	0.251	NA	NA	NA%	NA%
0132	0.271	0.249	NA	NA	NA%	NA%
0076	0.261	0.335	NA	NA	NA%	NA%
0077	0.253	0.210	NA	NA	NA%	NA%
0078	0.256	0.307	NA	NA	NA%	NA%
0079	0.256	0.222	NA	NA	NA%	NA%
0080	0.256	0.223	NA	NA	NA%	NA%
0081	0.259	0.240	NA	NA	NA%	NA%
0082	0.256	0.221	NA	NA	NA%	NA%
0083	0.256	0.221	NA	NA	NA%	NA%
0093	0.275	0.247	NA	NA	NA%	NA%
0096	0.298	0.249	NA	NA	NA%	NA%
0110	0.250	0.201	NA	NA	NA%	NA%
0111	0.255	0.256	NA	NA	NA%	NA%
0112	0.260	0.297	NA	NA	NA%	NA%
0113	0.249	0.273	NA	NA	NA%	NA%
0114	0.239	0.316	NA	NA	NA%	NA%
0115	0.256	0.254	NA	NA	NA%	NA%
0116	0.258	0.228	NA	NA	NA%	NA%
0117	0.251	0.282	NA	NA	NA%	NA%
0118	0.281	0.258	NA	NA	NA%	NA%
0119	0.255	0.218	NA	NA	NA%	NA%
0120	0.272	0.170	NA	NA	NA%	NA%
0121	0.272	0.170	NA	NA	NA%	NA%
0122	0.279	0.171	NA	NA	NA%	NA%
0124	0.249	0.234	NA	NA	NA%	NA%
0126	0.245	0.238	NA	NA	NA%	NA%
0127	0.247	0.221	NA	NA	NA%	NA%

Table 5: Model performance and validation statistics

Model	Index RMSE	Catch RMSE	Mohn's ρ	MASE	Coverage D/D_{MSY}	Coverage U/U_{MSY}
0128	0.268	0.221	NA	NA	NA%	NA%
0133	0.260	0.252	NA	NA	NA%	NA%
0134	0.259	0.250	NA	NA	NA%	NA%

Notes:

- Index RMSE: Root mean squared error for fit to standardized CPUE index (lower is better)
- Catch RMSE: Root mean squared error for fit to catch data (lower is better)
- Mohn's ρ : Retrospective bias statistic (values close to 0 indicate low bias)
- MASE: Mean Absolute Scaled Error from hindcast cross-validation (< 1 indicates model outperforms naïve predictor)
- Coverage D/D_{MSY} : Percentage of retrospective peels where relative depletion estimates fall within full model credible intervals
- Coverage U/U_{MSY} : Percentage of retrospective peels where relative exploitation rate estimates fall within full model credible intervals

Table 6: Posterior estimates (median and 95% credible intervals) of the leading model parameters for models in the final ensemble.

Model	$\log(K)$	R_{Max}	n	x_0	q_{eff}	ρ	σ_{qdev}	σ_P	$\sigma_{O,add}$	ν_{catch}	σ_F
0100	12.786 (12.323, 14.046)	0.396 (0.155, 0.866)	1.353 (0.500, 3.544)	1.000 (0.948, 1.053)	2.227 (0.602, 8.505)	0.993 (0.980, 0.997)	1.495 (1.013, 2.211)	0.058 (0.034, 0.094)	0.084 (0.029, 0.154)	3.797 (2.119, 7.667)	—
0102	12.631 (12.142, 14.378)	0.543 (0.185, 1.168)	1.090 (0.446, 3.042)	1.000 (0.952, 1.051)	2.502 (0.513, 9.714)	0.993 (0.979, 0.997)	1.478 (1.016, 2.210)	0.069 (0.038, 0.115)	0.184 (0.100, 0.291)	3.664 (1.967, 7.697)	—
0105	12.751 (12.198, 14.209)	0.322 (0.121, 0.722)	2.913 (1.172, 8.168)	1.002 (0.950, 1.045)	2.356 (0.596, 8.655)	0.993 (0.979, 0.997)	1.547 (1.041, 2.191)	0.059 (0.036, 0.101)	0.083 (0.029, 0.160)	3.747 (2.000, 7.582)	—
0107	12.571 (12.015, 14.428)	0.438 (0.153, 1.008)	2.351 (0.899, 6.335)	1.000 (0.954, 1.052)	2.673 (0.565, 9.271)	0.993 (0.980, 0.997)	1.503 (1.006, 2.147)	0.067 (0.038, 0.112)	0.184 (0.103, 0.291)	3.746 (1.988, 7.349)	—

10 Figures




Figure 1: Annual catch (numbers; individuals) of striped marlin and nominal longline effort (hooks fished; thousands) in the Southwest Pacific Ocean (1952-2022).




Figure 2: Standardized index (black points), and observation error (black bars) for alternative model indices. The trend in the indices is shown in blue using a loess-smooth.




Figure 3: Prior pushforward check showing simulated removals and population depletion trajectories under different filtering scenarios (median: colored lines) with 95% quantile of trajectories (shaded polygon).




Figure 4: Prior pushforward diagnostic pairs plot showing parameter correlations and marginal distributions for different filtering scenarios. The plot compares initial priors (blue), catch-filtered priors (cyan), cpue-filtered priors (lime green), and the final updated priors following fitting to the cpue-filtered distribution (orange) across key model parameters (logK, r, shape, x0, log_qeff, atan_rho, sigma_qdev). Diagonal panels show marginal density distributions with overlaid curves for each filtering scenario. Off-diagonal scatter plots display pairwise parameter relationships colored by filtering type. Correlation coefficients in the upper triangle quantify linear relationships between adjacent distributions, with statistical significance indicated by asterisks.




Figure 5: Model fit to catch data showing observed catch (black points with 95% error bars) and model-predicted catch (colored lines and shaded ribbons representing 95% credible intervals).




Figure 6: Posterior predictive check for catch data using probability integral transform empirical cumulative distribution function (PIT-ECDF). The solid blue line shows the empirical CDF of PIT values calculated from observed catch data relative to posterior predictive distributions. Light blue dashed lines indicate 99% simultaneous confidence bands under the assumption of correct model specification. The solid blue line should be contained within the 99% simultaneous confidence bands.




Figure 7: Model fit to the standardized index data showing observations (black points with 95% error bars) and model-predictions (colored lines and shaded ribbons representing 95% credible intervals).




Figure 8: Probability Integral Transform (PIT) residuals for catch data from the diagnostic model over time. Orange points with vertical lines show PIT residual values for each year, where values should be uniformly distributed between 0 and 1 under correct model specification. The ‘Runs test: Pass’ annotation indicates that the runs test for temporal autocorrelation in the residuals passed at the 5% significance level, suggesting no systematic patterns in the residuals over time.




Figure 9: Posterior predictive check for index data using probability integral transform empirical cumulative distribution function (PIT-ECDF). The solid blue line shows the empirical CDF of PIT values calculated from observed catch data relative to posterior predictive distributions. Light blue dashed lines indicate 99% simultaneous confidence bands under the assumption of correct model specification. The solid blue line should be contained within the 99% simultaneous confidence bands.




Figure 10: Retrospective analysis for the diagnostic case model (0100) with respect to time series of depletion D_t , depletion relative to depletion at MSY D_t/D_{MSY} , fishing mortality F_t , and fishing mortality relative to the rate of fishing mortality that produces MSY F_t/F_{MSY} . The base model with data included through 2022 (black line – median; dark shading – 50% credible interval; light shading – 95% credible interval) is shown relative to the retrospective models. Colored lines correspond to the last year of index data and the colored point indicates the estimate in the last year of the retrospective peel.




Figure 11: Standardized indices of relative abundance used in the diagnostic case model (0100). Open circles show observed values (standardized to mean of 1; black horizontal line) and the vertical bars indicate the observation error (95% confidence interval). One year ahead ‘model-free’ hindcast predictions are shown by the colored lines, where the color indicates the last year of index data seen by the model. The predicted value is shown one year-ahead with the colored point.




Figure 12: Prior-posterior comparison pairs plot for the diagnostic case model showing naïve prior (blue), realized prior (green), and posterior (orange) distributions across leading model parameters. Diagonal panels display marginal density distributions for each parameter. Off-diagonal scatter plots show pairwise parameter relationships colored by distribution type. Upper triangle correlation coefficients quantify linear relationships between distributions, with statistical significance indicated by asterisks. The comparison illustrates sequential parameter refinement from initial biological/expert knowledge (naïve prior) through model structure filtering (realized prior) to data-based learning (posterior).




Figure 13: Distributions of biological parameters (maximum age max_age , natural mortality reference M_ref , length at birth $L1$, asymptotic length $L2$, von Bertalanffy growth coefficient vbk , length at 50% maturity $l50$, sex ratio at birth sex_ratio , coefficient of variation in length cv_len , weight-length relationship parameters $weight_a$ and $weight_b$, and steepness h) used in the rmax sub-sampling process. Original biological prior distribution (blue shading), input prior distribution following the prior pushforward analysis (green shading), and final sub-sampled distribution selected to match the posterior R_{Max} distribution (orange shading).




Figure 14: Model-free hindcast performance for the DWFN CPUE index. The diagnostic model fitted to the full dataset (orange line and shading) is compared against retrospective models excluding the final 5, 10, 15, and 20 years of index data. Colored lines show model predictions with 95% credible intervals (shading). Observed index values are shown as black points with error bars. Colored points at the bottom indicate the terminal year of each retrospective peel: 2017 (yellow), 2012 (green), 2007 (cyan), and 2002 (blue).




Figure 15: Model-free hindcast performance for key population dynamics parameters. Time series show posterior distributions (median and 95% credible intervals) for depletion (D), population (P), fishing mortality (F), relative depletion (D/D_{MSY}), relative fishing mortality (F/F_{MSY}), removals, process error, nominal CPUE, and catchability deviates. The diagnostic model (orange) is compared against retrospective models terminating in 2017 (yellow), 2012 (green), 2007 (cyan), and 2002 (blue). Colored points indicate the terminal year of each retrospective peel.




Figure 16: One-off sensitivity group: [Index](#). Model fit to standardized index data showing observations (black points with 95% error bars) and model-predictions (colored lines and shaded ribbons representing 95% credible intervals). The diagnostic model ([0100](#)) is shown in orange.




Figure 17: One-off sensitivity group: [Index](#). Prior and posterior parameter distributions showing the evolution from prior (dotted line) to posterior (shaded) distributions for leading model parameters: carrying capacity ($\log K$), intrinsic growth rate (r), observation error (σ_{ao_add}), process error (σ_{map}), shape parameter, effective catchability (q_{eff}), autocorrelation in catchability (ρ), catchability deviation variability (σ_{qdev}), initial depletion (x_0), and catch likelihood degrees of freedom (ν_{catch}). The diagnostic model ([0100](#)) is shown in orange.




Figure 18: One-off sensitivity group: [Index](#) . Posterior time series distributions for key derived quantities over time (line = median, shading = 95% credible interval): depletion (D), absolute population size in numbers (P), fishing mortality (F), stock status relative to MSY reference points (D/D_{MSY} , F/F_{MSY}), total removals in numbers, process error, nominal CPUE (numbers caught per 1000 hooks), and catchability deviates. The diagnostic model (0100) is shown in orange.




Figure 19: One-off sensitivity group: **Catch observation error**. Prior and posterior parameter distributions showing the evolution from prior (dotted line) to posterior (shaded) distributions for leading model parameters: carrying capacity ($\log K$), intrinsic growth rate (r), observation error (σ_{ao_add}), process error (σ_{map}), shape parameter, effective catchability (q_{eff}), autocorrelation in catchability (ρ), catchability deviation variability (σ_{qdev}), initial depletion (x_0), and catch likelihood degrees of freedom (ν_{catch}). The diagnostic model (0100) is shown in orange.




Figure 20: One-off sensitivity group: **Catch observation error**. Posterior time series distributions for key derived quantities over time (line = median, shading = 95% credible interval): depletion (D), absolute population size in numbers (P), fishing mortality (F), stock status relative to MSY reference points (D/D_{MSY} , F/F_{MSY}), total removals in numbers, process error, nominal CPUE (numbers caught per 1000 hooks), and catchability deviates. The diagnostic model (**0100**) is shown in orange.




Figure 21: One-off sensitivity group: **Early data**. Prior and posterior parameter distributions showing the evolution from prior (dotted line) to posterior (shaded) distributions for leading model parameters: carrying capacity ($\log K$), intrinsic growth rate (r), observation error (σ_{ao_add}), process error (σ_{map}), shape parameter, effective catchability (q_{eff}), autocorrelation in catchability (ρ), catchability deviation variability (σ_{qdev}), initial depletion (x_0), and catch likelihood degrees of freedom (ν_{catch}). The diagnostic model (**0100**) is shown in orange.




Figure 22: One-off sensitivity group: **Early data**. Posterior time series distributions for key derived quantities over time (line = median, shading = 95% credible interval): depletion (D), absolute population size in numbers (P), fishing mortality (F), stock status relative to MSY reference points (D/D_{MSY} , F/F_{MSY}), total removals in numbers, process error, nominal CPUE (numbers caught per 1000 hooks), and catchability deviates. The diagnostic model (0100) is shown in orange.




Figure 23: One-off sensitivity group: **Scaling priors**. Prior and posterior parameter distributions showing the evolution from prior (dotted line) to posterior (shaded) distributions for leading model parameters: carrying capacity ($\log K$), intrinsic growth rate (r), observation error ($\sigma_{\text{ao_add}}$), process error (σ_{map}), shape parameter, effective catchability (q_{eff}), autocorrelation in catchability (ρ), catchability deviation variability (σ_{qdev}), initial depletion (x_0), and catch likelihood degrees of freedom (ν_{catch}). The diagnostic model (**0100**) is shown in orange.




Figure 24: One-off sensitivity group: **Scaling priors**. Model fit to standardized index data showing observations (black points with 95% error bars) and model-predictions (colored lines and shaded ribbons representing 95% credible intervals). The diagnostic model (**0100**) is shown in orange.




Figure 25: One-off sensitivity group: **Catch observation model**. Prior and posterior parameter distributions showing the evolution from prior (dotted line) to posterior (shaded) distributions for leading model parameters: carrying capacity ($\log K$), intrinsic growth rate (r), observation error (σ_{ao_add}), process error (σ_{map}), shape parameter, effective catchability (q_{eff}), autocorrelation in catchability (ρ), catchability deviation variability (σ_{qdev}), initial depletion (x_0), and catch likelihood degrees of freedom (ν_{catch}). The diagnostic model (0100) is shown in orange.




Figure 26: One-off sensitivity group: [Catch observation model](#). Posterior time series distributions for key derived quantities over time (line = median, shading = 95% credible interval): depletion (D), absolute population size in numbers (P), fishing mortality (F), stock status relative to MSY reference points (D/D_{MSY} , F/F_{MSY}), total removals in numbers, process error, nominal CPUE (numbers caught per 1000 hooks), and catchability deviates. The diagnostic model ([0100](#)) is shown in orange.




Figure 27: One-off sensitivity group: [Temporal resolution of catchability deviates](#). Model fit to catch data showing observed catch (black points with 95% error bars) and model-predicted catch (colored lines and shaded ribbons representing 95% credible intervals). The diagnostic model ([0100](#)) is shown in orange.




Figure 28: One-off sensitivity group: **Temporal resolution of catchability deviates**. Prior and posterior parameter distributions showing the evolution from prior (dotted line) to posterior (shaded) distributions for leading model parameters: carrying capacity ($\log K$), intrinsic growth rate (r), observation error ($\sigma_{\text{mao_add}}$), process error (σ_{map}), shape parameter, effective catchability (q_{eff}), autocorrelation in catchability (ρ), catchability deviation variability (σ_{qdev}), initial depletion (x_0), and catch likelihood degrees of freedom (ν_{catch}). The diagnostic model (0100) is shown in orange.




Figure 29: One-off sensitivity group: **Temporal resolution of catchability deviates**. Posterior time series distributions for key derived quantities over time (line = median, shading = 95% credible interval): depletion (D), absolute population size in numbers (P), fishing mortality (F), stock status relative to MSY reference points (D/D_{MSY} , F/F_{MSY}), total removals in numbers, process error, nominal CPUE (numbers caught per 1000 hooks), and catchability deviates. The diagnostic model (0100) is shown in orange.




Figure 30: One-off sensitivity group: **Catch reporting**. Prior and posterior parameter distributions showing the evolution from prior (dotted line) to posterior (shaded) distributions for leading model parameters: carrying capacity ($\log K$), intrinsic growth rate (r), observation error (σ_{ao_add}), process error (σ_{map}), shape parameter, effective catchability (q_{eff}), autocorrelation in catchability (ρ), catchability deviation variability (σ_{qdev}), initial depletion (x_0), and catch likelihood degrees of freedom (ν_{catch}). The diagnostic model (0100) is shown in orange.




Figure 31: One-off sensitivity group: [Catch reporting](#). Posterior time series distributions for key derived quantities over time (line = median, shading = 95% credible interval): depletion (D), absolute population size in numbers (P), fishing mortality (F), stock status relative to MSY reference points (D/D_{MSY} , F/F_{MSY}), total removals in numbers, process error, nominal CPUE (numbers caught per 1000 hooks), and catchability deviates. The diagnostic model (0100) is shown in orange.




Figure 32: One-off sensitivity group: **Process error** . Model fit to standardized index data showing observations (black points with 95% error bars) and model-predictions (colored lines and shaded ribbons representing 95% credible intervals). The diagnostic model (**0100**) is shown in orange.




Figure 33: One-off sensitivity group: **Process error**. Prior and posterior parameter distributions showing the evolution from prior (dotted line) to posterior (shaded) distributions for leading model parameters: carrying capacity ($\log K$), intrinsic growth rate (r), observation error (σ_{ao_add}), process error (σ_{map}), shape parameter, effective catchability (q_{eff}), autocorrelation in catchability (ρ), catchability deviation variability (σ_{qdev}), initial depletion (x_0), and catch likelihood degrees of freedom (ν_{catch}). The diagnostic model (0100) is shown in orange.




Figure 34: One-off sensitivity group: **Process error**. Posterior time series distributions for key derived quantities over time (line = median, shading = 95% credible interval): depletion (D), absolute population size in numbers (P), fishing mortality (F), stock status relative to MSY reference points (D/D_{MSY} , F/F_{MSY}), total removals in numbers, process error, nominal CPUE (numbers caught per 1000 hooks), and catchability deviates. The diagnostic model (**0100**) is shown in orange.




Figure 35: One-off sensitivity group: **Start year** . Posterior time series distributions for key derived quantities over time (line = median, shading = 95% credible interval): depletion (D), absolute population size in numbers (P), fishing mortality (F), stock status relative to MSY reference points (D/D_{MSY} , F/F_{MSY}), total removals in numbers, process error, nominal CPUE (numbers caught per 1000 hooks), and catchability deviates. The diagnostic model (**0100**) is shown in orange.




Figure 36: One-off sensitivity group: **Start year**. Prior and posterior parameter distributions showing the evolution from prior (dotted line) to posterior (shaded) distributions for leading model parameters: carrying capacity ($\log K$), intrinsic growth rate (r), observation error (σ_{ao_add}), process error (σ_{map}), shape parameter, effective catchability (q_{eff}), autocorrelation in catchability (ρ), catchability deviation variability (σ_{qdev}), initial depletion (x_0), and catch likelihood degrees of freedom (ν_{catch}). The diagnostic model (0100) is shown in orange.




Figure 37: One-off sensitivity group: [Inclusion of effort deviates](#). Model fit to catch data showing observed catch (black points with 95% error bars) and model-predicted catch (colored lines and shaded ribbons representing 95% credible intervals). The diagnostic model ([0100](#)) is shown in orange.




Figure 38: One-off sensitivity group: **Inclusion of effort deviates**. Prior and posterior parameter distributions showing the evolution from prior (dotted line) to posterior (shaded) distributions for leading model parameters: carrying capacity ($\log K$), intrinsic growth rate (r), observation error (σ_{ao_add}), process error (σ_{map}), shape parameter, effective catchability (q_{eff}), autocorrelation in catchability (ρ), catchability deviation variability (σ_{qdev}), initial depletion (x_0), and catch likelihood degrees of freedom (ν_{catch}). The diagnostic model (**0100**) is shown in orange.




Figure 39: One-off sensitivity group: [Inclusion of effort deviates](#). Posterior time series distributions for key derived quantities over time (line = median, shading = 95% credible interval): depletion (D), absolute population size in numbers (P), fishing mortality (F), stock status relative to MSY reference points (D/D_{MSY} , F/F_{MSY}), total removals in numbers, process error, nominal CPUE (numbers caught per 1000 hooks), and catchability deviates. The diagnostic model (0100) is shown in orange.




Figure 40: One-off sensitivity group: [Direct estimation of fishing mortality](#) . Model fit to catch data showing observed catch (black points with 95% error bars) and model-predicted catch (colored lines and shaded ribbons representing 95% credible intervals). The diagnostic model ([0100](#)) is shown in orange.




Figure 41: One-off sensitivity group: [Direct estimation of fishing mortality](#). Posterior time series distributions for key derived quantities over time (line = median, shading = 95% credible interval): depletion (D), absolute population size in numbers (P), fishing mortality (F), stock status relative to MSY reference points (D/D_{MSY} , F/F_{MSY}), total removals in numbers, process error, nominal CPUE (numbers caught per 1000 hooks), and catchability deviates. The diagnostic model (0100) is shown in orange.




Figure 42: One-off sensitivity group: **Alternative RMax prior**. Posterior time series distributions for key derived quantities over time (line = median, shading = 95% credible interval): depletion (D), absolute population size in numbers (P), fishing mortality (F), stock status relative to MSY reference points (D/D_{MSY} , F/F_{MSY}), total removals in numbers, process error, nominal CPUE (numbers caught per 1000 hooks), and catchability deviates. The diagnostic model (0100) is shown in orange.




Figure 43: **Model ensemble**. Model fit to catch data showing observed catch (black points with 95% error bars) and model-predicted catch (colored lines and shaded ribbons representing 95% credible intervals). The diagnostic model (0100) is shown in orange.




Figure 44: **Model ensemble..** Model fit to standardized index data showing observations (black points with 95% error bars) and model-predictions (colored lines and shaded ribbons representing 95% credible intervals). The diagnostic model (0100) is shown in orange.




Figure 45: Retrospective analysis for the four ensemble models with respect to time series of depletion D_t , depletion relative to depletion at MSY D_t/D_{MSY} , fishing mortality F_t , and fishing mortality relative to the rate of fishing mortality that produces MSY F_t/F_{MSY} . The base model with data included through 2022 (black line – median; dark shading – 50% credible interval; light shading – 95% credible interval) is shown relative to the retrospective models. Colored lines correspond to the last year of index data and the colored point indicates the estimate in the last year of the retrospective peel.




Figure 46: Standardized indices of relative abundance used in the model ensemble. Open circles show observed values (standardized to mean of 1; black horizontal line) and the vertical bars indicate the observation error (95% confidence interval). One year ahead 'model-free' hindcast predictions are shown by the colored lines, where the color indicates the last year of index data seen by the model. The predicted value is shown one year-ahead with the colored point.




Figure 47: **Model ensemble**. Posterior time series distributions for key derived quantities over time (line = median, shading = 95% credible interval): depletion (D), absolute population size in numbers (P), fishing mortality (F), stock status relative to MSY reference points (D/D_{MSY} , F/F_{MSY}), total removals in numbers, process error, nominal CPUE (numbers caught per 1000 hooks), and catchability deviates. The model ensemble is shown in orange.

11 Technical Annex

Additional technical detail of population dynamics components used to solve the Euler-Lotka equation for R_{Max} .

11.1 Equilibrium Age-Structured Population Dynamics

11.1.1 Survival

Survival schedule was calculated assuming constant natural mortality:

$$l_a = \begin{cases} 1 & \text{if } a = 1 \\ l_{a-1} \times \exp(-M_{ref}) & \text{if } 1 < a < A_{Max} \\ \frac{l_{A_{Max}-1}}{1 - \exp(-M_{ref})} & \text{if } a = A_{Max} \text{ (plus group)} \end{cases}$$

11.1.2 Fecundity

Reproductive output incorporated biological constraints:

$$f_a = \frac{\psi_a \times W_a \times s}{\rho}$$

where: - ψ_a = proportion mature at age a - W_a = weight at age a (proxy for reproductive capacity) - s = sex ratio (proportion female) - ρ = reproductive cycle (years between spawning events)

Maturity-at-age was derived from length-based logistic maturity:

$$\psi_L = \frac{\exp(a_{mat} + b_{mat} \times L)}{1 + \exp(a_{mat} + b_{mat} \times L)}$$

where $b_{mat} = -a_{mat}/L_{50}$, then integrated over the length-at-age distribution to obtain ψ_a .

Length-at-age followed the Francis parameterization:

$$L_a = L_1 + (L_2 - L_1) \times \frac{1 - \exp(-k(a - age_1))}{1 - \exp(-k(age_2 - age_1))}$$

with length variability modeled using probability density functions linking length bins to ages.

Weight-at-age used the allometric relationship:

$$W_a = a_w \times L_a^{b_w}$$

11.1.3 Stock-Recruitment Relationship

The slope at the origin of the stock-recruitment curve was:

$$\alpha = \frac{4h}{(1-h)\phi_0}$$

The calculation incorporated steepness (h) through the Beverton-Holt stock-recruitment relationship. Spawners-per-recruit in the unfished state was:

$$\phi_0 = \sum_{a=1}^{A_{Max}} l_a f_a$$

This α parameter links the intrinsic rate of increase to recruitment compensation, ensuring that R_{Max} estimates reflect realistic population productivity under the assumed stock-recruitment dynamics.

Successful simulations (those yielding $R_{Max} > 0$ and $R_{Max} < 1.5$) were filtered to create a plausible prior distribution. The resulting distribution was fitted to a lognormal prior for use in the BSPM.