



**SCIENTIFIC COMMITTEE
TWENTY-FIRST REGULAR SESSION**

Nuku'alofa, Tonga
13-21 August 2025

2025 Stock Assessment of Striped Marlin in the Southwest Pacific Ocean: Part II Data-moderate approach

**WCPFC-SC21-2025/SA-WP-07
16 July 2025**

N. Ducharme-Barth¹, C. Castillo-Jordán², P. Hamer²

¹NOAA Fisheries, Pacific Islands Fisheries Science Center, 1845 Wasp Boulevard, Bldg. 176, Honolulu HI 96818

²Oceanic Fisheries Programme of the Pacific Community

Table of contents

1	Executive summary	4
2	Introduction	5
3	Methods	6
3.1	Model framework	6
3.1.1	Input data	7
3.1.2	Population dynamics	7
3.1.3	Observation model	9
3.2	Prior development	9
3.2.1	Biological simulation framework	10
3.2.2	Fishing data simulation framework	11
3.2.3	Other priors	12
3.2.4	Prior pushforward	12
3.3	Model implementation	13
3.4	Model diagnostics	13
3.4.1	Convergence diagnostics	14
3.4.2	Leave-one-out cross-validation diagnostics	14
3.4.3	Data fits	14
3.4.4	Posterior predictive checks	15
3.4.5	Retrospective analysis	15
3.4.6	Hindcast cross-validation	15
3.5	Forecasts	16
4	Results	16
4.1	Diagnostic case	16
4.1.1	Model convergence and diagnostics	16
4.1.2	Population dynamics and stock status	17
4.2	One-off sensitivities	18
4.3	Model ensemble	18
4.3.1	Population dynamics and stock status	18
4.3.2	Biological reference points	18
4.3.3	Future projections	18
5	Discussion	19
5.1	General remarks	19
5.2	Challenges, limitations & key uncertainties	19
5.3	Future stock assessment modeling considerations	20
6	Acknowledgements	20
7	Declaration of generative AI use	20
8	References	20
9	Tables	23
10	Figures	27
11	Technical Annex	44
11.1	Equilibrium Age-Structured Population Dynamics	44
11.1.1	Survival	44

11.1.2 Fecundity	45
11.1.3 Stock-Recruitment Relationship	45

1 Executive summary

2 Introduction

Assessments of Southwest Pacific Ocean (SWPO) striped marlin (*Kajikia audax*) have been challenging. As documented in the joint NOAA-SPC modeling meeting report (Ducharme-Barth et al., 2025), key issues identified by WCPFC SC20 included poor fits to size composition data and relative abundance indices, conflicts between different data sources, and difficulties in estimating model initial conditions.

Discussions at the 2025 SPC Pre-Assessment Workshop highlighted additional challenges related to the estimation of total population scale. The low biomass scale indicated by previous versions of the assessment appears largely driven by the size composition from multiple fisheries. While estimation of low biomass scale is not inherently problematic, the resulting high ratio of fishing mortality (F) to natural mortality (M) that is maintained over multiple decades in the face of relative stability in catches and catch-per-unit-effort (CPUE) was noted as suspicious and flagged for further investigation.

The complexity of integrated age-structured models, while providing detailed population dynamics representation, can become problematic when fundamental data conflicts exist or when key biological parameters are uncertain. For SWPO striped marlin, these challenges are compounded by spatiotemporal heterogeneity in fleet coverage, potential non-representativeness in mixed-fleet composition data, and limitations in age data from opportunistic sampling. Changes to productivity assumptions (growth, natural mortality, maturity, and/or steepness) or selectivity patterns will impact biomass scaling through predictions of expected size composition data.

Scaling back model complexity to a data-moderate approach, like Bayesian surplus production models (BSPMs), offers analysts a simplified yet robust alternative for stock assessment when data limitations or conflicts present challenges for more complex models. These models have proven effective in recent pelagic fish stock assessments (Winker et al., 2018), and are routinely applied for WCPFC shark assessments (ISC, 2024; Neubauer et al., 2019, 2024). They facilitate a more tractable exploration of model assumptions by distilling productivity and fishing assumptions into a restricted parameter subset. The BSPM framework allows for explicit incorporation of parameter uncertainty through informative priors while maintaining computational efficiency and interpretability.

One advantage of the Bayesian approach is the use of prior information to propagate uncertainty in model assumptions. Biological simulation approaches can provide robust methods for parameterizing key population dynamics parameters (ISC, 2024; Pardo et al., 2016). For species with complex life histories like striped marlin, these simulation-based priors can incorporate uncertainty in growth, maturity, natural mortality, and reproductive parameters to derive realistic distributions for maximum intrinsic rate of increase and carrying capacity. Additionally, prior pushforward checks (Kim & Neubauer, 2025; Monnahan, 2024) can be used to further refine parameter distributions by identifying parameter combinations that produce clearly improbable outcomes.

This analysis presents a data-moderate approach for SWPO striped marlin that addresses some of the key technical challenges identified in previous assessments while providing a robust framework for stock status evaluation. The model incorporates biological uncertainty through simulation-based priors and includes approaches to address catch uncertainty and population depletion. It is important to note that this analysis complements the 2025 revised stock assessment report (Castillo-Jordan et al., 2025). While the relative simplicity of data-moderate approaches offers some advantages relative to more data-intensive approaches

like fully-integrated age-structured stock assessment models, particularly when uncertainty in data or key assumptions is high, this comes at the cost of simplifying the model fishery and population dynamics. Over-simplification of key age-structured processes, or lack of spatiotemporal specificity could result in biases in data-moderate approaches. Taking multiple modeling approaches into account can give managers a holistic view on the likely status and trajectory of the stock.

3 Methods

3.1 Model framework

To address some of the issues raised in Section 2, a series of Bayesian state-space surplus production models (BSPMs) spanning the period 1952-2022 were developed following the Fletcher-Schaefer production model framework (Edwards, 2024; Winker et al., 2020). Development of the BSPM followed the approach of (ISC, 2024; Neubauer et al., 2019) and incorporated recent best practices for surplus production models (Kokkalis et al., 2024) and Bayesian workflows for stock assessment (Monnahan, 2024) as guides for model development, analysis, and presentation.

The models were implemented in the Stan probabilistic programming language (Team, 2024b) to take advantage of enhanced convergence diagnostics, greater efficiency in posterior sampling through the no-U-turn (NUTS) Hamiltonian Monte Carlo (HMC) algorithm (Betancourt & Girolami, 2013), and greater flexibility with model configuration and prior specification. Implementation in R using the *rstan* package (Team, 2024a) provides connection to an ecosystem of R packages for model diagnostics and validation: *bayesplot* (Gabry & Mahr, 2024) for posterior visualization and *loo* (Vehtari et al., 2024).

In their baseline configuration (diagnostic case), the models began from an assumed unfished state in 1952 and incorporate process error in population dynamics, observation error in the relative abundance index and catch time series, and uncertainty in biological parameters. Population scale or carrying capacity $\log(K)$ is given in the units of the catch which is total numbers. The models fit to a single time series of relative abundance (e.g., standardized CPUE index) at a time. Rather than subtracting the catch directly, the fishing mortality necessary to produce the catch was estimated internally by fitting to a time series of observed catches. Computationally, this was necessary to avoid placing hard constraints on the posterior parameter space which can cause convergence issues for the HMC sampling algorithm. Additionally, rather than directly estimating the independent fishing mortality values required to produce the catch, fishing mortality was derived from an estimated time-varying catchability trend and a time series of observed longline fishing effort. The primary rationale for this modelling choice being that it is easier to develop a defensible prior for catchability rather than an arbitrary prior for fishing mortality, given that previous work (ISC, 2024) showed a strong correlation between estimated population scale and the fishing mortality prior. Unless otherwise mentioned, all models used the same prior parameterizations (Table 2) based on the biological simulation filtering described in Section 3.2.

Using the diagnostic model¹ as a starting point, a number of one-off sensitivity analyses were conducted to explore key model assumptions or data questions. These included:

- the choice of relative abundance index used to drive the model dynamics,

¹The Stan code for the diagnostic model `bspm_estqsimple_softdep_fullmvprior_x0_sttgamma_flexsigmaC_OPT` is publicly available on [GitHub](#).

- the likelihood used to model catch observation error,
- the magnitude and trend in catch observation error,
- sensitivity to assumptions in data errors for early model (1952-1954) catch and effort observations,
- alternative catch scenarios for under-reporting,
- structural assumptions related to the temporal resolution of time-varying catchability,
- complexity of the fishing mortality parametrization (e.g., direct estimation, temporal structure of catchability deviates and inclusion of effort deviates),
- model start year,
- and alternative priors assumptions (e.g., process error, catchability, population carrying capacity, and production function shape parameter).

Based on these results a structural uncertainty grid was developed for the provision of stock status and management reference points. Full model development code and input data are available online at the following GitHub repository: <https://github.com/N-DucharmeBarth-NOAA/2025-swpo-mls-bspm>.

3.1.1 Input data

The BSPMs were fitted to catch and standardized CPUE data for SWPO striped marlin. Effort was used as an input to drive the time series of estimated fishing mortality via the estimated catchability trend.

Annual catch data (in numbers of individuals) spanning 1952-2022 were compiled from the 2024 assessment (Castillo-Jordan et al., 2024) sources and aggregated into total removals by year. The catch series (Figure 1) shows initially low removals in 1952-1953, followed by a high peak in 1954. Overall, however, catches have been relatively stable though catches in recent decades have shown a slight decline.

Annual longline effort (in numbers of hooks) matching the SWPO assessment domain, as specified in Castillo-Jordan et al. (2024), was extracted from publicly available WCPFC data. Longline effort in the assessment domain (Figure 1) increased gradually from zero in the early 1950s to ~100 million hooks per year in the early 1970s where it stabilized between 100-150 million hooks per year through the 1990s. Beginning in 2000 effort increased substantially to ~350 million hooks per year in 2003, and has fluctuated around that level for the last 20 years.

A number of indices were investigated either as one-off sensitivities or included in the structural uncertainty grid (Figure 2). These included the composite distant water fishing nation (DWFN) longline standardized CPUE index (1988-2022) from the diagnostic case of the 2024 assessment (Castillo-Jordan et al., 2024), an Australian longline index, a New Zealand recreational sportfish index, and three different observer data based indices (Neubauer et al., 2025). The Pacific Islands observer program indices contained information from New Caledonia, Fiji, Tonga, and French Polynesia observer programs. Three alternative indices were constructed either by including data from all programs together, including data from just the western programs (New Caledonia, Fiji, and Tonga) or including data just from French Polynesia. The DWFN, Australia and observer without French Polynesia indices show recent increases; the New Zealand and observer with only French Polynesia indices show recent declines; and the all observer program index shows no trend.

3.1.2 Population dynamics

The population dynamics follow a hybrid Fletcher-Schaefer surplus production model (Edwards, 2024; ISC, 2024) with state-space formulation where population depletion x_t (relative to carrying capacity K) evolves

according to:

$$x_t = (x_{t-1} + \text{surplus production}_{t-1}) \times \epsilon_t \times \exp(-F_{t-1})$$

where surplus production is defined as:

$$\text{surplus production}_{t-1} = \begin{cases} R_{Max} \cdot x_{t-1} \left(1 - \frac{x_{t-1}}{h}\right), & x_{t-1} \leq D_{MSY} \\ \gamma \cdot m \cdot x_{t-1} \left(1 - x_{t-1}^{n-1}\right), & x_{t-1} > D_{MSY} \end{cases}$$

Here, R_{Max} is the maximum intrinsic rate of increase, n is the shape parameter, F_{t-1} is the fishing mortality in year $t-1$, and ϵ_t represents multiplicative process error with $\epsilon_t = \exp(\delta_t - \sigma_P^2/2)$ and $\delta_t \sim N(0, \sigma_P)$. The intermediate parameters are: $D_{MSY} = (1/n)^{1/(n-1)}$, $h = 2D_{MSY}$, $m = R_{Max} \cdot h/4$, and $\gamma = n^{n/(n-1)}/(n-1)$.

Fishing mortality F_t is in turn defined as a function of fishing effort and time-varying catchability:

$$F_t = q_{eff} \times \exp(\delta_{q,t} - \sigma_{qdev}^2/2) \times E_t$$

where q_{eff} is the base catchability coefficient, E_t is the fishing effort² in year t , and $\delta_{q,t}$ represents temporal variation in catchability effectiveness.

The catchability deviations $\delta_{q,t}$ follow a structured temporal process organized into discrete periods of length n_{step} years. In the diagnostic model $n_{step} = 1$ which allows for annual catchability deviates across 70³ periods p . Catchability remains constant within periods p , and evolves between periods according to an AR(1) process:

$$\delta_{q,p} = \rho \times \delta_{q,p-1} + \epsilon_{q,p} \times \sigma_{qdev} \times \sqrt{1 - \rho^2}$$

where ρ is the autocorrelation parameter, σ_{qdev} is the standard deviation of catchability innovations, and $\epsilon_{q,p} \sim N(0, 1)$. The initial period is constrained such that $\delta_{q,1} = 0$. The term $\sqrt{1 - \rho^2}$ normalizes the variance of $\delta_{q,p}$ to remain stationary at σ_{qdev}^2 regardless of the value of ρ .

For each year t , the corresponding period assignment is:

$$\text{period}(t) = \min \left(\left\lfloor \frac{t-1}{n_{step}} \right\rfloor + 1, n_{periods} \right)$$

and the catchability deviation for year t is assigned as $\delta_{q,t} = \delta_{q,\text{period}(t)}$. This fishing mortality parametrization is a simplification of the *catch-errors* formulation in MULTIFAN-CL (Fournier et al., 1998) and allows catchability to vary smoothly over time while maintaining temporal structure and avoiding overfitting to annual fluctuations in the effort-catch relationship.

²When input in the model the raw input effort, defined as *hundred hooks* in the WCPFC database is divided by 1e6 to re-scale catchability so that it is numerically on the same scale as other estimated parameters. This improves convergence of the HMC sampling of the posterior distribution.

³There are 71 total periods in the model period 1952-2022 and fishing mortality is not estimated in the terminal year.

Model initial conditions are defined for $t = 1$ as $x_t = x_0 \times \epsilon_t$ where x_0 is the estimated depletion relative to unfished at the start of the model period. The prior specified for x_0 in the diagnostic case assumes an unfished initial condition with some uncertainty (Table 2).

3.1.3 Observation model

The model fits to a standardized CPUE index using a lognormal likelihood:

$$I_t \sim \text{Lognormal}(\log(q \times x_t) - \frac{\sigma_{O,t}^2}{2}, \sigma_{O,t})$$

where catchability q is analytically derived assuming an uninformative uniform prior, and total observation error $\sigma_{O,t} = \sigma_{O,input} + \sigma_{O,add}$ combines fixed input uncertainty with an estimated additional error component.

3.1.3.1 Catch treatment

Population removals (catch) were calculated in each time step as:

$$\text{Removals}_{t-1} = (x_{t-1} + \text{surplus production}_{t-1}) \times \epsilon_t \times (1 - \exp(-F_{t-1})) \times \exp(\log K)$$

As described in Section 3.1.2, instantaneous fishing mortality F_t was transformed to a discrete proportion of the population that survived fishing, survivorship, through the exponential decay function $\exp(-F_{t-1})$. Removals are simply calculated as the complementary proportion $(1 - \exp(-F_{t-1}))$ of the total production (including surplus production and process error) that are removed due to fishing and converted to absolute units, numbers, through $\exp(\log K)$.

The model fits to the observed total catch in numbers using a Student's t-distribution likelihood:

$$\log(\text{Removals}_{obs,t}) \sim \text{Student-t} \left(\nu_c, \log(\text{Removals}_t) - \frac{\sigma_{C,t}^2}{2}, \sigma_{C,t} \right)$$

where the degrees of freedom parameter ν_c controls the tail behavior⁴, and the observation error $\sigma_{C,t}$ allows for time-varying uncertainty in catch observations. However, in the diagnostic model configuration $\sigma_{C,t} = 0.2$ for all t .

3.2 Prior development

The baseline prior distributions for leading estimated model parameters are specified in Table 2. A description of the derivation of these prior distributions follows.

Informed prior distributions for R_{Max} , and the Pella-Tomlinson shape parameter n were developed using numerical simulation based on biological knowledge. Informed prior distributions for $\log(K)$, and baseline catchability q_{eff} were developed using numerical simulation based on existing fishing data observations (e.g., catch and effort). Priors for other estimated leading model parameters (initial depletion x_0 , AR1 correlation

⁴As $\nu_c \rightarrow \infty$ the likelihood equation converges to the lognormal distribution $\text{Lognormal} \left(\log(\text{Removals}_t) - \frac{\sigma_{C,t}^2}{2}, \sigma_{C,t} \right)$

in catchability deviates ρ , variability in catchability deviates σ_{qdev} , process error σ_P , index observation error $\sigma_{O,add}$, and degrees of freedom of the catch likelihood ν_{catch}) were based on expert opinion and values observed in the literature. The naïve or input prior distributions were refined using a prior pushforward analysis.

3.2.1 Biological simulation framework

Informative priors for key population parameters were developed through biological simulation using Monte Carlo sampling from biologically plausible parameter distributions⁵. The simulation framework incorporated uncertainty and correlations in life history parameters:

- **Growth parameters:** Francis parameterization with independent sampling
- **Natural mortality:** Reference mortality M_{ref} negatively correlated with maximum age ($r = -0.3$)
- **Maturity:** Length-based logistic function parameters
- **Reproduction:** Steepness parameter for stock-recruit relationship and sex ratio
- **Weight-length relationship:** Correlated allometric parameters ($r = -0.5$) with biological constraints
- **Fishery selectivity:** Length at first capture for depletion prior

Parameter combinations (Table 1) were filtered to ensure biological realism and included correlations between life history parameters reflecting biological trade-offs (e.g., shorter lived individuals having higher natural mortality rates).

3.2.1.1 Maximum intrinsic rate of increase (R_{Max}) prior

R_{Max} was calculated by numerically solving the Euler-Lotka equation using bounded optimization (nlminb function in R) to find the value of R_{Max} that satisfies the equation within convergence criteria. Starting values and bounds were set to ensure biologically realistic solutions. Formulation of the Euler-Lotka equation followed the approach implemented in openMSE (Hordyk et al., 2024; Stanley et al., 2009):

$$\alpha \sum_{a=1}^{A_{Max}} l_a f_a \exp(-R_{Max} \times a) = 1$$

where α is the slope at the origin of the stock-recruitment curve, l_a is the probability of survival to age a , and f_a is the fecundity (reproductive output) at age a . Additional technical detail can be found in Section 11.1 of the Technical Annex. Repeating this calculation across the 500,000 parameter combinations resulted in a prior distribution of R_{Max} conditioned on plausible biology and life-history characteristics. The resulting distribution was first filtered to values of $R_{Max} < 1.5$ representative of what the literature (Gravel et al., 2024; Hutchings et al., 2012) indicates is most likely for teleosts. A lognormal distribution was fit to the filtered distribution to derive input prior parameters for the Stan model.

3.2.1.2 Pella-Tomlinson shape parameter (n) prior

Similarly, a prior distribution for the Pella-Tomlinson shape parameter n was derived from the filtered R_{Max} values and associated biological parameter combinations through a two-step process. First, the inflection point of the production function was estimated using the relationship established by Fowler (1988):

⁵The R code for the simulation `01-01-define-rmax-prior.r` is publicly available on [GitHub](#).

$$\phi = 0.633 - 0.187 \times \log(T_G \times R_{Max})$$

where ϕ is the inflection point (representing the population level at which maximum sustainable yield occurs as a fraction of carrying capacity), T_G is the generation time calculated following Grant & Grant (1992) as $T_G = \sum_{a=1}^{A_{Max}} a \times l_a \times f_a / \sum_{a=1}^{A_{Max}} l_a \times f_a$, and R_{Max} is the maximum intrinsic rate of increase.

The Pella-Tomlinson shape parameter n was then derived from the inflection point through the mathematical relationship $\phi = (1/n)^{1/(n-1)}$, solved numerically using bounded optimization in R as done for R_{Max} . This transformation links the biologically-derived inflection point to the production function curvature, where $n = 2$ corresponds to the symmetric Schaefer model (e.g., MSY achieved at depletion $D = 0.5$). Lower values of n ($n < 2$) indicate more productive stocks that achieves MSY at $D < 0.5$ and higher values of n indicate less productive stocks with MSY occurring at $D > 0.5$. A lognormal distribution was fit to the distribution of n to derive input prior parameters for the Stan model. Additionally, linking the shape parameter n to R_{Max} necessitated a multivariate prior structure so the correlation between the two parameters was estimated as well.

3.2.2 Fishing data simulation framework

Prior distributions for carrying capacity $\log(K)$ and initial catchability coefficient q_{eff} were derived using catch and effort data combined with the biologically-informed parameter distributions⁶.

3.2.2.1 Carrying capacity prior ($\log(K)$)

The carrying capacity prior was constructed by assuming the maximum observed catch approximates maximum sustainable yield (MSY). This provides a conservative assumption for the upper limit of population scale⁷ as larger MSY correlates positively with larger population scale. To account for uncertainty in the maximum catch $\sim$ MSY relationship (e.g., true MSY was above or below the maximum observed catch), candidate MSY values were sampled from a lognormal distribution centered on the maximum observed catch with moderate levels of uncertainty ($MSY \sim \text{Lognormal}(\log(\text{max catch}), 0.5)$).

For each combination of MSY, biologically-derived R_{Max} and shape parameter n , carrying capacity was calculated as:

$$K = \frac{MSY}{r \times \phi \times (1 - \phi)^{n-1}}$$

where $\phi = (1/n)^{1/(n-1)}$ represents the inflection point of the production function. For the Fox model special case where $n = 1$, the relationship simplifies to $K = e \times MSY/r$. Non-finite values and extreme outliers ($\log(K) > 20$) were replaced through resampling from the valid parameter space to ensure numerical stability. A lognormal distribution was fit to the filtered distribution to derive input prior parameters for the Stan model.

⁶This is applied in the R script [03-06-new-logK-prior-pushforward-bspm_estqsimple_softdep_mvprior_x0.r](#) lines 64-92 which is publicly available on [GitHub](#).

⁷The upper limit of population scale is typically more challenging to estimate than the lower population limit as population scales of infinite sizes can support any observed catches. However, there typically is a hard floor to population scale in that it must be sufficiently large to maintain a population that produces the time series of observed catches.

3.2.2.2 Initial effort catchability prior (q_{eff})

Next, the initial catchability coefficient prior was developed using the relationship between CPUE and stock abundance:

$$q_{eff} = \frac{CPUE}{K}$$

where CPUE was calculated from early fishery data (first 10 years of catch and effort records) when the stock was assumed to be closer to carrying capacity. The prior distribution was constructed by randomly pairing CPUE values sampled from the early period with carrying capacity values from the $\log(K)$ prior distribution to generate a distribution of q_{eff} values incorporating both the uncertainty in population scale and the variability observed in early fishery CPUE. Again, a lognormal distribution was fit to the filtered distribution to derive input prior parameters for the Stan model.

3.2.3 Other priors

As mentioned previously, initial prior distributions for the remaining leading model parameters (initial depletion x_0 , AR1 correlation in catchability deviates ρ , variability in catchability deviates σ_{qdev} , process error σ_P , index observation error $\sigma_{O,add}$, and degrees of freedom of the catch likelihood ν_{catch}) were specified either using expert opinion or literature.

Given that the limited large-scale industrial fishing existed in the SWPO prior to the expansion of the Japanese distant-water longline fleet in 1952, the population was assumed to start at an unfished condition with some uncertainty. Initial depletion x_0 was modeled with a lognormal distribution centered on 1. A lognormal prior was used for the standard deviation of the process error where the parameters were converted from the JABBA default prior for process error (Winker et al., 2018). The degrees of freedom ν_{catch} parameter assumed a gamma prior based on suggestions in the literature that it serves as a sensible default choice (Juárez & Steel, 2010). The prior for index observation error $\sigma_{O,add}$ assumed a half-normal distribution following ISC (2024) which provides sufficient support for a broad range of extra estimated observation error. For the autocorrelation coefficient in catchability deviations (ρ) and the standard deviation of catchability deviations (σ_{qdev}), initial priors were derived from data-filtering simulation results and specified as a bivariate distribution on the transformed scale: $\tanh^{-1}(\rho)$ and $\log(\sigma_{qdev})$.

3.2.4 Prior pushforward

Following Kim & Neubauer (2025), a prior pushforward analysis was conducted to refine the naïve prior distributions derived from the biological and fishing data simulation frameworks. Random parameter combinations drawn from the input prior distributions were passed through the diagnostic Stan model used for inference but with the likelihoods turned off (`fit_to_data = 0L`), generating simulated population trajectories and fishing dynamics under the model structure. This approach ensures that simulated trajectories are consistent with the actual model structure used for inference.

Parameter combinations were filtered through sequential criteria based on population viability and fishery realism:

- **Initial survival filter** (`iter_surve`): Terminal year depletion $> 1\%$ (`mindep > 0.01`), minimum

population size $> 15,000$ individuals which matches recent observed catch levels (`minN > 15000`), and biologically reasonable shape parameter (`shape < 20`)

- **Fishing mortality filter (`iter_f`)**: Maximum fishing mortality < 2.5 (`maxF < 2.5`) and average fishing mortality < 0.8 (`avgF < 0.8`) to ensure realistic exploitation levels
- **Catch consistency filter (`iter_catch`)**: Simulated catches within realistic bounds relative to observed data: maximum catch between $0.5\text{-}5.0\times$ observed maximum, minimum catch $> 0.5\times$ observed minimum, and average catch between $0.5\text{-}5.0\times$ observed average
- **CPUE consistency filter (`iter_cpue`)**: Early period CPUE (first 10 years) between $0.5\text{-}5.0\times$ observed early CPUE average to ensure realistic abundance levels during the initial fishery development

Parameter combinations passing all filtering stages (`iter_cpue`) were used to fit updated multivariate prior distributions using maximum likelihood estimation. Given that a number of parameters showed correlations following the prior pushforward, a 7-dimensional multivariate prior was developed for core parameters: $\log(K)$, $\log(R_{Max})$, $\log(n)$, $\log(x_0)$, $\log(q_{eff})$, $\tanh^{-1}(\rho)$ and $\log(\sigma_{qdev})$. Correlations with absolute values < 0.1 were reset to zero to avoid weak spurious correlations, retaining only meaningful parameter relationships. Though $\log(x_0)$ was included in the multivariate prior, for convenience, it was treated as an unchanged univariate prior from the initial naïve prior distribution since the pushforward did not provide an update to this parameter.

Prior distributions for process error σ_P , index observation error $\sigma_{O,add}$, and degrees of freedom of the catch likelihood ν_{catch} were unchanged following the prior pushforward analysis and remained as univariate priors. Final prior distributions are listed in Table 2.

3.3 Model implementation

The models were implemented using the No-U-Turn Sampler (NUTS) in Stan. Each model was run using 5 independent Markov chains with randomly generated starting values to ensure robust exploration of the posterior parameter space. Each chain was sampled for 3,000 iterations, with the first 1,000 iterations discarded as warmup to allow for algorithm adaptation. To reduce posterior sample autocorrelation and storage requirements, every 10^{th} samples was retained from the post-warmup iterations, yielding a final sample size of 1,000 draws from the joint posterior distribution across all chains. The NUTS algorithm was configured with strict adaptation parameters to ensure reliable convergence. The adaptation delta was set to 0.99 to reduce the step size and minimize divergent transitions, while the maximum tree depth was increased to 12 to allow for longer trajectories during the dynamic sampling process.

Model convergence and performance were assessed using multiple diagnostic criteria recommended for Bayesian stock assessment workflows (Monnahan, 2024). The potential scale reduction factor ($\hat{R}$) was required to be less than 1.01 for all parameters, indicating convergence across chains. Effective sample sizes were required to exceed 500 to ensure adequate precision in posterior estimates. Additionally, posterior predictive checks were conducted to evaluate model adequacy through comparison of observed versus predicted index and catch values, while parameter identifiability was assessed through visual examination of posterior distributions and correlation structures among estimated parameters (Gabry et al., 2019).

3.4 Model diagnostics

Model performance was comprehensively evaluated using multiple diagnostic criteria recommended for Bayesian stock assessment workflows (Monnahan, 2024). The diagnostic framework included Stan model

convergence criteria, data fits, posterior predictive checks, retrospective analysis, and hindcast cross-validation.

3.4.1 Convergence diagnostics

Conventional Stan model convergence diagnostics and thresholds were employed to identify whether posterior distributions were representative and unbiased (Monnahan, 2024). Models were considered to have ‘converged’ to a stable, unbiased posterior distribution if they satisfied the following criteria:

- **Potential scale reduction factor** ($\hat{R}$) less than 1.01 for all leading model parameters, indicating convergence across chains
- **Bulk effective sample size** (N_{eff}) greater than 500 for all leading model parameters, ensuring adequate precision in posterior estimates
- **No divergent transitions** during the NUTS sampling process, which can indicate problematic posterior geometry
- **Maximum tree depth** not exceeded during sampling, ensuring adequate exploration of parameter space

Additionally, trace plots and rank plots were examined to visually assess mixing and convergence behavior across chains (Gabry et al., 2019).

3.4.2 Leave-one-out cross-validation diagnostics

Model predictive performance and influential observations were assessed using leave-one-out cross-validation (LOO-CV) implemented through the *loo* package (Vehtari et al., 2017; Vehtari et al., 2024). LOO-CV provides an estimate of the expected log pointwise predictive density (ELPD) while identifying problematic observations that may unduly influence model predictions.

LOO-CV was also used as a factor to compare performance between models. Following the recommendations of Sivula et al. (2020), models were considered to have meaningfully different performance when the ELPD difference exceeded 4 units. For differences larger than 4 units, statistical significance was assessed by comparing the ELPD difference to twice its standard error, with differences greater than $2 \times$ the standard error considered statistically significant.

The effective number of parameters (p_{loo}) was examined as an additional diagnostic measure, with values substantially larger than the nominal number of model parameters indicating potential model misspecification. Catch and index observations were categorized as having low influence ($\text{Pareto-}k \leq 0.7$), high influence ($0.7 < \text{Pareto-}k \leq 1.0$), or very high influence ($\text{Pareto-}k > 1.0$). Observations with high or very-high influence were investigated to identify potential outliers, data entry errors, or systematic model misspecification that could compromise predictive performance.

3.4.3 Data fits

Model fits to the abundance index and catch time series were each quantified using normalized root-mean-squared error (NRMSE):

$$RMSE = \sqrt{\frac{\sum_{i=1}^n (y_i - \hat{y}_i)^2}{n}}$$

$$NRMSE = \frac{RMSE}{\sum_{i=1}^n y_i / n}$$

where y_i are the observed values and $\hat{y}_i$ are the model predictions. The average NRMSE across all posterior samples was calculated for each data component. Additionally, Probability Integral Transform (PIT) style residuals were calculated for the fit to the index.

3.4.4 Posterior predictive checks

Posterior predictive checks were conducted to evaluate model adequacy by comparing observed data with data simulated from the fitted model. For each posterior sample, synthetic datasets were generated using the estimated parameters, and agreement between observed and predicted datasets was assessed visually.

3.4.5 Retrospective analysis

Retrospective analysis was conducted for each model by sequentially peeling off a year from the terminal end of the fitted index and re-running the model. Data were removed for up to five years, from 2022 back to 2018. Estimates of x in the terminal year of each retrospective peel were compared to the corresponding estimate of x from the full model run to better understand any potential biases or uncertainty in terminal year estimates. The Mohn's ρ statistic (Mohn, 1999) was calculated and presented. This statistic measures the average relative difference between an estimated quantity from an assessment (e.g., depletion in final year) with a reduced time-series of information and the same quantity estimated from an assessment using the full time-series.

Additionally, following Kokkalis et al. (2024) the proportion of retrospective peels (or coverage) where the relative exploitation rate (U/U_{MSY}) and relative depletion (D/D_{MSY}) were inside the credible intervals of the full model run was calculated. This metric provides an additional perspective on retrospective consistency by evaluating whether the uncertainty estimates from the full model appropriately capture the range of estimates obtained when using reduced datasets.

3.4.6 Hindcast cross-validation

Hindcast cross-validation (Kell et al., 2021) was conducted for each index to determine model performance in predicting the observed CPUE I one-step-ahead into the future relative to a naïve predictor. Briefly, the ‘model-free’ approach to hindcast cross-validation was used, and used the same set of five retrospective peels described in Section 3.4.5.

The ‘model-free’ hindcast calculation is described using the model from the last peel $BSPM_{2017}$ as an example. This model fit to index data through 2017 but included catch through 2022. The model estimates of predicted CPUE in 2018 based on $BSPM_{2017}$ is the ‘model-free’ hindcast for 2018, $\hat{I}_{2018}$. The naïve prediction of CPUE in 2018 is simply the observed CPUE from 2017, $I_{2017} = \check{I}_{2018}$. The absolute scaled error (ASE) of the prediction is:

$$ASE_{2018} = \frac{|I_{2018} - \hat{I}_{2018}|}{|I_{2018} - \hat{I}_{2018}|}$$

Repeating this calculation across all retrospective peels for years 2019-2022 and taking the average across ASE values gives the mean ASE or MASE for the model. An MASE value less than one indicates that the model has greater predictive skill than the naïve predictor.

3.5 Forecasts

Simple, status quo catch-based stochastic projections over a 10-year window are provided for each model. For each model, projected catch was taken as the average catch over the terminal 5 years (2018-2022). Process error in the projection period was randomly resampled from the estimated process error in the main model period.

4 Results

4.1 Diagnostic case

4.1.1 Model convergence and diagnostics

All three BSPM model variants achieved satisfactory convergence based on standard Hamiltonian Monte Carlo diagnostics (Table 3). All models met convergence criteria with $\hat{R}$ values well below the 1.01 threshold and effective sample sizes exceeding the recommended minimum of 500 (100 samples per chain). No divergent transitions were observed across chains, and maximum tree depth was not exceeded during sampling, indicating successful exploration of the posterior space.

4.1.1.1 Model fit to data

4.1.1.1.1 Catch data fit

All models fitted the observed catch time series (1952-2022) with a coefficient of variation of 0.2 to account for uncertainty in catch estimates. Since fishing mortality required to generate the catch was directly estimated, model fit to the catch was good (Figure 10). The models consistently predicted lower than observed catch for the large catch event in 1954.

4.1.1.1.2 CPUE index fit

Model fit to the standardized CPUE index (1988-2022) was evaluated using normalized root-mean-squared error (NRMSE). All three models provided reasonable fits to the observed index given the high variability and observation error (Table 3). The `0003-2024cpueFPrior` model showed marginally improved fit with lower RMSE values. All models successfully captured the general declining trend in standardized CPUE from 1988-2022. However, the estimated combination of observation and process error meant models were not tightly constrained to individual observations, resulting in persistent negative residuals in the last decade (Figure 11). Posterior predictive checks confirmed that models replicated key features of the observed CPUE data (Figure 12).

4.1.1.2 Model validation

Overall retrospective bias was very low and close to zero for all models (Table 3; Figure 13), indicating lack of persistent bias in terminal estimates as successive years of index data were removed. Estimates of relative exploitation rate (U/U_{MSY}) and relative depletion (D/D_{MSY}) fell within the credible intervals of the full model run 100% of the time (Table 3). For hindcast cross-validation, all models exceeded the performance of the naïve predictor, with the 0003-2024cpueFPrior model showing the best performance (Table 3; Figure 14). This provides support that models successfully identified and estimated a production function for the stock.

4.1.2 Population dynamics and stock status

In all models, most leading parameters ($\log(K)$, R_{Max} , σ_P , n , σ_F , and $\sigma_{O,add}$) showed posterior updates from their respective prior distributions (Figure 15), indicating sufficient information in the data to inform parameter estimates. The shape parameter n showed little deviation from the Schaefer-model prior. All models estimated similar levels of σ_P and additional observation error $\sigma_{O,add}$ (Table 4), which is unsurprising given they fitted the same catch and standardized index. Although estimates of σ_P were elevated from the prior, models primarily reconciled index variability by estimating substantial additional observation error $\sigma_{O,add}$, resulting in approximately a 3:1 ratio of total observation error σ_O to process error σ_P . Similar to findings in ISC (2024), there was a trade-off between $\log(K)$ and σ_F where higher population scale was associated with lower fishing mortality and vice versa.

All models showed substantial posterior updates for the R_{Max} parameter, estimating productivity on the lower end of the biological prior. These median estimates of R_{Max} , between 0.2-0.3 (Table 4), are consistent with estimates for an analogous species in the Atlantic Ocean, white marlin (*Kajikia albida*). The most recent ICCAT assessment (ICCAT, 2020), implemented using JABBA (Winker et al., 2018), estimated R_{Max} at 0.163 (95% credible interval: 0.122-0.215), which falls within the posterior estimates of all three models (Table 4). Notably, the Atlantic white marlin JABBA model showed remarkable consistency with stock status and trend estimates from a fully integrated, length-structured model implemented in Stock Synthesis (ICCAT, 2020). In the current analysis, the 0003-2024cpueFPrior model estimated lower productivity R_{Max} with reduced uncertainty (Table 4), consistent with assuming a more depleted stock. Inverse transform sampling to match the R_{Max} posterior distribution from this model reveals that lower productivity is characterized by pronounced shifts toward lower steepness values h and, to a lesser extent, smaller von Bertalanffy k values (Figure 16).

The estimated population depletion time series shows a decline from the assumed unfished state in 1952 to minimum levels around the early 1990s (Figure 17). All three models estimated similar depletion trajectories, with differences primarily related to estimated population scale. The 0003-2024cpueFPrior model estimated slightly lower depletion levels throughout the time series compared to the other models. From the minimum depletion period, all models estimated gradual recovery through the 2000s and 2010s, with populations stabilizing at moderate depletion levels in recent years. Credible intervals around depletion estimates are relatively wide, particularly in the early period when no abundance index data were available to inform the model, reflecting inherent uncertainty in reconstructing historical population dynamics. The estimated population time series mirrors these depletion patterns. The 0001-2024cpueExPrior model estimated the highest population scale throughout the time series, while the 0002-2024cpueDepPrior and

0003-2024cpueFPrior models estimated lower population levels. While uncertainty in absolute population size is substantial, as reflected in wide credible intervals, relative trends are more precisely estimated.

Process error estimates (Figure 17) showed consistent temporal variability across models, with the largest deviations occurring during periods of rapid population change in the 1970s and 1980s. This indicates that removals alone could not explain the observed changes in the standardized index. The estimated process error variability σ_P (Table 4), while similar across all models, was lowest for the 0003-2024cpueFPrior model, indicating greater internal consistency in the model dynamics.

4.2 One-off sensitivities

4.3 Model ensemble

4.3.1 Population dynamics and stock status

4.3.2 Biological reference points

MSY-based reference points were derived from the Fletcher-Schaefer production function for all three BSPM variants. The relative stock status time series (Figure 17) shows consistent trends across all models, albeit at different relative levels. Median estimates of the D/D_{MSY} ratio indicate the stock was above the MSY reference level for the entire model period. Only the 0003-2024cpueFPrior model indicates risk of falling below the MSY-based reference point in recent years. Similarly, median estimates of the F/F_{MSY} ratio were consistently below the overfishing reference point, with episodic periods of overfishing risk indicated by the 0003-2024cpueFPrior model. However, recent trends indicate declining overfishing risk, consistent with estimated stock recovery.

4.3.3 Future projections

Ten-year stochastic projections (2023-2032) were conducted for all three BSPM variants under a status quo catch scenario, with projected removals set to the average catch over 2018-2022 (Figure 18). Under status quo catch levels, all models projected continued population recovery through 2032. Depletion trajectories show gradual increases from current levels, with the 0001-2024cpueExpPrior model projecting the most optimistic recovery due to higher estimated population scale. The 0002-2024cpueDepPrior and 0003-2024cpueFPrior models showed more modest but consistent increases in depletion over the projection period.

Relative to MSY-based reference points, projections indicate the stock will likely remain above D_{MSY} throughout the projection period for the baseline and depletion-constrained models. The 0003-2024cpueFPrior model showed greater variability and some risk of declining below D_{MSY} in the near term, though the median trajectory suggests stabilization above the reference point by the end of the projection period. Projected fishing mortality rates remained well below F_{MSY} for all models under the status quo catch scenario, indicating that current harvest levels are likely sustainable and consistent with continued stock recovery. Projection uncertainty increased over time, reflecting both parameter uncertainty and stochastic variability in future population dynamics.

5 Discussion

5.1 General remarks

The three BSPM variants provide a consistent picture of stock trajectory and current status. All models estimated similar depletion patterns: decline from unfished conditions in 1952 to minimum levels around the early 1990s, followed by gradual recovery through recent decades. This trajectory suggests the stock is currently recovering. While median estimates indicate the stock was not overfished or undergoing overfishing during the assessment period, the `0003-2024cpueFPrior` model suggests some recent risk of falling below these thresholds.

The BSPM approach addresses key limitations encountered in previous integrated assessments while providing robust stock status estimates. The primary advantage lies in condensing complex biological and fishery uncertainties into a manageable parameter set. Integrated assessments can be influenced by size composition data, which are sensitive to assumptions about growth, mortality, selectivity, and reproduction parameters. When these assumptions are uncertain or mis-specified, population scale estimates informed by size composition data can be biased. Production models circumvent this issue by aggregating these uncertainties into productivity parameters, allowing more efficient exploration of plausible population dynamics while maintaining biological realism through informed priors.

5.2 Challenges, limitations & key uncertainties

Several important limitations affect this analysis. The assumption of no age-specific population dynamics may be problematic when mortality or reproductive potential varies significantly by age. This simplification prevents incorporation of biological information beyond production function parameters and excludes explicit use of size composition data.

The aggregated nature of surplus production models can potentially bias MSY estimates when age-specific processes differ from production function assumptions. For species with complex life histories like striped marlin, managers should consider these limitations when applying MSY-based reference points derived from surplus production models.

Data limitations also constrain the analysis. Fitting to a single standardized abundance index limits evaluation of uncertainty in abundance trends. Multiple indices would enable model ensemble approaches to better characterize this uncertainty.

The incorporation of depletion information from New Zealand recreational size data through a 1988 depletion prior, while novel, represents a simplified approach. More sophisticated modeling frameworks like delay-difference or age-structured models could explicitly fit size composition time series in a likelihood-based framework, providing more elegant integration of this information.

Modeling assumptions introduce additional uncertainties. Current modeling assumes the same level of catch uncertainty across all time periods. Future versions could allow time-varying catch uncertainty (e.g., greater uncertainty in historical versus recent observations) or accommodate effort-driven removals similar to catch-errors with effort deviation approaches available in MULTIFAN-CL (Fournier et al., 1998).

Population scale estimates were highly uncertain and sensitive to model configuration choices, while depletion

estimates were more precisely estimated. The choice of prior on σ_F was particularly influential. Although this prior was developed using prior pushforward analysis (Kim & Neubauer, 2025), it represents a key uncertainty. Similarly, uncertainty in biological parameters (growth, maturity, natural mortality) due to limited sample sizes contributed to broad initial priors for R_{Max} . While the posterior distribution for R_{Max} was substantially updated from the prior, reducing uncertainty in key biological productivity parameters could help reduce model uncertainty estimates and facilitate development of age-structured models.

5.3 Future stock assessment modeling considerations

Future assessment efforts should progressively build toward full age-structure through a systematic approach: delay-difference models, age-structured production models, and then fully integrated age-structured models. This development should occur within a Bayesian framework to inform parameter estimates where prior information exists or can be derived using prior pushforward approaches (Kim & Neubauer, 2025).

Developing Bayesian age-structured modeling approaches can provide more stable estimation of increased parameters due to the use of priors, explicitly account for parameter uncertainty rather than making fixed assumptions, potentially enable informed estimation of difficult-to-observe biological relationships (e.g., stock-recruit relationships), address BSPM limitations by explicitly considering age-structured processes and fishery selectivity, and estimate leading parameters with likelihood components for indices and size composition.

While simplified models offer advantages when data conflicts exist, over-simplification of age-structured processes could result in biases. Future modeling must balance complexity with data quality and conflicts. The transition should be guided by careful evaluation of whether increased model complexity is supported by available data and whether it improves the reliability of management advice.

6 Acknowledgements

The authors appreciate feedback received on earlier versions of this analysis, and are thankful to the following for their insights and thoughtful discussion which greatly improved the assessment: K. Kim (SPC), N. Davies (Te Takina Ltd.), S. Hoyle (Hoyle Consulting), R. Ahrens (PIFSC), M. Nadon (PIFSC), and F. Carvalho (PIFSC).

7 Declaration of generative AI use

A generative artificial intelligence (AI) assistant, Anthropic Claude Sonnet 4.0, was used to parse model code in the preparation of an initial draft of this report.

8 References

- Betancourt, M. J., & Girolami, M. (2013). *Hamiltonian Monte Carlo for Hierarchical Models*. <https://doi.org/10.48550/ARXIV.1312.0906>
- Castillo-Jordan, C., Day, J., Tears, T., Davies, N., Hampton, J., McKechnie, S., Magnusson, A., Peatman, T., Vidal, T., Williams, P., & Hamer, P. (2024). *Stock Assessment of Striped Marlin in the Southwest*

- Pacific Ocean: 2024* (WCPFC-SC20-2024/SA-WP-03 (Ver. 2.01)). <https://meetings.wcpfc.int/node/23120>
- Castillo-Jordan, C., Day, J., Teears, T., Davies, N., Hampton, J., McKechnie, S., Magnusson, A., Peatman, T., Vidal, T., Williams, P., & Hamer, P. (2025). *Revised Stock Assessment of Striped Marlin in the Southwest Pacific Ocean: 2025* (WCPFC-SC21-2025/SA-WP-XX).
- Ducharme-Barth, N., Castillo-Jordan, C., Sculley, M., Ahrens, R., & Carvalho, J. (2025). *Summary report from the NOAA-SPC assessment model meeting on SWPO striped marlin* (WCPFC-SC21-2025/SA-IP-XX).
- Edwards, C. (2024). *Bdm: Bayesian biomass dynamics model* (R package version 0.0.0.9036).
- Farley, J., Eveson, P., Krusic-Golub, K., & Kopf, K. (2021). *Southwest Pacific Striped Marlin Population Biology (Project 99)* (WCPFC-SC-17/SA-IP-11). <https://meetings.wcpfc.int/node/12569>
- Fournier, D. A., Hampton, J., & Sibert, J. R. (1998). MULTIFAN-CL: A length-based, age-structured model for fisheries stock assessment, with application to South Pacific albacore, *Thunnus alalunga*. *Canadian Journal of Fisheries and Aquatic Sciences*, 55(9), 2105–2116. <https://doi.org/10.1139/f98-100>
- Fowler, C. W. (1988). Population dynamics as related to rate of increase per generation. *Evolutionary Ecology*, 2(3), 197–204. <https://doi.org/10.1007/BF02214283>
- Gabry, J., & Mahr, T. (2024). *Bayesplot: Plotting for Bayesian Models* (R package version 1.11.1). <https://mc-stan.org/bayesplot/>
- Gabry, J., Simpson, D., Vehtari, A., Betancourt, M., & Gelman, A. (2019). Visualization in Bayesian Workflow. *Journal of the Royal Statistical Society Series A: Statistics in Society*, 182(2), 389–402. <https://doi.org/10.1111/rssa.12378>
- Grant, P. R., & Grant, B. R. (1992). Demography and the Genetically Effective sizes of Two Populations of Darwin’s Finches. *Ecology*, 73(3), 766–784. <https://doi.org/10.2307/1940156>
- Gravel, S., Bigman, J. S., Pardo, S. A., Wong, S., & Dulvy, N. K. (2024). Metabolism, population growth, and the fast-slow life history continuum of marine fishes. *Fish and Fisheries*, 25(2), 349–361. <https://doi.org/10.1111/faf.12811>
- Hamel, O. S., & Cope, J. M. (2022). Development and considerations for application of a longevity-based prior for the natural mortality rate. *Fisheries Research*, 256, 106477. <https://doi.org/10.1016/j.fishres.2022.106477>
- Hordyk, A., Huynh, Q., & Carruthers, T. (2024). *openMSE: Easily Install and Load the 'openMSE' Packages* (R package version 1.0.1). <https://openmse.com/>
- Hutchings, J. A., Myers, R. A., García, V. B., Lucifora, L. O., & Kuparinen, A. (2012). Life-history correlates of extinction risk and recovery potential. *Ecological Applications*, 22(4), 1061–1067. <https://doi.org/10.1890/11-1313.1>
- ICCAT. (2020). Report of the 2019 ICCAT white marlin stock assessment meeting. *Collect. Vol. Sci. Pap. ICCAT*, 78(4), 97–181. https://www.iccat.int/Documents/CVSP/CV076_CV076_2019/colvol76.html#
- ISC. (2024). *Stock Assessment of Shortfin Mako Shark in the North Pacific Ocean through 2022* (WCPFC-SC20-2024/SA-WP-14). <https://meetings.wcpfc.int/node/22828>
- Juárez, M. A., & Steel, M. F. J. (2010). Model-Based Clustering of Non-Gaussian Panel Data Based on Skew-*t*Distributions. *Journal of Business & Economic Statistics*, 28(1), 52–66. <https://doi.org/10.1198/jbes.2009.07145>
- Kell, L. T., Sharma, R., Kitakado, T., Winker, H., Mosqueira, I., Cardinale, M., & Fu, D. (2021). Validation of stock assessment methods: Is it me or my model talking? *ICES Journal of Marine Science*, 78(6), 2244–2255. <https://doi.org/10.1093/icesjms/fsab104>
- Kim, K., & Neubauer, P. (2025). Examining potential biases through prior predictive checks: Prior mis-

- specifications and their impact on Bayesian stock assessments. *Fisheries Research*, 288, 107405. <https://doi.org/10.1016/j.fishres.2025.107405>
- Kokkalis, A., Berg, C. W., Kapur, M. S., Winker, H., Jacobsen, N. S., Taylor, M. H., Ichinokawa, M., Miyagawa, M., Medeiros-Leal, W., Nielsen, J. R., & Mildenberger, T. K. (2024). Good practices for surplus production models. *Fisheries Research*, 275, 107010. <https://doi.org/10.1016/j.fishres.2024.107010>
- Mohn, R. (1999). The retrospective problem in sequential population analysis: An investigation using cod fishery and simulated data. *ICES Journal of Marine Science*, 56(4), 473–488. <https://doi.org/10.1006/jmsc.1999.0481>
- Monnahan, C. C. (2024). Toward good practices for Bayesian data-rich fisheries stock assessments using a modern statistical workflow. *Fisheries Research*, 275, 107024. <https://doi.org/10.1016/j.fishres.2024.107024>
- Neubauer, P., Castillo-Jordan, C., Day, J., & Hamer, P. (2025). *Exploring the potential for observer CPUE for southwest Pacific swordfish (Xiphias gladius) and striped marlin (Kajikia audax)* (WCPFC-SC21-2025/SA-IP-13). <https://meetings.wcpfc.int/node/26539>
- Neubauer, P., Kim, K., Large, K., & Brouwer, S. (2024). *Stock Assessment of Silky Shark in the Western and Central Pacific Ocean 2024* (WCPFC-SC20-2024/SA-WP-04-Rev2). <https://meetings.wcpfc.int/node/23088>
- Neubauer, P., Richard, Y., & Tremblay-Boyer, L. (2019). *Alternative assessment methods for oceanic whitetip shark* (WCPFC-SC15-2019/SA-WP-13).
- Pardo, S. A., Kindsvater, H. K., Reynolds, J. D., & Dulvy, N. K. (2016). Maximum intrinsic rate of population increase in sharks, rays, and chimaeras: The importance of survival to maturity. *Canadian Journal of Fisheries and Aquatic Sciences*, 73(8), 1159–1163. <https://doi.org/10.1139/cjfas-2016-0069>
- Sivula, T., Magnusson, M., Matamoros, A. A., & Vehtari, A. (2020). *Uncertainty in Bayesian Leave-One-Out Cross-Validation Based Model Comparison*. arXiv. <https://doi.org/10.48550/ARXIV.2008.10296>
- Stanley, R. D., McAllister, M., Starr, P., & Olsen, N. (2009). *Stock assessment for bocaccio (Sebastes paucispinis) in British Columbia waters* (Canadian {Science} {Advisory} {Secretariat} Research Document 2009/055). Fisheries; Oceans Canada. https://www.dfo-mpo.gc.ca/csas-sccs/publications/resdocs-docrech/2009/2009_055-eng.htm
- Team, S. D. (2024a). *RStan: The R interface to Stan* (R package version 2.32.6). <https://mc-stan.org/>
- Team, S. D. (2024b). *Stan Modeling Language Users Guide and Reference Manual, v 2.32.6*.
- Vehtari, A., Gabry, J., Magnusson, M., Yao, Y., Burkner, P., Paananen, T., & Gelman, A. (2024). *Loo: Efficient leave-one-out cross-validation and WAIC for Bayesian models* (R package version 2.7.0). <https://mc-stan.org/loo/>
- Vehtari, A., Gelman, A., & Gabry, J. (2017). Practical Bayesian model evaluation using leave-one-out cross-validation and WAIC. *Statistics and Computing*, 27(5), 1413–1432. <https://doi.org/10.1007/s11222-016-9696-4>
- Winker, H., Carvalho, F., & Kapur, M. (2018). JABBA: Just Another Bayesian Biomass Assessment. *Fisheries Research*, 204, 275–288. <https://doi.org/10.1016/j.fishres.2018.03.010>
- Winker, H., Mourato, B., & Chang, Y. (2020). Unifying parameterizations between age-structured and surplus production models: An application to atlantic white marlin (*Kajiki albidia*) with simulation testing. *Collect. Vol. Sci. Pap. ICCAT*, 76(4), 219–234.

9 Tables

Table 1: Biological parameter distributions and sources used in simulation framework

Parameter	Symbol	Distribution	Mean/Mode	CV/Range	Correlation	Source/Notes
Length at age 1	L_1	Lognormal	60 cm	0.2	Independent	(Ducharme-Barth et al., 2025)
Length at age 10	L_2	Lognormal	210 cm	0.2	Independent	(Ducharme-Barth et al., 2025)
Growth coefficient	k	Beta(6.5, 3.5)	~0.65	-	Independent	(Ducharme-Barth et al., 2025)
Length CV	cv_{len}	Uniform	-	[0.05, 0.25]	-	Growth variability
Maximum age	A_{Max}	Lognormal	15 years	0.2	$\rho = -0.3$ with M_{ref}	(Farley et al., 2021)
Reference mortality	M_{ref}	Lognormal	0.36 yr^{-1}	0.44	$\rho = -0.3$ with A_{Max}	(Hamel & Cope, 2022)
Maturity slope	a_{mat}	Normal	-20	0.2	-	(5.4/ A_{Max}) Logistic steepness
Length at 50% maturity	L_{50}	Lognormal	184 cm	0.2	-	(Farley et al., 2021)
Weight coefficient	a_w	Lognormal	5.4×10^{-7}	0.05	$\rho = -0.5$ with b_w	(Castillo-Jordan et al., 2024)
Weight exponent	b_w	Normal	3.58	0.05	$\rho = -0.5$ with a_w	(Castillo-Jordan et al., 2024) Constrained: 150-350 kg at 300 cm
Steepness	h	Beta(3, 1.5)	~0.73	-	-	Censored to [0.2, 1.0]
Sex ratio (prop. female)	s	Normal	0.5	0.05	-	Constrained [0.01, 0.99]
Reproductive cycle	-	Fixed	1 year	-	-	Annual spawning
Reference ages	age_1, age_2	Fixed	0, 10 years	-	-	Francis parameterization

Table 2: Prior distributions for effort-based Bayesian surplus production model leading parameters.

Parameter	Distribution	Mean	SD	Description	Key Correlations
Core Population Parameters					
K	Lognormal	$\log(1,110,533)$	0.59	Carrying capacity (N)	$\rho(K, R_{Max}) = -0.22$; $\rho(K, q_{eff}) = -0.241$
R_{Max}	Lognormal	$\log(0.6206)$	0.58	Maximum intrinsic rate of increase	$\rho(R_{Max}, n) = -0.881$
n	Lognormal	$\log(1.16)$	0.54	Pella-Tomlinson shape parameter	
x_0	Lognormal	$\log(1)$	0.025	Initial depletion	(Independent)
q_{eff}	Lognormal	$\log(0.5019)$	0.95	Effort catchability coefficient	$\rho(q_{eff}, \sigma_{qdev}) = 0.559$; $\rho(q_{eff}, \rho) = 0.243$
ρ	Transformed Normal	$\tanh^{-1}(0.981)$	0.62	Autocorrelation in catchability deviations	$\rho(\rho, \sigma_{qdev}) = 0.148$
σ_{qdev}	Lognormal	$\log(1.619)$	0.3	Catchability variability standard deviation	
Process and Observation Error					
σ_P	Lognormal	$\log(0.0533)$	0.27	Process error standard deviation	(Independent)
$\sigma_{O,add}$	Half-Normal	0	0.2	Additional estimated observation error	(Independent)
Catch Likelihood					
ν_{catch}	Gamma	Shape = 2	Rate = 0.1	Student-t degrees of freedom for catch	(Independent)

Table 3: Model summary

Model	Max $\hat{R}$	Min ESS	Divergent	Tree Depth	BFMI Issues	Index RMSE	Mohn's ρ	MASE	Coverage D/D_{MSY}	Coverage U/U_{MSY}
0001	1.009	724	0	0	0	0.28	0.006	0.966	100%	100%
0002	1.009	674	0	0	0	0.27	-0.005	0.863	100%	100%
0003	1.008	728	0	0	0	0.265	-0.026	0.779	100%	100%

Notes:

0001 corresponds to the 0001-2024cpueExPrior model

0002 corresponds to the 0002-2024cpueDepPrior model

0003 corresponds to the 0003-2024cpueFPrior model

- Max $\hat{R}$: Maximum potential scale reduction factor across all parameters (should be < 1.01)
- Min ESS: Minimum effective sample size across all parameters (should be > 500)
- Divergent: Number of divergent transitions during sampling (should be 0)
- Tree Depth: Whether maximum tree depth was exceeded during sampling
- BFMI Issues: Whether low Bayesian Fraction of Missing Information was detected
- Index RMSE: Root mean squared error for fit to standardized CPUE index
- Mohn's ρ : Retrospective bias statistic (values close to 0 indicate low bias)
- MASE: Mean Absolute Scaled Error from hindcast cross-validation (< 1 indicates model outperforms naïve predictor)
- Coverage D/D_{MSY} : Percentage of retrospective peels where relative depletion estimates fall within full model credible intervals
- Coverage U/U_{MSY} : Percentage of retrospective peels where relative exploitation rate estimates fall within full model credible intervals

Table 4: Posterior estimates with 95% credible intervals for all BSPM parameters

Model	$\log(K)$	R_{Max}	σ_P	$Shape\ (n)$	σ_F	$\sigma_{O,add}$	σ_O	x_{1988}
0001	13.8 (13.2-14.6)	0.27 (0.13-0.76)	0.07 (0.04-0.11)	1.97 (1.62-2.37)	0.03 (0.01-0.07)	0.10 (0.04-0.19)	0.27 (0.20-0.35)	0.87 (0.59-1.06)
0002	13.7 (13.2-14.5)	0.23 (0.12-0.56)	0.07 (0.04-0.12)	1.96 (1.62-2.35)	0.04 (0.02-0.08)	0.09 (0.03-0.17)	0.26 (0.20-0.34)	0.75 (0.56-0.94)
0003	13.3 (12.8-14.0)	0.23 (0.13-0.45)	0.06 (0.03-0.10)	1.97 (1.65-2.38)	0.07 (0.03-0.13)	0.09 (0.03-0.16)	0.26 (0.19-0.33)	0.67 (0.50-0.89)

Parameter Definitions:

0001 corresponds to the 0001-2024cpueExPrior model

0002 corresponds to the 0002-2024cpueDepPrior model

0003 corresponds to the 0003-2024cpueFPrior model

- $\log(K)$: Natural logarithm of carrying capacity (total numbers)
- R_{Max} : Maximum intrinsic rate of population increase (per year)
- σ_P : Process error standard deviation
- $Shape\ (n)$: Pella-Tomlinson production function shape parameter
- σ_F : Fishing mortality variability standard deviation
- $\sigma_{O,add}$: Extra estimated observation error component
- σ_O : Total observation error (input + additional)
- x_{1988} : Population depletion in 1988 (year 37 of model period)

10 Figures




Figure 1: Annual catch (numbers; individuals) of striped marlin and nominal longline effort (hooks fished; thousands) in the Southwest Pacific Ocean (1952-2022).




Figure 2: Standardized index (black points), and observation error (black bars) for alternative model indices. The trend in the indices is shown in blue using a loess-smooth.




Figure 3: Prior pushforward check showing simulated population depletion trajectories under different filtering scenarios. Lines represent population trajectories colored by maximum intrinsic rate of increase (R_{Max}). Panels show progressively stricter filtering criteria from left to right.




Figure 4: Prior distributions for maximum intrinsic rate of population increase (R_{Max}). Upper panel shows baseline filtered values (gray histogram) with fitted lognormal distribution (red line). Middle panel shows extreme filtered values with fitted distribution (dotted red line). Lower panel compares original simulation values (gray), viable populations (blue), and final lognormal priors (red).




Figure 5: Observed versus predicted mean weights from New Zealand recreational fishery data. Points show observed mean weights with error bars, colored lines show model predictions from the transitional mean weight model colored by relative spawning stock biomass (SSB) depletion.




Figure 6: Prior distribution for relative depletion in 1988 derived from New Zealand recreational weight data. Light blue histogram shows depletion estimates from biological parameter combinations that successfully fit the transitional mean weight model. Blue dashed line shows kernel density estimate, red line shows fitted lognormal distribution used as prior in the depletion-constrained BSPM.




Figure 7: Prior pushforward check for population depletion trajectories under different filtering scenarios. Each line represents a simulated population trajectory colored by fishing mortality variability (σ_F).




Figure 8: Simulated removals from the fishing mortality variability (σ_F) prior pushforward analysis. Solid blue line indicates the median simulated removals with 95th percentile (shaded region). Observed removals are shown by the solid black line.




Figure 9: Prior distributions for fishing mortality variability (σ_F) under different filtering scenarios. Top three panels show progressively stricter filtering (baseline, extreme, catch-based) with fitted distributions (red lines). Bottom panel compares all distributions: original simulation (gray), viable populations (blue), final priors (red), and prior used in models 0001 and 0002 (green).




Figure 10: Time series of observed catch (black dots with uncertainty bounds) and model-predicted catch for three Bayesian surplus production models fitted to SWPO striped marlin data from 1950-2022. The left panel shows model `0001-2024cpueExpPrior` (baseline model with expert priors), the central panel shows model `0002-2024cpueDepPrior` (depletion-constrained model incorporating 1988 depletion prior from New Zealand recreational weight data), the right panel shows model `0003-2024cpueFPrior` which assumes a less restrictive prior on fishing mortality variability. Colored lines represent posterior median predictions with shaded uncertainty bands showing 95% credible intervals.




Figure 11: Probability-integral transform (PIT) style residuals from each model (colored points) fit to the standardized index.




Figure 12: Standardized index (black points), observation error (black bars), and posterior predicted model fits (colored lines) with associated 95% credible interval (colored shading).




Figure 13: Retrospective analysis for all models with respect to time series of depletion D_t , depletion relative to depletion at MSY D_t/D_{MSY} , exploitation rate U_t , and exploitation rate relative to the rate of exploitation that produces MSY U_t/U_{MSY} . The base model with data included through 2022 (black line – median; dark shading – 50% credible interval; light shading – 95% credible interval) is shown relative to the retrospective models. Colored lines correspond to the last year of index data and the colored point indicates the estimate in the last year of the retrospective peel.




Figure 14: Standardized indices of relative abundance used in the Bayesian Surplus Production Model. Open circles show observed values (standardized to mean of 1; black horizontal line) and the vertical bars indicate the observation error (95% confidence interval). One year ahead ‘model-free’ hindcast predictions are shown by the colored lines, where the color indicates the last year of index data seen by the model. The predicted value is shown one year-ahead with the colored point.




Figure 15: Posterior parameter distributions (solid lines) for leading parameters ($\log(K)$, R_{Max} , σ_P , shape (n), σ_F , $\sigma_{O,add}$, and $\sigma_{O,sc}$), relative to their assumed prior distributions (dotted lines) for all models.




Figure 16: Distributions of biological parameters (maximum age max_age , natural mortality reference M_ref , length at birth $L1$, asymptotic length $L2$, von Bertalanffy growth coefficient vbk , length at 50% maturity $l50$, sex ratio at birth sex_ratio , coefficient of variation in length cv_len , weight-length relationship parameters $weight_a$ and $weight_b$, and steepness h) used in the rmax sub-sampling process. Original biological prior distribution (orange shading), distribution based on extreme filtering criteria following the prior pushforward analysis (green shading), and final sub-sampled distribution selected to match the posterior R_{Max} distribution (blue shading).




Figure 17: Posterior time series estimates (1952-2022) for key population metrics across all three BSPM models. Solid lines show median estimates with 95% credible intervals (shading). Metrics include depletion (D), population (P), relative depletion (D/D_{MSY}), relative fishing mortality (F/F_{MSY}), removals, and process error.




Figure 18: Ten-year stochastic projections (2023-2032) under status quo catch scenario. Colored lines show median projections with 95% credible intervals for all three models. Projections assume average 2018-2022 catch levels with resampled process error. Black vertical line indicates start of projection period. Metrics include depletion (D), population (P), relative depletion (D/D_{MSY}), relative fishing mortality (F/F_{MSY}), removals, and process error.

11 Technical Annex

Additional technical detail of population dynamics components used to solve the Euler-Lotka equation for R_{Max} .

11.1 Equilibrium Age-Structured Population Dynamics

11.1.1 Survival

Survival schedule was calculated assuming constant natural mortality:

$$l_a = \begin{cases} 1 & \text{if } a = 1 \\ l_{a-1} \times \exp(-M_{ref}) & \text{if } 1 < a < A_{Max} \\ \frac{l_{A_{Max}-1}}{1 - \exp(-M_{ref})} & \text{if } a = A_{Max} \text{ (plus group)} \end{cases}$$

11.1.2 Fecundity

Reproductive output incorporated biological constraints:

$$f_a = \frac{\psi_a \times W_a \times s}{\rho}$$

where: - ψ_a = proportion mature at age a - W_a = weight at age a (proxy for reproductive capacity) - s = sex ratio (proportion female) - ρ = reproductive cycle (years between spawning events)

Maturity-at-age was derived from length-based logistic maturity:

$$\psi_L = \frac{\exp(a_{mat} + b_{mat} \times L)}{1 + \exp(a_{mat} + b_{mat} \times L)}$$

where $b_{mat} = -a_{mat}/L_{50}$, then integrated over the length-at-age distribution to obtain ψ_a .

Length-at-age followed the Francis parameterization:

$$L_a = L_1 + (L_2 - L_1) \times \frac{1 - \exp(-k(a - age_1))}{1 - \exp(-k(age_2 - age_1))}$$

with length variability modeled using probability density functions linking length bins to ages.

Weight-at-age used the allometric relationship:

$$W_a = a_w \times L_a^{b_w}$$

11.1.3 Stock-Recruitment Relationship

The slope at the origin of the stock-recruitment curve was:

$$\alpha = \frac{4h}{(1 - h)\phi_0}$$

The calculation incorporated steepness (h) through the Beverton-Holt stock-recruitment relationship. Spawners-per-recruit in the unfished state was:

$$\phi_0 = \sum_{a=1}^{A_{Max}} l_a f_a$$

This α parameter links the intrinsic rate of increase to recruitment compensation, ensuring that R_{Max} estimates reflect realistic population productivity under the assumed stock-recruitment dynamics.

Successful simulations (those yielding $R_{Max} > 0$ and $R_{Max} < 1.5$) were filtered to create a plausible prior distribution (Figure 3). The resulting distribution was fitted to a lognormal prior for use in the BSPM (Figure 4).