

Assignment 1

NYALAPOGULA MANASWINI(CS20btech11035)

March 15, 2021

QUESTION 3.3:

Suppose X has a binomial distribution . Show that X = 3 is the most likely outcome.(Hint : P(X = 3) is the maximum among all P(x_i), (x_i= 0,1,2,3,4,5,6).Assume p=0.5

SOLUTION:

Let X be a binomial random variable which has probability p=0.5

then differentiate it with respect to p.

$$\log P = \log\left[\binom{n}{k} \times p^k \times (1-p)^{n-k}\right] \quad (0-5)$$

$$= \log\left[\frac{n!}{(n-k)! \times k!}\right] + k \times \log p + (n-k) \times \log(1-p) \quad (0-6)$$

Differentiate eq (0-6) with respect to p

$$p(X) = 0.5 \quad (0-1)$$

Given number of times event(X) is performed(n)=6

$$n(x) = 6 \quad (0-2)$$

Given probability of event(p)=0.5

Therefore probability that event(X) does not occur is (1-p)=1-0.5=0.5

$$1 - p(x) = 0.5 \quad (0-3)$$

We know that binomial probability

$$P(X = k) = \binom{n}{k} p^k (1-p)^{n-k} \quad (0-4)$$

For P(X=k) to be most likely outcome(highest probability), P(X=k) should be maximum ,where k={0,1,2,3,4,5,6}

To find maximum of P(X=k) ,let us apply logarithm on both sides for equation(0-4) and

$$\begin{aligned} \frac{d \log P}{dp} &= \frac{d \log\left[\frac{n!}{(n-k)! \times k!}\right]}{dp} + k \times \frac{d \log p}{dp} \\ &+ (n-k) \times \frac{d \log(1-p)}{dp} \\ &= 0 + \frac{k}{p} - \frac{n-k}{1-p} \end{aligned} \quad (0-7)$$

To find maximum ,substitute $\frac{d \log P}{dp} = 0$ in (0-7)

$$\begin{aligned} \frac{k}{p} &= \frac{n-k}{1-p} \\ \frac{n-k}{k} &= \frac{1-p}{p} \\ \frac{n}{k} - 1 &= \frac{1}{p} - 1 \\ \frac{n}{k} &= \frac{1}{p} \\ k &= n \times p \end{aligned} \quad (0-8)$$


substituting n=6,p=1-p= $\frac{1}{2}$ in (0-8)

We get $k = 6 \times 0.5 = 3$ Therefore P(X = 3) is maximum, therefore P(X=3) is most likely outcome.

Hence proved.

Submitted by Student unknown on .

Number of times $(X=k)(k=0,1,2,3,4,5,6)$ has occurred out of 10000000 experiments



No course label specified No exercise sheet title specified

No semester specified