30530135	Student Report Ash	C5 ¹ 0 ¹
D.		o138R23C51
0138R22	Name 510138 123551 1386135 5510138 235510 1386135 5510138	200
	N SUCHITRA	cs
		367,2
,BR13C51	3BR23C\$107	22
EX Tit	KRERIMENT 38 CONTROLLEM CONTROLLE	SCS 1 38 R. 2 S. 1
୍ୟ	Pescription You are competing in a basketball contest. In this contest the score for each successful shot depends on both the distance from the backet and the player's position. The half is chet N times successfully. You are given an array A containing the	o1 3V
38R1	age of the state o	, o
, \$P. ? 3C5 ?	distance of a player from basket for N shots. The index of array represents the position of the player. Score is calculated by	3551018
,61	Your task is to find and return an integer value, representing the maximum possible score you can achieve by choosing a contiguous subarray of size K from the given array.	(013BR2°
13	Note:	
3057078	* A subarray is a contiguous part of array.	38R73C51
	* Assume 1 based indexing.	BRIL
222	* The array contains both negative and positive values.	
(013BR25	* Assume the player is standing on a cartesian plane.	7013
`	Input Format	3057013
,8R23C5	- input1:An integer value N representing the number of shots made by the player	
BRIST	- input2 : An integer K representing the size of subarray	013BR25
	in the second se	,01
200	Sample Input	
3057013	Sample input	
	2	A Balley
3BR2	12345	
30		Service Services
	14	30
;	Source Code: ** 3HR1 ³ C5101 3	A CANAL CANA
	arthing and are a second are a second and are a second and are a second are a second and ar	o Barry

```
goals=int(input())
size=int(input())
l=list(map(int,input().split()))
mx=0
for i in range(0,len(1)):
    sub=l[i:i+size]
    k=1
    s=0
    for j in sub:
        s+=(j*k)
        k+=1
        if s>mx:
        mx=s
print(mx)

RESULT

From Table 20

RESULT

RESULT

From Table 20

RESULT

From Table
```