

The KRC Planetary ORBit (PORB) System Users Guide

Hugh H. Kieffer File=-/krc/PUG.tex 2019dec10

December 10, 2019

Contents

1	Introduction	1
2	Outline	2
3	Files input to PORB	2
3.1	Solar system Planets	2
3.2	Minor planets / asteroids	2
3.2.1	Web sources	3
3.3	Comets	3
3.4	exoPlanets	4
3.5	Obsolete	4
4	PORBEL: get elements	4
A	Cookbook	6
B	The geometry matrix	7
C	Routine list and call tree	8

This file is a guide to using PORB and is intended for the general user who wants to apply KRC to someplace other than Mars. A separate document *-/krc/porb.tex* describes the design of the PORB system; it should not be needed by the general user.

1 Introduction

This document describes Version 2.2.2 and later; it applies to KRC Version 2.2.1 and later; 2.2.2 includes additional source files *small.tab* and *exoplan.tab*. The intention is to allow a user to create a **geometry matrix**, 7 rigidly formatted lines of numbers, for inclusion in a KRC input file that would apply to any planet, satellite, asteroid, comet or exoplanet the user may wish; see §B.

Version 2 of KRC uses dates relative to J2000.0 = 2000Jan01 noon UTC. To change from version 1 dates, subtract 11545.0

2 Outline

Check *PORBCM.mat* to see if it already contains the matrix you want. If so, simply copy that into the KRC input file. Otherwise proceed....

The executable of **porbm**n and the elements data files must be in the same directory. Copy them all into a directory of your choice.

Determine if the elements files to be used need any edits for a new object. If so, do the edits.

Run PORBMN to create a geometry matrix, see §A; cut-and-paste it into your KRC input file. Then run KRC.

3 Files input to PORB

Orbital element files formatted for the PORB system cover four categories, each file has its own format:

Planets and satellites: one file for orbits and another for spin-axes

Minor (or small) bodies: orbits and pole direction for each object. Two versions.

Comets: Orbit size specified by perihelion distance.

ExoPlanets: Orbit specified by semi-major axis, period and eccentricity; each commonly known. Relative orientation of the body spin axis specified by obliquity and Ls at periapsis, neither is commonly known.

All files contain adequate information in the header to easily add additional objects.

Reading of these files uses Fortran list-directed format, which uses white-space (blank or tab) to separate items and will terminate with a slash '/'. Thus, names and numbers must not have any internal space or /. On each line, anything after the number of items in the READ list is ignored, so comments may be entered. The first body in each file has a "/" just after the items read on each line to indicate how many items are read. All but the planets files contain a skeleton form for adding objects.

3.1 Solar system Planets

standish.tab Standish06=[4] Keplerian elements and rates of change for 9 planets in mean ecliptic and equinox of J2000 valid for 1800 to 2050. This file is valid for 1800 to 2050; it should not need any edits

Planet 10 is Mars derived from Allison00=[1] Table 2.

spinaxis.tab Archinal11=[2] Planets: Direction of the pole and rates of change in ICRF, which differs from J2000 equatorial system by less than 0.1 arcsecond. The planets and a few satellites are included; more satellites can be added.

3.2 Minor planets / asteroids

There are two PORB-system ASCII files that differ slightly in format. Both contain one element per line and include spin-axis.

minor.tab Generally older entries.

small.tab Format designed to allow cut-and-paste from the JPL Small-Body Database Browser. Rotation period in hours. Must get spin axis from other sources.

All entries were created 2013Aug 22 or later.

includes comet Halley.

Look up object with JPL Small-Body Database Browser

<http://ssd.jpl.nasa.gov/sbdb.cgi>

Copy the number and name line. There must be no blanks or / within the name

Append to this line the epoch

copy elements lines e through M

Click on: Physical Parameters and copy 3 columns for the rotation period. E.g.,

```

    rot_per 7.210 h
Append 3 lines for: [data not in the JPL database]
0. / Right Ascension of Pole, J2000 [deg]
0. / Declination of Pole, J2000 [deg]
0. / Prime meridian at epoch [deg]
    Then replace the zeros if you have real values.

```

3.2.1 Web sources

There are several sources on the web. Some I found did not have unambiguous definition of the coordinate systems used; these were established through email exchanges

PDS Small Bodies Node <http://pdssbn.astro.umd.edu/>

Primary source for spin axis. Values are in ecliptic coordinates of equinox 1950

```

PDS Small Bodies Node http://pdssbn.astro.umd.edu/
DATA ARCHIVES> Archived at SBN > by target > Asteroids(all)
> Photometry and Lightcurves > EAR-A-5-DDR-DERIVED-LIGHTCURVE-V13.0
Either download the set or: > Browse > data > lc_spinaxis.tab (230 kb)

```

The explanation of this file is in `lc_spinaxis.lbl`
Then scroll through. They are in numerical order with multiple entries per asteroid;
the first line is 'summary' of the period (hours) and amplitude of the light curve.
4th and 5th columns to the right of column containing 'summary' contains
the 'best guess' of the longitude and latitude of the spin axis for each observer.
You get to choose your favorite!

Minor Planet Center <http://www.minorplanetcenter.net/iau/info/OrbElsExplanation.html>

ArgPeri, Node and Incln are in heliocentric ecliptic J2000.0
Must specify date for elements that is within requested ephemeris range

JPL Small-Body Database Browser <http://ssd.jpl.nasa.gov/sbdb.cgi>

Orbit elements are in heliocentric ecliptic J2000. Rotation period in hours. No spin-axis data.

Ted Bowells database <ftp://ftp.lowell.edu/pub/elgb/astorb.html>

Enormous; more than 1/2 million objects. Frequent updates.
ArgPeri, Node and Incln are in J2000.0, must be ecliptic. No spin-axis data.

3.3 Comets

Elements from Ted Bowell and Lowell observatory

comet.tab Some comets: orbit and spin.

WARNING: Consistency of source orientation systems uncertain
Specify perihelion and eccentricity.
Includes orbit and pole. Mostly compiled in 1980:1992

Current values are available at:

```

http://www.minorplanetcenter.net/iau/Ephemerides/Comets/Soft00Cmt.txt
Columns   F77      Use
1 -   4   i4      Periodic comet number
5         a1      Orbit type (generally 'C', 'P' or 'D')
6 -  12   a7      Provisional designation (in packed form)
15 - 18   i4      Year of perihelion passage
20 - 21   i2      Month of perihelion passage
23 - 29   f7.4    Day of perihelion passage (TT)
31 - 39   f9.6    Perihelion distance (AU)
42 - 49   f8.6    Orbital eccentricity

```

52 -	59	f8.4	Argument of perihelion, J2000.0 (degrees)
62 -	69	f8.4	Longitude of the ascending node, J2000.0 (degrees)
72 -	79	f8.4	Inclination in degrees, J2000.0 (degrees)
82 -	85	i4	Year of epoch for perturbed solutions
86 -	87	i2	Month of epoch for perturbed solutions
88 -	89	i2	Day of epoch for perturbed solutions
92 -	95	f4.1	Absolute magnitude
97 -	100	f5.1	Slope parameter
103 -		a	Name

J2000.0 was confirmed to be heliocentric ecliptic.

3.4 exoPlanets

A few objects grabbed from the literature

exoplan.tab Major differences from solar system objects are:

Need both semi-major axis and orbital period.

Direction of spin axis is specified by planet obliquity and L_S of periapses

Both these are rarely known, variations are illustrative.

A skeleton for additions is at the bottom of the file.

porbmn run will print an estimated factor for the “solar constant” . This value times the normal SOLCON in KRC input should replace SOLCON. Currently the PORB system does not make use of the host star spectral type.

The length of the synodic day (assuming prograde motion) is also printed; it should be used as PERIOD in the KRC input.

One list of stars with planets is: http://en.wikipedia.org/wiki/List_of_exoplanetary_host_stars

3.5 Obsolete

sturms.tab Sturms71=[5] earliest source

seidelOld.tab Seidelmann74=[3] without Mars updates

seidel.tab Seidelmann74=[3] Mars updated to 2008

MPcomets.tab =[] List of 251 objects from

<http://www.minorplanetcenter.net/iau/Ephemerides/Comets/Soft00Cmt.txt>

Saved 2013 Jun 19

4 PORBEL: get elements

Will read elements and spin-vector data for one object from a (or two) data file, move several values into common and return others.

For planets, reads first the orbital elements and adjusts to the requested epoch.

Standish tables list the following with respect to the J2000 ecliptic system, in order, at J2000.0 and their derivative in time:

- 1 a : semi-major axis [au, au/century]
 - 2 e : eccentricity [radians, radians/century]
 - 3 I : inclination [degrees, degrees/century]
 - 4 L : mean longitude [degrees, degrees/century]
- $L = \Omega + \omega + M$ is a compound angle measured in two planes:
From VE to node in plane of reference

- then to periapsis and on to mean anomaly in orbital plane
- 5 ϖ : longitude of perihelion [degrees, degrees/century] ($\varpi = \Omega + \omega$)
 ϖ is a compound angle measured in two planes:
 ω , the argument of periapsis, measured in the plane of orbit
- 6 Ω : longitude of the ascending node [degrees, degrees/century]
measured in the reference plane.

References

- [1] M. Allison and M. McEwen. A post-Pathfinder evaluation of areocentric solar coordinates with improved timing recipes for Mars seasonal/diurnal climate studies. *Plan. Space Sci.*, 48:215–235, February 2000. See <http://www.giss.nasa.gov/tools/mars24/help/algorithm.html>.
- [2] B. A. Archinal, M. F. A’Hearn, E. Bowell, A. Conrad, G. J. Consolmagno, R. Courtin, T. Fukushima, D. Hestroffer, J. L. Hilton, G. A. Krasinsky, G. Neumann, J. Oberst, P. K. Seidelmann, P. Stooke, D. J. Tholen, P. C. Thomas, and I. P. Williams. Report of the IAU Working Group on Cartographic Coordinates and Rotational Elements: 2009. *Celestial Mechanics and Dynamical Astronomy*, 109:101–135, 2011. Available online at <http://www.dtic.mil/cgi-bin/GetTRDoc?AD=ADA538254>.
- [3] P. K. Seidelmann, L. E. Doggett, and M. R. Deluccia. Mean elements of the principal planets. *Astron. Jour.*, 79:57–+, January 1974.
- [4] E.M. Standish and J.G. Williams. Keplerian elements of the approximate positions of the major planets. In *Explanatory supplement to the Astronomical Almanac*. online at <http://iau-comm4.jpl.nasa.gov/keplerformulae/kepform.pdf>, 2006. Section 8.10.
- [5] F. M. Sturms Jr. Polynomial expressions for planetary equators and orbit elements with respect to the mean 1950.0 coordinate system. Jet Propulsion Laboratory Technical Report, 32-1508, 1971.

A Cookbook

A Cookbook to running the KRC Planetary ORBit system: PORB

cd into the directory containing the probmn executable and the *.tab files

> porbmn To start the main program

Will be asked where to send output; normally enter: T <CR>

Will be asked which of 5 options:

Enter: 1

Will be asked which source data.

Normally enter : 1 (planets) or 4 for asteroids or 6 for exoplanets

Will be asked which item in the file. Should have looked at the chosen elements file to know this

Enter the proper number. E.g., 4 is Mars

Only if doing planets, will be asked for the epoch in centuries after 2000.

E.g., for 2012, enter 0.12

[Version 3+ only Will be asked which Debug print, enter 0 or /]

At this point, the calculations are done and a brief report will be appear

If and only if doing Planets, will be asked for which pole position

a / will use the planet pole, else, enter a satellite name

If that is not in the spinaxis table, will use the planets pole.

If doing exoPlanets, then note the solar power factor and the synodic day.

Will be asked which of 5 options:

4 will print the values in vector format and come back to this same prompt

2 Will begin the required output process: Then

Will be asked to read or save. Enter: 2

Will be asked output format. Enter: 2

Will be asked for an output file name:

A single "/" with no leading spaces will use the default "PORBCM.mat" and
append the latest set to whatever is already in that file.

Else enter a new file name and <CR>

Will be asked which of same 5 options:

Enter: 0 to quit OR to another object.

You then copy the 7 lines in the output file into a KRC input file.

B The geometry matrix

30 items are loaded into the common PORBCM. These are listed in *porbc8m.f*, extracted below (2019dec04)

```

      REAL*8 PLANUM ! 1 Body number. 100*fileNumber+itemNumber
      &,TC           ! 2 Time in centuries from reference date (2000.0)
      &,RODE         ! 3 Longitude of the rising node, radian
      &,CLIN         ! 4 Inclination, radian
      &,ARGP         ! 5 Argument of periapsis, radian
      &,XECC         ! 6 Eccentricity
      &,SJA          ! 7 Semi-major axis (Astronomical Units)
      &,EOBL         ! 8 Obliquity of the Earth's axis, radians
      &,SFLAG        ! 9 Input system flag 0:default, +1: pole=ec +10:orb=eq
      &,ZBAA         ! 10 Body pole: declination, radian in J2000 eq.
      &,ZBAB         ! 11 " " " Right Ascension, radian in J2000 eq.
      &,WDOT         ! 12 Siderial rotation rate; degrees/day (used only in reconc.f)
      &,W0           ! 13 Position of prime meridian at 2000.0, degree

```

C Above are from orbital element tables Below are derived

```

      REAL*8 OPERIOD ! 14 Period of the orbit (days)
      &,TJP          ! 15 J2000 Date of perihelion
      &,SIDAY        ! 16 Siderial rotation period of body (hours)
      &,spar17       ! 17 unused
      &,TAV          ! 18 True anomaly at spring equinox, radians
      &,BLIP         ! 19 Body obliquity, radians
      &,PBUG         ! 20 Debug Flag
      &,spar21       ! 21 Unused
      &,BFRM(9)      ! 22 Rotation matrix from orbital to seasonal

```

C --- end of the 30 real variables generated by PORB system

C Routine list and call tree

Call tree produced by IDL forcalltree

i	name	NumTimesCalled
	PORBMN	
0	CATIME	1
1	PORBIG	1
2	PORBEL	1
3	UPCASE	1
4	COCOMC	1
5	ROTV	1
6	ROTVEC	1
7	COCOCM	1
8	YMD2J2	1
9	ROTORB	1
10	ROTSHO	3
11	ROTAX	1
12	ROTCOL	1
13	COCOSC	1
14	VDOT	1
15	TRANS3	1
16	ROTVEC	1
17	VCROSS	2
18	VNORM	1
19	VSHOW	4
20	ROTEST	1
21	PORBIO	1
22	EPHEMR	1
23	PORBIT	1
24	ORBIT8	1
25	ECCANOM8	1
26	VNEG	1
27	ROTVEC	1
28	COCOCS	1
29	CALDATE	1
30	CALDAT	1
31	JULDAY	1
32	SPCREV	1
33	PRTPCOM	1
34	ROTSHO	1
35	ROTEST	1

Unfound= cococm cococs cocomc cocosc rotax rotcol rotest rotorb rotsho rotv rotvec trans3
vcross vdot vneg vnorm vshow

porbm.f PROBMN Main program of planetary orbit (PORB) software system. R*8
eccanom8.f ECCANOM Iterative solution of Kepler's equations for eccentric orbit.
ephemr.f EPHEMR prints orbital position and date table. PORB system R8
orbit8.f ORBIT Compute radius and coordinates for elliptical orbit. DefPrec.
porbel.f PORBEL read planetary orbital element file, compute basic constants
porbig.f PROBIG Read orbital elements via PORBEL. Initiate porbcm
porbio.f PORBIO read/write porb common to disk file with name = porbcm.dat
porbit.f PORBIT Converts between date and Ls. Also returns DAU and Sdec R*8
prtpcom.f PRTPCOM Print (write) the PORB system version 3 common to IOP
spcrev.f SPCREV: returns spacecraft revolution number for Viking
Math families

cocodp8.f COCODP8 General coordinate conversion package, 8 routines. DP
rotmdp8.f ROTMDP General 3-dimension rotation matrix geometry package. DP, 18 routines
vadddp8.f VADD General vector arithmetic package, 13 routines

Utility

caldat.f CALDAT Convert Julian date to year,month,day. From NumRec
caldate.f CALDATE convert MJD or julian date to year, month, day, day-of-week/year
catime.f CATIME Returns current date & time as " yyyy mon dd hh:mm:ss "
julday.f JULDAY Compute integer Julian day from month, day,year, From NumRec
upcase.f UPCASE Converts all characters to uppercase.