

# 1 Introduction

## 1.1 Outline of proposed research project

The text Counterexamples in Topology by Steen and Seebach has been a fabulous resource for students and researchers in Topology since its publication in 1970. The book was the product of an undergraduate research project funded by NSF and supervised by Steen and Seebach (and including then student Gary Gruenhage) to systematically survey important topological counterexamples. More recently James Dabbs has implemented a database on Github based on the Steen and Seebach textbook called Pi-Base (see <https://topology.pi-base.org/>) and it is currently being maintained by Dabbs and Stephen Clontz. This resource has great potential to both researchers and advanced undergraduate and graduate students at the start of their research careers. There are still big gaps in the database's subject matter, especially in relation to research in and around Frechet-Urysohn spaces. There is a significant body of work, and especially interesting counterexamples, concerning Michael's class of bisequential spaces, Arhangel'skii's alpha-i spaces and several game theoretic formulations of convergence which do not yet appear in the Pi-Base. The project has two goals. The first, and most accessible, is to give a systematic survey of the recent research which will be implemented into the Pi-Base database. The second half of the project will be devoted to open problems related to a recent class of examples defined from ladder systems (more generally on so called square-sequence) described in two

## 2 Meeting Log

### Monday April 21

1. Frechet Fan -  $S_\omega$ :  $\omega \times (\omega + 1)/\omega \times \{\infty\}$  (i.e.  $\omega \times (\omega + 1)$  with the points at infinity identified). Show that
  - $S_\omega$  is not first countable ✓
  - $S_\omega$  is Fréchet. ✓
2. Product of Fréchet spaces not always Fréchet: take  $(\omega + 1) \times S_\omega$ . Let  $A = \{(m, (m, n)) : m, n \in \omega\}$ . Show that
  - $(\omega + 1, \infty) \in \overline{A}$  ✓
  - No sequence in  $A$  converges to  $(\omega + 1, \infty)$ . ✓
3. Right way to think about sequences:  $A \subseteq X$  converges to  $a \in X$  if  $|A| = \aleph_0$  and for all neighbourhoods  $U_x \subseteq X$ ,  $|A \setminus U_x| < \aleph_0$ .
4. Right way to think about Fréchet space: take sequential closure once same as closure.
5. Another exercise: which  $\alpha_i$  properties does  $S_\omega$  have? ✓

### Thursday May 1

1. Go over example 3.9; why is  $S_\omega$  Fréchet? ✓
2. Give example of space that is  $\alpha_2$  but not  $\alpha_1$  ✓,  $\alpha_3$  but not  $\alpha_2$  etc.
3. Under which set theoretic assumptions do such examples exist? Look at the paper by Nyikos: Subsets of  ${}^\omega\omega$  and the Fréchet Urysohn and  $\alpha_i$  properties. **Good grasp at sections 1, 2, 5; partial understanding section 3; skipped section 4, 6.**
4. For example it is consistent that both  $\alpha_1$  and countable imply first countable, and  $\alpha_2$  and countable imply  $\alpha_1$ , yet there exists a countable  $\alpha_2$  space that is not first countable.

5. We talked a little bit about topological groups (a topological space equipped with group operation that is continuous) and some of the nice structure they have: the group operation being continuous and invertible implies  $G \times G \rightarrow G$  is always a homeomorphism, in particular  $aG$  is the homeomorphic image of  $G$  by left multiplication of  $a \in G$ , which in often cases lets us define a nbhd base at the identity and "send" it to all points of  $G$  in order to define the topology.
6. Look into what it takes to contribute to the pi-base, in particular add the  $\alpha_i$  spaces. **Forked database to github account; properties are stored as markdown text files, should not be difficult to add properties as well as if/then.**  
 Things to add:
  - $\alpha_i$  and if  $\alpha_i$  then  $\alpha_{i+1}$
  - bisequential
  - v-space (w-space exists already)
  - if 1st countable then  $\alpha_1$  Frechet
  - if bisequential then Frechet
  - w-space iff Frechet and  $\alpha_2$
7. Some more exercises
  - $X$  is  $\alpha_1$  and Fréchet,  $Y$  is Fréchet, then  $X \times Y$  is Fréchet. **this is not true, take  $(\omega + 1) \times S_\omega$ .**
  - $\alpha_2$  is equivalent to  $A \cap B \neq \emptyset$  for all  $A \in \xi$  whenever  $\xi$  is countable collection of sequences at  $x$ . ✓

## Thursday May 15

1. When is the square of an  $\alpha_i$  space still  $\alpha_i$ ? What  $\alpha$  properties does  $\alpha_i \times \alpha_j$  have?
2. I asked for an example of a tower, and Paul proved that towers exist, but said that theres not really specific constructions of towers.
3. historical note: the cardinals that lay between  $\omega_1$  and  $\mathcal{P}(\omega)$  came from attempts to solve CH, for example, if  $\mathfrak{t} \leq \mathfrak{c}$  then, finding the cardinality for  $\mathfrak{t}$  would help in finding cardinality of  $\mathfrak{c}$ .
4. Proximal game
5. Uniformities
6. Homogeneity in Top group used to determine uniformity for player 1 in prox game
7. FUF and 2FUF
8. exercises:
  - prove lemmas used in existence proof of towers.
  - If  $\{A_\alpha : \alpha \subseteq \gamma\}$  is a tower and  $\{\alpha_\xi : \xi < \lambda\}$  is increasing and cofinal in  $\gamma$  then  $A_{\alpha_\xi} : \xi < \lambda$  is a tower.
  - $S_\omega$  is neither FUF or 2FUF
  - Proximal game is equivalent to Gruenhage game.
9. note: send Paul changes to databse before pushing to github

## 3 Set Theory

### 3.1 Some Interesting Cardinals

#### 3.1.1 Subsets of ${}^\omega\omega$ and Almost Disjoint Families

**Definition 3.1.** Let  $\mathcal{A}$  be an infinite family of infinite subsets of  $\omega$ . The family  $\mathcal{A}$  is said to be an almost disjoint family (ADF) of subsets of  $\omega$  if  $A \cap B$  is finite for any  $A, B \in \mathcal{A}$  with  $A \neq B$ . By Zorn's lemma we can assume  $\mathcal{A}$  lives inside of a maximal (w.r.t. inclusion) family that is also pairwise disjoint. Such a family is a maximal almost disjoint family (MADF).

**Definition 3.2.** Define an order  $\leq^*$  on  ${}^\omega\omega$  where  $f <^* g$  if  $f(n) < g(n)$  for all but finitely many  $n \in \omega$

**Definition 3.3.** Let  $(A, \prec)$  be a linearly ordered set. We say that  $B \subseteq A$  is unbounded in  $A$  if there does not exist  $y \in A$  such that  $x \prec y$  for all  $x \in B$ . We say that  $C \subseteq A$  is cofinal (or dominating) in  $A$  if for all  $x \in A$  there exists  $z \in C$  such that  $x \prec z$ .

**Definition 3.4.**

- $\mathfrak{a} = \min\{|\mathcal{A}| : \mathcal{A} \text{ is infinite MADF}\}$
- $\mathfrak{b} = \min\{|B| : B \text{ is } \leq^* \text{-unbounded subset of } {}^\omega\omega\}$
- $\mathfrak{d} = \min\{|D| : D \text{ is } \leq^* \text{-cofinal in } {}^\omega\omega\}$

Each of  $\mathfrak{a}, \mathfrak{b}, \mathfrak{d}$  are at least  $\omega_1$  but at most  $\mathcal{P}(\omega)$ .

#### 3.1.2 Towers and Pseudo-Intersections

**Definition 3.5.** Define an order on  $\mathcal{P}(X)$  for some set  $X$ . We say  $A \subseteq^* B$  if  $B$  contains all but finitely many points of  $A$ . If  $\mathcal{F} \subseteq \mathcal{P}(X)$  is a collection of (infinite) sets, then  $A$  is a pseudo-intersection of  $\mathcal{F}$  if  $A \subseteq^* F$  for all  $F \in \mathcal{F}$ .

**Definition 3.6.** We call  $\mathcal{T} \subseteq [\omega]^\omega$  a tower if  $\mathcal{T}$  is well ordered by  $\subseteq^*$  and has no infinite intersection.

**Lemma 3.7.** If  $\mathcal{A} = \{A_n : n \in \omega\} \subseteq [\omega]^\omega$  such that for all  $n \in \omega$

$$A_0 \supseteq^* A_1 \supseteq^* \dots \supseteq^* A_n$$

then  $\mathcal{A}$  has infinite pseudo-intersection.

**Theorem 3.8.** *There exists a tower.*

*Proof.* Define  $\langle A_\alpha : \alpha < \mathfrak{c} \rangle$  as follows:

1.  $A_\alpha = \omega \setminus n$  for  $\alpha < \omega$
2. If  $\alpha$  is a successor ordinal, define  $A_\alpha$  such that  $|A_{\alpha-1} \setminus A_\alpha| = \aleph_0$
3. If  $\alpha$  is a limit ordinal then define  $A_\alpha$  such that  $A_\alpha \subseteq^* A_\beta$  for all  $\beta < \alpha$

Suppose  $\gamma$  is given and  $\langle A_\alpha : \alpha < \gamma \rangle$  has been defined. If  $\gamma$  is a successor then define  $A_\gamma$  as in (2). Otherwise  $\gamma$  is a limit ordinal: if (3) isn't possible then  $\langle A_\alpha : \alpha < \gamma \rangle$  is a tower; otherwise define  $A_\gamma$  as in (3) and continue.

This process eventually terminates, otherwise we obtain  $\mathfrak{c}$  many distinct subsets of  $\omega$ , a contradiction.  
TODO: elaborate □

**Definition 3.9.**

- $\mathfrak{t} = \min\{|\mathcal{T}| : \mathcal{T} \text{ is a tower}\}$
- $\mathfrak{p} = \min\{|\mathcal{P}| : \mathcal{P} \text{ is a filter base for a free filter on } \omega \text{ with no infinite pseudo-intersection}\}$

## 4 Topology

### 4.1 Sequential and Fréchet Spaces

**Definition 4.1.** For a topological space  $X$  and any set  $A \subseteq X$ , the *sequential closure* of  $A$  is

$$[A]_{\text{seq}} := \left\{ x \in X : \exists (x_n) \in A \left( \lim_{n \rightarrow \infty} x_n = x \right) \right\}.$$

In general we can repeat this operation recursively  $[[[A]_{\text{seq}}]_{\text{seq}} \dots]_{\text{seq}}$  by which is meant the *total sequential closure* of  $A$ .

**Fact 4.2.** In general it takes at most  $\omega_1$  many iterations of the sequential closure to get a closed set.

**Definition 4.3.** A space  $X$  is said to be Fréchet if  $[A]_{\text{seq}} = \overline{A}$  for all  $A \subseteq X$ .

**Example 4.4.** Let  $X = \omega_1 + 1$  with the order topology.  $X$  is not Fréchet, since any sequence  $(x_n) \in \omega_1$  cannot converge to  $\infty$ , as otherwise  $\omega_1 = \sup\{x_n : n \in \mathbb{N}\}$ , a contradiction.

**Definition 4.5.** A space  $X$  is sequential if any closed set  $A \subseteq X$  is equal to its total sequential closure.

Then a space that is Fréchet is also sequential. The following example shows that the converse is not true.

**Example 4.6.** Let  $X^* = \omega \times (\omega + 1)$  be given the order topology and let  $X = X^* \cup \{\infty\}$  where the neighbourhoods of  $\infty$  are such that there exists  $p \in \omega$  such that  $|\{(m, n) : m > p, n \in \omega + 1\} \setminus U_\infty| < \aleph_0$ . Then  $X$  is sequential but not Fréchet. To see this, note that for all  $m \in \omega$  the sequence  $A_m = \{(m, n) : n \in \omega\}$  converges to  $(m, \omega + 1)$  and moreover  $B = \{(m, \omega + 1) : m \in \omega\}$  is a sequence that converges to  $\infty$ . Then  $A = \bigcup_{m \in \omega} A_m$  is such that  $[[A]_{\text{seq}}]_{\text{seq}} = X$ , hence  $X$  is sequential. On the other hand there is no sequence in  $A$  that converges to  $\infty$ . Suppose there were, say some  $\gamma \rightarrow \infty$ . Then for all  $m \in \omega$ ,  $U_m = X \setminus \{(m, n) : n \in \omega + 1\}$  is a neighbourhood of  $\infty$  such that  $|\gamma \setminus U_m| < \aleph_0$ . Hence  $\gamma$  has only finitely many terms belonging to each column. If  $\alpha_m = \max\{\gamma \cap \{(m, n) : n \in \omega\}\}$ , then  $U = X \setminus \bigcup_{m \in \omega} \{(m, n) : n \leq \alpha_m\}$  is a neighbourhood of  $\infty$  disjoint from  $\gamma$ , a contradiction. Hence  $X$  is not Fréchet.

**Proposition 4.7.** If  $X$  is first countable then  $X$  is Fréchet.

*Proof.* Let  $A \subseteq X$  and let  $x \in \overline{A}$ . Then  $x$  has a countable neighbourhood base  $N_x$  such that  $U \cap A \neq \emptyset$  for all  $U \in N_x$ . Enumerating the neighbourhoods of  $x$  as  $U_1, U_2, \dots$  then the sequence  $(x_n)_{n \geq 1}$  where  $x_n \in U_n \cap A$  for each  $n \in \omega$  is such that  $(x_n)_{n \geq 1}$  converges to  $x$ .  $\square$

The following example shows that the converse is not true.

**Example 4.8 (Fréchet Fan).** Let  $S_\omega$  be the quotient of  $\omega \times (\omega + 1)$  obtained by identifying all the points  $\{(m, \omega + 1) : m \in \omega\}$  as  $\infty^*$ . More precisely  $S_\omega$  has the quotient topology induced by the map  $h : \omega \times (\omega + 1) \rightarrow S_\omega$  where  $h(x) = x$  for all  $x \in \omega \times \omega$  and  $h(x) = \infty^*$  for all  $x \in \omega \times \{\omega + 1\}$ . Then  $S_\omega$  is Fréchet but not 1st countable.

To see that  $S_\omega$  is Fréchet, let  $A \subseteq S_\omega$  such that  $\infty \in \overline{A}$ .  $A$  must meet at least one column of  $S_\omega$  in an infinite set, otherwise we could find a nbhd of  $\infty^*$  disjoint from  $A$ . Then  $A$  restricted to that column will be a convergent sequence to  $\infty$ .

Now assuming that  $S_\omega$  was countable, we would have a countable neighbourhood base at  $\infty^*$ . For each  $k \in \omega$  let  $B_k = \bigcup_{m \in \omega} \{m\} \times (f_k(m), \infty^*]$  for some  $f_k : \omega \rightarrow \omega$  determining the startpoints of each interval. Suppose  $\mathcal{B} = \{B_k : k \in \omega\}$  is a base at  $\infty^*$ , then let  $f^* : \omega \rightarrow \omega$  be defined by  $f^*(m) = f_m(m) + 1$  for all  $m \in \omega$ . Letting  $B^* = \bigcup_{m \in \omega} \{m\} \times (f^*(m), \infty^*]$  then  $B^*$  is an open neighbourhood of  $\infty^*$  but it is clear by construction that  $B_k \not\subseteq B^*$  for all  $k \in \omega$ . Hence  $\mathcal{B}$  cannot be a neighbourhood base and  $S_\omega$  is not first countable.

As the following example shows, the product of Fréchet spaces need not be Fréchet.

**Example 4.9.** Let  $X = (\omega + 1) \times S_\omega$ , and consider the set  $A = \{(m, (m, n)) : m, n \in \omega\}$ . If  $\infty^*$  is the identified point of  $S_\omega$ , let  $\infty = \{\omega + 1\} \times \infty^*$ . Then  $\infty \in \bar{A}$  but  $\infty \notin [A]_{\text{seq}}$ . The open neighbourhoods of  $\infty$  are of the form  $(\alpha, \omega + 1] \times (\bigcup_{m \in \omega} \{m\} \times (f(m), \infty^*])$ , which clearly always has non empty intersection with  $A$ . Hence  $\infty \in \bar{A}$ . To see that  $\infty \notin [A]_{\text{seq}}$ , suppose  $\gamma$  is a sequence in  $A$  that converges to  $\infty$ . Since the sets  $U_k = (k, \omega + 1] \times (\bigcup_{m \in \omega} \{m\} \times (f(m), \infty^*])$  are open neighbourhoods of  $\infty$  it must be the case that  $|\gamma \setminus U_k| < \aleph_0$  for all  $k \in \omega$ . Thus  $\gamma \cap (\{k\} \times \{k\} \times (1, \omega + 1])$  is finite for every  $k$ . Let  $h : \omega \rightarrow \omega$  be defined by  $h(k) = \max\{\pi_3(\gamma \cap (\{k\} \times \{k\} \times (1, \infty^*]))\} + 1$  for  $k \in \omega$ . Pictorially,  $h$  is picking the point on each spine beyond which no elements of  $\gamma$  exist. Thus

$$W = (1, \omega + 1] \times \left( \bigcup_{n \in \omega} \{n\} \times (h(n), \infty^*] \right)$$

is an open neighbourhood of  $\infty$  which by construction is disjoint from  $\gamma$ . Hence  $\gamma$  cannot converge to  $\infty$  showing that  $X$  is not Fréchet.

## 4.2 Finite Frechet-Urysohn spaces

**Definition 4.10.** A network of a space  $X$  is a collection  $\mathcal{F} \subseteq \mathcal{P}(X)$  such that for all  $x \in X$  and for all nbhds  $U_x$  there is  $F \in \mathcal{F}$  such that  $x \in F \subseteq U_x$ .

**Definition 4.11.**  $X$  is Fréchet-Urysohn Finite (FUF) if whenever  $\mathcal{F}$  is a network in  $X$  there exists  $\{F_n : n \in \omega\} \subseteq \mathcal{F}$  such that for all nbhds  $U$  there exists  $n \in \omega$  such that  $x \in F_n \subseteq U$  for all  $m > n$ .

## 4.3 $\alpha_i$ notions of convergence

**Definition 4.12.** Let  $X$  be a topological space and  $\xi$  be a countable family of sequences converging to a point  $x \in X$ . We say that  $x$  is an  $\alpha_i$  point for  $i = 1, 2, 3, 4$  if there exists a sequence  $B$  such that

- $\alpha_1$ :  $|A \setminus B| < \aleph_0$  for every  $A \in \xi$ ;
- $\alpha_2$ :  $|A \cap B| = \aleph_0$  for every  $A \in \xi$ ;
- $\alpha_3$ :  $|A \cap B| = \aleph_0$  for infinitely many  $A \in \xi$ ;
- $\alpha_4$ :  $A \cap B \neq \emptyset$  for infinitely many  $A \in \xi$ .

Then  $X$  is an  $\alpha_i$  space if every  $x \in X$  is an  $\alpha_i$  point. Note that if a space is  $\alpha_i$  then it is  $\alpha_{i+1}$  for  $i = 1, 2, 3$ .

**Example 4.13.**  $S_\omega$  is not even  $\alpha_4$ . For each  $m \in \omega$  let  $A_m = \{m\} \times (1, \infty)$ . Then  $\xi = \{A_m : m \in \omega\}$  is a countable collection of sequences converging to  $\infty$ . Suppose  $B$  is a sequence that converges to  $\infty$  such that  $A \cap B \neq \emptyset$  for infinitely many  $A \in \xi$ . In particular let  $\alpha \leq \omega$  be such that  $A_i \cap B \neq \emptyset$  for all  $i \in \alpha$  and let  $f : \alpha \rightarrow \omega$  be defined by  $f(k) \in B \cap A_k$  for all  $k \in \alpha$ . Then

$$U = \left( \bigcup_{k \in \alpha} \{k\} \times (f(k) + 1, \infty] \right) \times \left( \bigcup_{m \in \omega \setminus \alpha} \{m\} \times (1, \infty] \right)$$

is such that  $|B \cap U^C| = \aleph_0$ , hence  $B$  does not converge to  $\infty$ .

**Exercise 4.14.**  $X$  is  $\alpha_2$  iff whenever  $\xi$  is a countable collection of sequences converging to  $x$  there exists  $B \rightarrow x$  such  $A \cap B \neq \emptyset$  for all many  $A \in \xi$ .

**Solution.** The forward direction is obvious. Conversely, let  $\xi = \{A_1, A_2, \dots\}$  be a countable collection of sequences converging to  $x$ . For every  $n \in \omega$  let  $\{a_{nm} : m \in \omega\}$  enumerate the elements of  $A_n$  and define  $A_{nm} = A_n \setminus \{a_{n1}, a_{n2}, \dots, a_{n,m-1}\}$  for each  $n, m \in \omega$ . Then the  $A_{nm}$  still converge to  $x$  and  $\mathcal{A} = \{A_{nm} : n, m \in \omega\}$  is a sheaf at  $x$ . By hypothesis there exists a  $B$  converging to  $x$  that meets each  $A_{nm}$ . Thus  $B$  meets each  $A_n$  in an infinite set.

#### 4.4 $\Psi$ -like spaces

**Definition 4.15.** Let  $D$  be a countable set and let  $\mathcal{A} \subseteq \mathcal{P}(D)$  be an ADF. Define a topology on  $X = D \cup \mathcal{A}$  such that the points of  $D$  are isolated and  $D$  is dense in  $X$ . Then for every  $A \in \mathcal{A}$  attach a point  $z_A$  where the nbhds of  $z_A$  consist of all sets of the form  $B(A, F) = \{z_A\} \cup (A \setminus F)$  where  $F \subseteq A$  is finite.

**Proposition 4.16.** Let  $X$  be as above. Then  $X$  is Hausdorff, first countable and locally compact. If  $X$  is constructed as above but  $\mathcal{A}$  is MADF then  $X$  is also pseudo-compact (i.e. all continuous real functions on  $X$  are bounded).

*Proof.* To check Hausdorffness it suffices to show  $Z = X \setminus D$  is Hausdorff. Let  $z_1, z_2 \in Z$ . Then  $z_1, z_2$  correspond to sets  $A_1, A_2$  such that  $A_1 \cap A_2 = I$  is finite. Then  $B(A_1, I)$  and  $B(A_2, I)$  are disjoint open sets containing  $z_1, z_2$  respectively.

To see that  $X$  is locally compact, note that  $\{x\}$  is a compact neighbourhood base for all  $x \in D$ . Local compactness of  $Z$  follows from the fact that by design each  $z_A \in Z$  is the limit of  $A$  viewed as a sequence. In other words, whenever  $B(A_z, F)$  is a basic nbhd of  $z_A$ , any open cover will have a smaller nbhd  $B(A_z, F')$  that omits only finitely many points.  $\square$

**Proposition 4.17.** The one point compactification of a  $\Psi$ -like space of cardinality  $< \mathfrak{a}$  is Fréchet.

#### 4.5 Constructing Examples of $\alpha_i$ Spaces

**Example 4.18.** Let  $D = \omega \times \omega$ , let  $\mathcal{F} = \{f_\alpha : \alpha < \mathfrak{b}\} \subseteq {}^\omega\omega$  be  $<^*$ -well-ordered and  $<^*$ -unbounded, and let  $\mathcal{C} = \{C_n = \{n\} \times \omega : n \in \omega\}$ . Construct a  $\Psi$ -like from  $X = \mathcal{F} \cup \mathcal{C}$ , where we attach compactification points  $\{n\} \times (\omega + 1)$  for each  $C_n$  and compactification points  $p_\alpha$  for each  $f_\alpha$ . Then the resulting one point compactification,  $X + \infty$ , is  $\alpha_2$  and Fréchet but not  $\alpha_1$ .  $X + \infty$  becomes  $\alpha_1$  if  $\mathcal{F}$  is cofinal and well ordered.

#### 4.6 Bisequential Spaces

**Definition 4.19.**

#### 4.7 Topological Groups

#### 4.8 Games