

National Biodiversity Assessment 2025

The status of South Africa's ecosystems and biodiversity

Andrew L. Skowno Carol J. Poole Domitilla C. Raimondo
Kerry J. Sink Lara van Niekerk Linda R. Harris
Krystal A. Tolley Tsungai A. Zengeya

2024-10-12

Table of contents

Preface	8
1 Introduction	9
2 Summary	10
I About	11
3 About the National Biodiversity Assessment	12
3.1 Why do we do the NBA	12
3.2 How do we do the NBA	12
3.3 When is the NBA updated	12
3.4 What the NBA is not	12
3.5 How to cite the NBA	12
3.6 Acknowledgements	12
4 How the NBA is used	13
4.1 Introduction	13
4.2 How the NBA input datasets are used	13
4.2.1 Overview of how NBA input datasets are used	13
4.2.2 Detailing how NBA realm specific input datasets are used?	14
4.2.3 How the NBA findings and key messages can be used	14
4.3 Direct links between NBA indicators and MEAs indicators	14
4.3.1 KBA analysis and indicators?	14
4.3.2 Indicators database to serve NBA and MEAs	14
5 South Africa's biodiversity profile	15
5.1 Overview of South Africa's biodiversity profile	15
5.2 Biodiversity benefits	15
5.2.1 Benefits of biodiversity in this realm?	15

II	Key messages	17
6	South Africa's biodiversity provides benefits to people	18
6.1	Healthy ecosystems are essential for water security	18
6.2	Water flowing into the sea provides multiple benefits to people	18
6.3	Small high-value ecosystem types take up just 5% of South Africa's territory, but provide disproportionate benefits to people	18
6.4	Benefits from fishing are at risk, including food and job security	19
6.5	Climate change is impacting on people and ecosystems; in spite of this, healthy ecosystems can help us adapt to climate change	19
7	South Africa's biodiversity is under pressure, but solutions are at hand	20
7.1	Estuaries and wetlands are the most threatened and least protected ecosystems in South Africa	20
7.2	Coastal biodiversity assets, including beaches, are at risk	20
7.3	Protected areas: investment success in the ocean and on land	20
7.4	Protected areas: providing effective protection for many species	20
7.5	Freshwater fishes are the most threatened species group in South Africa	21
7.6	Trends in threat status show rapid declines in some of South Africa's species, especially freshwater species and butterflies	21
7.7	Areas where pressures are concentrated should be priorities for spatial planning	21
7.8	Biological invasions threaten biodiversity and human wellbeing	21
7.9	Cooperative governance is essential for healthy landscapes and seascapes	22
8	The NBA stimulates work to address knowledge gaps	23
8.1	South Africa's new seamless map of ecosystem types paves the way for improved assessment, planning and monitoring	23
8.2	New indicators developed during the NBA 2018 advance South Africa's ability to report on the status of biodiversity	23
8.3	Evaluation of genetic diversity brings new value to the NBA	23
8.4	Investment in strategic and collaborative biodiversity monitoring programmes is crucial to inform management and decision making and for biodiversity assessments	24
III	Integrated findings	25
9	National Headline Indicators	26
9.1	Overview	26
9.1.1	Red List Index of Ecosystems	26
9.1.2	Protection Level	26
9.2	Species indicators	26
9.2.1	Red List Index of Species	26
9.2.2	Species Protection Level	26

9.3	Genetic diversity	27
9.3.1	Genetic diversity at population level	27
9.3.2	Genetic diversity at landscape level	27
10	Pressures	28
10.1	Synthesis	28
10.1.1	Climate change	28
10.1.2	Land cover change	28
10.1.3	Degradation	28
10.1.4	Alien invasive species	28
10.1.5	Pesticides	28
10.2	Ecological condition overview	29
10.2.1	Ecological condition per realm	29
IV	Findings by realm	31
11	Terrestrial realm	32
11.1	Overview	32
11.2	Input data and method for the terrestrial realm	32
11.2.1	Ecosystem assessments	32
11.2.2	Species assessments	32
11.3	Key drivers and pressures impacting terrestrial ecosystems	32
11.3.1	Habitat loss	32
11.3.2	Land degradation	33
11.3.3	Biological invasions	33
11.3.4	Climate change	33
11.3.5	Biological resource use	33
11.4	Terrestrial ecosystems	33
11.4.1	Map of ecosystem types	33
11.4.2	Overview of South African biomes	33
11.5	Ecological condition	34
11.6	Overview	34
11.6.1	Ecological condition per realm	35
11.7	Red List of Ecosystems	36
11.7.1	Red List of Ecosystems per biome	36
11.7.2	Red List of Ecosystems - Prince Edward Island	37
11.8	Protection Level	37
11.8.1	Protection level per biome	37
11.9	Terrestrial species	39
11.9.1	Utilized terrestrial species resource assessments	39

12 Freshwater realm	40
12.1 Overview	40
12.1.1 Rivers	40
12.1.2 Wetlands	40
12.2 Ecosystems	40
12.2.1 Rivers	40
12.2.2 Wetlands	40
12.2.3 Riparian	41
12.3 Ecological condition	41
12.3.1 Rivers	41
12.3.2 Wetlands	41
12.4 Red List of Ecosystems	41
12.4.1 Red List of Ecosystems - rivers	41
12.4.2 Red List of Ecosystems - wetlands	41
12.5 Protection level	41
12.5.1 Protection level of rivers	41
12.5.2 Protection level of wetlands	42
12.6 Species	42
13 Estuarine realm	43
13.1 Overview	43
13.2 Ecological condition	43
13.2.1 Condition	43
13.2.2 Pressures	43
13.3 Ecosystems	43
13.3.1 Habitats and vegetation	43
13.4 Red List of Ecosystems	44
13.5 Protection Level	44
13.6 Species	44
14 Marine realm	45
14.1 Overview	45
14.2 Input data and method for the marine realm	45
14.2.1 Ecosystem assessments	45
14.2.2 Species assessments	45
14.3 Key drivers and pressures impacting marine ecosystems	45
14.3.1 Habitat loss	45
14.3.2 Degradation	46
14.3.3 Biological invasions	46
14.3.4 Climate change	46
14.3.5 Biological resource use	46
14.4 Ecosystems	46
14.4.1 Ecosystems in the marine realm	46

14.4.2	Ecosystems around Prince Edward Island	46
14.5	Ecological condition	46
14.5.1	Marine pressure datasets	46
14.6	Red List of Ecosystems	47
14.7	Protection level	47
14.8	Species	47
15	Coast cross-cutting realm	48
15.1	Overview	48
15.2	Ecosystems	48
15.2.1	Map of ecosystem types	48
15.3	Ecological condition	48
15.4	Red List of Ecosystems	48
15.5	Protection level	48
15.6	Species	49
V	Species	50
16	Species	51
16.1	Amphibians	51
16.2	Birds	51
16.2.1	Terrestrial	51
16.2.2	Estuarine	51
16.3	Fish	51
16.3.1	Inland aquatic	51
16.3.2	Estuarine	52
16.4	Invertebrates	52
16.4.1	Overview of invertebrate groups assessed to date	52
16.4.2	Bees	52
16.4.3	Branchiopoda	52
16.4.4	Butterflies	52
16.4.5	Crabs	52
16.4.6	Dragonflies	52
16.4.7	Mollusca	52
16.4.8	Spiders	53
16.4.9	Estuary invertebrates	53
16.5	Mammals	53
16.5.1	Terrestrial	53
16.5.2	Inland aquatic	53
16.5.3	Marine	53
16.6	Plants	53
16.7	Reptiles	53

VI Priority actions and responses	54
17 Priority actions for improving future NBAs	55
17.1 Overview	55
17.2 Knowledge Gaps and Research Priorities for Strengthening the NBA	55
17.2.1 Review of NBA 2018 Research, Monitoring and Data Management Priorities	55
17.2.2 NBA 2025 Research, Monitoring and Data Management Priorities	55
17.3 Review of tools used to measure genetic diversity	56
17.3.1 State of DNA barcoding	56
17.3.2 Use of eDNA in South Africa	56
17.3.3 State of SA's Biobanks	56
18 South Africa's responses for managing and conserving biodiversity	57
18.1 Overview	57
18.2 Avoid loss	57
18.3 Protect	57
18.3.1 Protection over time indices?	57
18.4 Restore	57
18.5 Resources for biodiversity conservation	57
18.6 The need for evidence on the effectiveness and impact of responses	58
References	59

Preface



I am proud to present South Africa's National Biodiversity Assessment 2025, a product of high scientific calibre that summarises the state of our biodiversity by drawing on a broad foundation of information compiled by over a hundred authors. This body of knowledge should be used as a basis for policy, planning and decision making regarding the wise use and conservation of our country's biodiversity assets and the management and restoration of ecological infrastructure. **Shonisani Munzhedzi** South African National Biodiversity Institute - Chief Executive Officer

The NBA is a tool not only for the environment sector, but for all sectors of government that share the responsibility for the sustainable development of our landscapes and seascapes, and is prepared as part of the South African National Biodiversity Institute (SANBI) mandate under the National Environmental Management: Biodiversity Act (Act 10 of 2004). South Africa is a special country, with diverse cultures, remarkable geological wealth and exceptional biodiversity, much of which is unique to our nation. With this rich endowment comes the responsibility and challenge of ensuring our species and ecosystems are conserved and used sustainably to the benefit of all South Africans and future generations. This biodiversity wealth gives our people tangible benefits like food, clean water, medicine and materials; it supports agricultural and fisheries production, and helps protect us from natural hazards like floods and droughts; and it provides the basis of a vibrant tourism industry while offering natural spaces for recreational and cultural activities.

Biodiversity is central to South Africa's national objectives of addressing poverty, inequality and unemployment, and supports increased economic growth and improved service delivery for all its citizens. Examining biodiversity in the context of social and economic change shows us that investing in ecological infrastructure is as important as investing in built infrastructure, and safeguarding the delivery of services from ecosystems can support service delivery from all spheres of government.

1 Introduction

This is a book created from markdown and executable code.

See Knuth (1984) for additional discussion of literate programming.

```
1 + 1
```

```
[1] 2
```

2 Summary

In summary, this book has no content whatsoever.

1 + 1

[1] 2

Part I

About

3 About the National Biodiversity Assessment

3.1 Why do we do the NBA

Mandate etc

3.2 How do we do the NBA

Who, number of people

3.3 When is the NBA updated

Modules are updated at different times

3.4 What the NBA is not

Explanation of what NBA is not

3.5 How to cite the NBA

Recommended citation

Reference in scientific publications:

Reference in non-scientific publications:

3.6 Acknowledgements

4 How the NBA is used

4.1 Introduction

The NBA is the primary tool for monitoring and reporting on the state of biodiversity in South Africa. It is prepared as part of the South African National Biodiversity Institute's (SANBI) mandate to monitor and report regularly on the status of South Africa's biodiversity, and is a collaborative effort from many institutions and individuals. The NBA focusses primarily on assessing biodiversity at the ecosystem and species level, with efforts being made to include genetic level assessments. Two headline indicators that are applied to both ecosystems and species are used in the NBA: threat status and protection level. The products of the NBA include seven technical reports, a technical synthesis report and several popular outputs.

The primary purpose of the NBA is to provide a highlevel summary of the state of South Africa's biodiversity at regular points in time, with a strong focus on spatial information. Each NBA builds on decades of research and innovation by South African scientists, and makes that science available in a useful form to users both inside and outside of the biodiversity sector. As a body of work the NBA is not prescriptive; it presents important information that can be adopted by government and civil society in various decision making processes to support socio-economic imperatives, human wellbeing, and the best management and conservation of South Africa's biodiversity.

4.2 How the NBA input datasets are used

4.2.1 Overview of how NBA input datasets are used

add text

4.2.2 Detailing how NBA realm specific input datasets are used?

4.2.2.1 Terrestrial realm

add text

4.2.2.2 Freshwater (Inland aquatic) realm

add text

4.2.2.3 Estuarine realm

add text

4.2.2.4 Marine realm

add text

4.2.2.5 Coast cross-cutting realm

add text

4.2.3 How the NBA findings and key messages can be used

add text

4.3 Direct links between NBA indicators and MEAs indicators

4.3.1 KBA analysis and indicators?

add text

4.3.2 Indicators database to serve NBA and MEAs

add text

5 South Africa's biodiversity profile

5.1 Overview of South Africa's biodiversity profile

South Africa has exceptional biodiversity, characterised by high species richness, high levels of species endemism and a wide variety of ecosystems. South Africa's diversity and richness are not limited to biodiversity. Within its borders are also diverse cultures and languages, and exceptional geological and climatic diversity.

Identified as one of the world's 17 megadiverse nations, South Africa ranks as one of the top ten nations globally for plant species richness and third for marine species endemism. With a landmass of 1.2 million km² and surrounding seas of 1.1 million km², South Africa is among the smaller of the world's megadiverse countries – which together contain more than two-thirds of the world's biodiversity. South Africa also holds three of the world's 35 biodiversity hotspots (a measure of biological diversity combined with vulnerability to threats): the Cape Floristic Region, Succulent Karoo biome, and the Maputaland–Pondoland–Albany centre of endemism.

5.2 Biodiversity benefits

Add text

5.2.1 Benefits of biodiversity in this realm?

Benefits and how to link to the importance of SA's biodiversity

5.2.1.1 Terrestrial realm

Add text

5.2.1.2 Freshwater (inland aquatic) realm

Add text

5.2.1.3 Estuarine realm

Add text

5.2.1.4 Marine realm

Add text

5.2.1.5 Coast cross-cutting realm

Add text

Part II

Key messages

6 South Africa's biodiversity provides benefits to people

#3 Biodiversity provides jobs

South Africa's biodiversity provides substantial employment in a range of sectors (established but incomplete). Continued investment in managing and conserving biodiversity is essential so that jobs that depend on biodiversity can continue to increase.

6.1 Healthy ecosystems are essential for water security

Rivers, wetlands and their catchment areas are crucial ecological infrastructure for water security, often complementing built infrastructure, but the benefits from some of these ecosystems are currently compromised by their poor ecological condition (well established). Water security can be improved through integrated management of natural resources in Strategic Water Source Areas as well as other key catchments, including protection and restoration in some cases.

6.2 Water flowing into the sea provides multiple benefits to people

Freshwater flowing from rivers through estuaries into the sea is not wasted, and is essential for coastal and marine food production, livelihoods, tourism and future climate change resilience (established but incomplete). Through appropriate management, South Africa can maintain the vital freshwater flows that reach the coast.

6.3 Small high-value ecosystem types take up just 5% of South Africa's territory, but provide disproportionate benefits to people

Certain small ecosystem types function as crucial ecological infrastructure and, despite their small footprint, provide multiple benefits to society (established but incomplete). Managing, protecting and restoring these small, high-value.

6.4 Benefits from fishing are at risk, including food and job security

Estuarine and marine ecosystems provide South Africans with food and livelihoods by providing a basis for fishing – whether commercial, subsistence or recreational. Yet many fish stocks are overexploited and many fish species are threatened (well established). While a range of plans are in place to ensure that fisheries are sustainable, better practices to rebuild stocks of priority species are needed, as well as reliable data and sufficient capacity for undertaking regular stock assessments.

6.5 Climate change is impacting on people and ecosystems; in spite of this, healthy ecosystems can help us adapt to climate change

The impacts of climate change are evident across all realms and within most species groups. Biodiversity provides resilience against the worst effects of climate change (*established but incomplete*). Restoring ecosystems and maintaining them in a good ecological condition means they are better able to support natural adaptation and mitigation processes, offering increased protection to human communities and reducing the economic burden of future climate disasters.

7 South Africa's biodiversity is under pressure, but solutions are at hand

7.1 Estuaries and wetlands are the most threatened and least protected ecosystems in South Africa

Estuarine and inland wetland ecosystems face many pressures and are highly threatened (*established but incomplete*). The restoration and protection of estuaries and inland wetlands will secure essential benefits and deliver large return on investment.

7.2 Coastal biodiversity assets, including beaches, are at risk

Sixty per cent of coastal ecosystem types are threatened – a result of the many pressures concentrated on the coast (*well established*). Judicious coastal development that avoids sensitive areas can minimise further damage, maintain ecological infrastructure and reduce climate risks.

7.3 Protected areas: investment success in the ocean and on land

Protected areas have expanded in the ocean and on land, and are a source of pride for South Africans (*well established*). Continued expansion will help to ensure biodiversity conservation, ecological sustainability and even more social and economic benefits from biodiversity.

7.4 Protected areas: providing effective protection for many species

South Africa's protected areas are generally providing good protection for species, as shown by new protection level indicators for species (*established but incomplete*). The results provide important feedback for protected area expansion strategies and for protected area management.

7.5 Freshwater fishes are the most threatened species group in South Africa

Freshwater fishes are the most threatened of all species groups that have been fully assessed in South Africa, and half of South Africa's freshwater fish species are found nowhere else in the world (*established but incomplete*). Effective management and conservation strategies to halt the decline and promote recovery of threatened fish species are needed, focussed on the rivers and catchments where these fish occur.

7.6 Trends in threat status show rapid declines in some of South Africa's species, especially freshwater species and butterflies

Changes in species threat status over time were tracked for eight taxonomic groups using the IUCN Red List Index (RLI). Increased extinction risk is evident for most groups, but freshwater species and butterflies, in particular, show a steep decline (*established but incomplete*). For the RLI to be more comprehensive, repeat assessments are required for species in the marine and estuarine realms, and invertebrates in general.

7.7 Areas where pressures are concentrated should be priorities for spatial planning

The spatial distribution of pressures on biodiversity across the landscape and seascape is uneven. Pressure hotspots, where many different pressures converge, require strategic spatial planning and focussed management (*established but incomplete*).

7.8 Biological invasions threaten biodiversity and human wellbeing

Over 100 alien species have a severe impact on South Africa's biodiversity and, in some cases, on human wellbeing (*well established*). Although some successes in the management of biological invasions have been achieved, the adoption of a national strategy for managing biological invasions, improved project-level planning for prevention and management, and enhanced spatially explicit data will greatly increase effectiveness of current efforts.

7.9 Cooperative governance is essential for healthy landscapes and seascapes

Biodiversity patterns and ecological processes are connected in complex ways that cross realms as well as human-constructed boundaries. At the same time, human activities in a range of different sectors that have separate policies and legislation and are separately managed, can impact on the same biodiversity or ecological infrastructure (*established but incomplete*). To deal with this interconnectedness, cooperative governance and cross-sectoral planning and decision making are essential.

8 The NBA stimulates work to address knowledge gaps

8.1 South Africa's new seamless map of ecosystem types paves the way for improved assessment, planning and monitoring

Substantial progress made in classifying and mapping ecosystem types seamlessly across all realms has unlocked comprehensive and systematic assessment and planning for all of South Africa's territory, providing improved information to inform policy and decision making.

8.2 New indicators developed during the NBA 2018 advance South Africa's ability to report on the status of biodiversity

The collaborative process of undertaking the NBA builds on innovations and advances in the biodiversity sector to produce new techniques and advances in knowledge. In the NBA 2018, several new indicators have been developed to provide a more comprehensive picture of the state of ecosystems and species.

8.3 Evaluation of genetic diversity brings new value to the NBA

Genetic diversity enables species to evolve and adapt within an ever-changing environment. The development of potential genetic diversity indicators for national-level assessments and monitoring has added value to the NBA and South Africa's international reporting commitments.

8.4 Investment in strategic and collaborative biodiversity monitoring programmes is crucial to inform management and decision making and for biodiversity assessments

Investment in existing and future strategic and cooperative biodiversity monitoring programmes is essential to strengthen our ability to detect and report on trends, plan accordingly and manage effectively.

Part III

Integrated findings

9 National Headline Indicators

9.1 Overview

This section brings together the findings from components of the NBA 2018 and presents them in an integrated fashion. The headline indicators are compared across realms and taxonomic groups, key pressures that affect all realms are highlighted, and genetic diversity – that applies to all realms and taxonomic groups – is addressed.

9.1.1 Red List Index of Ecosystems

Trends in ecosystem threat status

9.1.2 Protection Level

add text

9.2 Species indicators

add text

9.2.1 Red List Index of Species

Trends in species threat status

9.2.2 Species Protection Level

add text

9.3 Genetic diversity

9.3.1 Genetic diversity at population level

add text

9.3.2 Genetic diversity at landscape level

add text

10 Pressures

10.1 Synthesis

The International Union for the Conservation of Nature (IUCN) proposes a threats classification scheme with a hierarchical structure for various pressures on biodiversity. The NBA 2018 adopted this approach with some minor language adaptations and the resulting pressure plots are used throughout the NBA 2018 reports. The species plots were based on a meta-analysis of species Red List assessments (which document pressures on each species) and the cross-realm plot was informed by the species meta-analysis and expert inputs.

10.1.1 Climate change

add text

10.1.2 Land cover change

add text

10.1.3 Degradation

add text

10.1.4 Alien invasive species

add text

10.1.5 Pesticides

add

10.2 Ecological condition overview

Ecological condition is estimated using a range of different approaches across the realms, but essentially depends on the ability to spatially represent the various pressures exerted on biodiversity. Ecological condition in the terrestrial realm relies primarily on land cover change data; cumulative pressure mapping is used in the marine realm; and a multi-criteria ecological condition framework is used in the estuarine and inland aquatic realms. The different systems were aligned as far as possible in the NBA to allow for crossrealm comparisons and unified terminology.

The marine and terrestrial realms are similar in terms of their relatively high percentage of natural/near-natural ecosystem extent ($\pm 80\%$). In these extensive realms, ecosystem modification tends to be focussed in pressure hotspots, usually linked to regional characteristics such as high productivity, accessibility and valuable natural resources; while large areas remain relatively unmodified or intact. For example, the Cape lowlands have extensive winter field crops while the mountainous areas of the Cape see far less intensive agriculture; all bay ecosystem types, the shelf edge and the KwaZulu-Natal Bight are subject to multiple pressures while many deep sea ecosystems

10.2.1 Ecological condition per realm

add text

10.2.1.1 Terrestrial realm

add text

10.2.1.2 Freshwater (inland aquatic) realm

add text

10.2.1.3 Estuarine realm

add text

10.2.1.4 Marine realm

add text

10.2.1.5 Coast cross-realm

add text

Part IV

Findings by realm

11 Terrestrial realm

Recommended citation:

11.1 Overview

add text

11.2 Input data and method for the terrestrial realm

add text

11.2.1 Ecosystem assessments

add text

11.2.2 Species assessments

add text

11.3 Key drivers and pressures impacting terrestrial ecosystems

add text

11.3.1 Habitat loss

add text

11.3.2 Land degradation

add text

11.3.3 Biological invasions

add text

11.3.4 Climate change

add text

11.3.5 Biological resource use

add text

11.4 Terrestrial ecosystems

11.4.1 Map of ecosystem types

add text

11.4.2 Overview of South African biomes

add text

11.4.2.1 Albany Thicket

add text

11.4.2.2 Azonal Vegetation

add text

11.4.2.3 Desert

add text

11.4.2.4 Forests

add text

11.4.2.5 Fynbos

add text

11.4.2.6 Grassland

add text

11.4.2.7 Indian Ocean Coastal Belt

add text

11.4.2.8 Nama-Karoo

add text

11.4.2.9 Savanna

add text

11.4.2.10 Succulent Karoo

add text

11.5 Ecological condition

11.6 Overview

add text

11.6.1 Ecological condition per realm

add text

11.6.1.1 Albany Thicket

add text

11.6.1.2 Azonal Vegetation

add text

11.6.1.3 Desert

add text

11.6.1.4 Forests

add text

11.6.1.5 Fynbos

add text

11.6.1.6 Grassland

add text

11.6.1.7 Indian Ocean Coastal Belt

add text

11.6.1.8 Nama-Karoo

add text

11.6.1.9 Savanna

add text

11.6.1.10 Succulent Karoo

add text

11.7 Red List of Ecosystems

add text

11.7.1 Red List of Ecosystems per biome

add text

11.7.1.1 Albany Thicket

add text

11.7.1.2 Azonal Vegetation

add text

11.7.1.3 Desert

add text

11.7.1.4 Forests

add text

11.7.1.5 Fynbos

add text

11.7.1.6 Grassland

add text

11.7.1.7 Indian Ocean Coastal Belt

add text

11.7.1.8 Nama-Karoo

add text

11.7.1.9 Savanna

add text

11.7.1.10 Succulent Karoo

add text

11.7.2 Red List of Ecosystems - Prince Edward Island

add text

11.8 Protection Level

add text

11.8.1 Protection level per biome

add text

11.8.1.1 Albany Thicket

add text

11.8.1.2 Azonal Vegetation

add text

11.8.1.3 Desert

add text

11.8.1.4 Forests

add text

11.8.1.5 Fynbos

add text

11.8.1.6 Grassland

add text

11.8.1.7 Indian Ocean Coastal Belt

add text

11.8.1.8 Nama-Karoo

add text

11.8.1.9 Savanna

add text

11.8.1.10 Succulent Karoo

add text

11.9 Terrestrial species

11.9.1 Utilized terrestrial species resource assessments

add text

12 Freshwater realm

Recommended citation:

12.1 Overview

add text

12.1.1 Rivers

Ecological condition of rivers

12.1.2 Wetlands

Ecological condition of wetlands

12.2 Ecosystems

add text

12.2.1 Rivers

Map ecosystem types - rivers

12.2.2 Wetlands

Map of ecosystem types - wetlands

12.2.3 Riparian

Map of ecosystem types - riparian

12.3 Ecological condition

add text

12.3.1 Rivers

Ecological condition of rivers

12.3.2 Wetlands

Ecological condition of wetlands

12.4 Red List of Ecosystems

add text

12.4.1 Red List of Ecosystems - rivers

add text

12.4.2 Red List of Ecosystems - wetlands

add text

12.5 Protection level

add text

12.5.1 Protection level of rivers

add text

12.5.2 Protection level of wetlands

add text

12.6 Species

add

13 Estuarine realm

Recommended citation:

13.1 Overview

add text

13.2 Ecological condition

add text

13.2.1 Condition

add text

13.2.2 Pressures

Estuary pressure information and condition dataset

Estuarine Invasive Alien Aquatic Plants

13.3 Ecosystems

13.3.1 Habitats and vegetation

add text

13.3.1.1 Functional zone

Estuary Functional Zone update

KZN Estuary Functional Zone

13.3.1.2 Classification updates

DWS Great Fish to Tsitsikamma Classification

13.4 Red List of Ecosystems

add text

13.5 Protection Level

add text

13.6 Species

add text

14 Marine realm

Recommended citation:

14.1 Overview

add text

14.2 Input data and method for the marine realm

add text

14.2.1 Ecosystem assessments

add text

14.2.2 Species assessments

add text

14.3 Key drivers and pressures impacting marine ecosystems

add text

14.3.1 Habitat loss

add text

14.3.2 Degradation

add text

14.3.3 Biological invasions

add text

14.3.4 Climate change

add text

14.3.5 Biological resource use

add text

14.4 Ecosystems

add text

14.4.1 Ecosystems in the marine realm

add text

14.4.2 Ecosystems around Prince Edward Island

add text

14.5 Ecological condition

add text

14.5.1 Marine pressure datasets

add text

14.6 Red List of Ecosystems

add text

14.7 Protection level

add text

14.8 Species

add text

15 Coast cross-cutting realm

Recommended citation:

15.1 Overview

add text

15.2 Ecosystems

add text

15.2.1 Map of ecosystem types

add text

15.3 Ecological condition

add text

15.4 Red List of Ecosystems

add text

15.5 Protection level

add text

15.6 Species

add text

Part V

Species

16 Species

Recommended citation:

16.1 Amphibians

Taxonomic group summary

16.2 Birds

add text

16.2.1 Terrestrial

Taxonomic group summary

16.2.2 Estuarine

Estuary birds

16.3 Fish

add text

16.3.1 Inland aquatic

Taxonomic group summary - Freshwater fish

16.3.2 Estuarine

Estuary fish

16.4 Invertebrates

16.4.1 Overview of invetebtrate groups assessed to date

16.4.2 Bees

Taxonomic group summary page

16.4.3 Branchiopoda

Taxonomic group summary - Large Branchiopoda

16.4.4 Butterflies

Taxonomic group summary

16.4.5 Crabs

Taxonomic group summary

16.4.6 Dragonflies

Taxonomic group summary

16.4.7 Mollusca

16.4.7.1 Terrestrial

Taxonomic group summary

16.4.7.2 Inland aquatic

16.4.8 Spiders

Taxonomic group summary

16.4.9 Estuary invertebrates

Taxonomic group summary

16.5 Mammals

16.5.1 Terrestrial

Taxonomic group summary

16.5.2 Inland aquatic

add text

16.5.3 Marine

add text

16.6 Plants

Taxonomic group summary

16.7 Reptiles

Taxonomic group summary

Part VI

Priority actions and responses

17 Priority actions for improving future NBAs

17.1 Overview

add text

17.2 Knowledge Gaps and Research Priorities for Strengthening the NBA

17.2.1 Review of NBA 2018 Research, Monitoring and Data Management Priorities

add text

17.2.2 NBA 2025 Research, Monitoring and Data Management Priorities

add text

17.2.2.1 Terrestrial realm

add text

17.2.2.2 Freshwater (inland aquatic) realm

add text

17.2.2.3 Estuarine realm

add text

17.2.2.4 Marine realm

add text

17.2.2.5 Coast cross-realm

add text

17.3 Review of tools used to measure genetic diversity

add text

17.3.1 State of DNA barcoding

Genetic resources and tools

17.3.2 Use of eDNA in South Africa

add text

17.3.3 State of SA's Biobanks

add text

18 South Africa's responses for managing and conserving biodiversity

18.1 Overview

add text

18.2 Avoid loss

add text

18.3 Protect

18.3.1 Protection over time indices?

Protected area layer, consolidated annual time steps for PL and Accounts

18.4 Restore

Response initiatives map

Recording where and what restoration is happening in the country

18.5 Resources for biodiversity conservation

add text

18.6 The need for evidence on the effectiveness and impact of responses

add text

References

Knuth, Donald E. 1984. “Literate Programming.” *Comput. J.* 27 (2): 97–111. <https://doi.org/10.1093/comjnl/27.2.97>.