


Lucile Soler PhD
Nima Rafati PhD

Bacterial Genome Annotation

https://nbisweden.github.io/workshop-genome_annotation_elixir/labs/prokaryote_annotation



Genome assembly and annotation course
2021

Bacterial genome characteristics



- A bacterial genome is a single "circular" DNA molecule with several million base pairs in size
- Bacteria can contains plasmids (small and circular DNA molecules, that contain (usually) non-essential genes)
- Genomes contain a few thousand genes.
- "Gene density" is much higher than in humans, one million base pairs of bacterial DNA contains about 500 to 1000 genes.
 - bacterial genes have no introns (intron-less),
 - the average number of codons in bacterial genes is less than in human genes,
 - neighboring genes are very close together throughout the genome

Bacterial feature types




- protein coding genes and associated features
 - promoter (-10, -35)
 - ribosome binding site (RBS)
 - coding sequence (CDS)
 - signal peptide, protein domains, structure
 - terminator
- non coding genes
 - transfer RNA (tRNA)
 - ribosomal RNA (rRNA)
 - non-coding RNA (ncRNA)
- other
 - repeat patterns, origin of replication, ...




Two strategies for identifying coding genes:

- **sequence alignment**
 - find known protein sequences in the contigs
 - transfer the annotation across
 - will miss proteins not in your database
 - may miss partial proteins
- ***ab initio* gene finding**
 - find candidate open reading frames
 - build model of ribosome binding sites
 - predict coding regions
 - may choose the incorrect start codon
 - may miss atypical genes, overpredict small genes

Some good existing tools



Software	<i>ab initio</i>	align- ment	Availability	Speed
RAST	yes	yes	web only	12-24 hours
xBASE	yes	no	web only	>4 hours
BG7	no	yes	standalone	>10 hours
PGAAP (NCBI)	yes	yes	email / we	>1 month



- Fast
 - exploits multi-core computers (aim < 15min)
- Convenient
 - Does structural and functional annotation in one go
 - Help submitting to NCBI and ENA
- Standards compliant
 - GFF3/GBK for viewing, TBL/FSA for Genbank.
- Provenance
 - Keep record of where/how/why it was annotated
- Also annotates archaea, mitochondria, and viruses

- Complicated to install
 - many dependencies (available on conda)

Feature prediction tools used by Prokka :


Tool (reference)	Features predicted
Prodigal (Hyatt 2010)	Coding sequence (CDS)
RNAmmer (Lagesen <i>et al.</i> , 2007)	Ribosomal RNA genes (rRNA)
Aragorn (Laslett and Canback, 2004)	Transfer RNA genes
SignalP (Petersen <i>et al.</i> , 2011)	Signal leader peptides
Infernal (Kolbe and Eddy, 2011)	Non-coding RNA

Seemann T. *Prokka: rapid prokaryotic genome annotation*. **Bioinformatics**. 2014 Jul 15;30(14):2068-9. [PMID:24642063](https://pubmed.ncbi.nlm.nih.gov/24642063/)



- Prodigal identifies the coordinates of candidates genes
- Compares with a database of known sequences
 - Small trustworthy database: the user provides a set of annotation proteins (optional)
 - Genus-specific proteome (optional)
 - Medium-size domain specific database: Uniprot-Swissprot
 - Curated model of protein families: all proteins from finished bacterial genomes in Refseq
 - HMMs profile: Pfam, TIGRFAMS (with HMMER)
 - If nothing is found, is labeled as 'hypothetical protein'

Prokka pipeline (simplified)



- Only one parameter mandatory :
Input fasta format
 - `prokka [options] <contigs.fasta>`
- More than 30 different options available
 - `prokka --help`

Command line options



SciLifeLab

NBS
NATIONAL BIOINFORMATICS
INFRASTRUCTURE SWEDEN

```
General:
  --help          This help
  --version       Print version and exit
  --docs          Show full manual/documentation
  --citation      Print citation for referencing Prokka
  --quiet         No screen output (default OFF)
  --debug         Debug mode: keep all temporary files (default OFF)

Setup:
  --listdb        List all configured databases
  --setupdb       Index all installed databases
  --cleandb       Remove all database indices
  --depends        List all software dependencies

Outputs:
  --outdir [X]    Output folder [auto] (default '')
  --force         Force overwriting existing output folder (default OFF)
  --prefix [X]    Filename output prefix [auto] (default '')
  --addgenes      Add 'gene' features for each 'CDS' feature (default OFF)
  --locustag [X]  Locus tag prefix (default 'PROKKA')
  --increment [N] Locus tag counter increment (default '1')
  --gffver [N]    GFF version (default '3')
  --compliant     Force Genbank/ENA/DDJB compliance: --genes --mincontiglen 200 --centre XXX (default OFF)
  --centre [X]    Sequencing centre ID. (default '')

Organism details:
  --genus [X]     Genus name (default 'Genus')
  --species [X]   Species name (default 'species')
  --strain [X]    Strain name (default 'strain')
  --plasmid [X]   Plasmid name or identifier (default '')

Annotations:
  --kingdom [X]   Annotation mode: Archaea|Bacteria|Mitochondria|Viruses (default 'Bacteria')
  --gcode [N]     Genetic code / Translation table (set if --kingdom is set) (default '0')
  --gram [X]      Gram: -/neg +/pos (default '')
  --usegenus      Use genus-specific BLAST databases (needs --genus) (default OFF)
  --proteins [X]  Fasta file of trusted proteins to first annotate from (default '')
  --hmms [X]      Trusted HMM to first annotate from (default '')
  --metagenome    Improve gene predictions for highly fragmented genomes (default OFF)
  --rawproduct    Do not clean up /product annotation (default OFF)

Computation:
  --fast          Fast mode - skip CDS /product searching (default OFF)
  --cpus [N]      Number of CPUs to use [0=all] (default '8')
  --mincontiglen [N] Minimum contig size [NCBI needs 200] (default '1')
  --evaluate [n.n] Similarity e-value cut-off (default '1e-06')
  --rfam          Enable searching for ncRNAs with Infernal+Rfam (SLOW!) (default '0')
  --norrna        Don't run rRNA search (default OFF)
  --notrna        Don't run tRNA search (default OFF)
  --rnammer       Prefer RNAmmer over Barrnap for rRNA prediction (default OFF)
```

Prokka output

Extension	Description
.gff	This is the master annotation in GFF3 format, containing both sequences and annotations. It can be viewed directly in Artemis or IGV.
.gbk	This is a standard Genbank file derived from the master .gff. If the input to prokka was a multi-FASTA, then this will be a multi-Genbank, with one record for each sequence.
.fna	Nucleotide FASTA file of the input contig sequences.
.faa	Protein FASTA file of the translated CDS sequences.
.ffn	Nucleotide FASTA file of all the prediction transcripts (CDS, rRNA, tRNA, tmRNA, misc_RNA)
.sqn	An ASN1 format "Sequin" file for submission to Genbank. It needs to be edited to set the correct taxonomy, authors, related publication etc.
.fsa	Nucleotide FASTA file of the input contig sequences, used by "tbl2asn" to create the .sqn file. It is mostly the same as the .fna file, but with extra Sequin tags in the sequence description lines.
.tbl	Feature Table file, used by "tbl2asn" to create the .sqn file.
.err	Unacceptable annotations - the NCBI discrepancy report.
.log	Contains all the output that Prokka produced during its run. This is a record of what settings you used, even if the --quiet option was enabled.
.txt	Statistics relating to the annotated features found.
.tsv	Tab-separated file of all features: locus_tag,ftype,gene,EC_number,product

<https://github.com/tseemann/prokka#output-files>

Prokka output



GFF format

```

Chromosome   Prodigal:2.6   CDS      7846    8796    .      +      0      ID=KFDOKKAG_00008;int
8;product=hypothetical protein
Chromosome   Prodigal:2.6   CDS      8812    9714    .      -      0      ID=KFDOKKAG_00009;eC
on:Prodigal:2.6,similar to AA sequence:UniProtKB:067644;locus_tag=KFDOKKAG_00009;product=Ribonuclease
Chromosome   Prodigal:2.6   CDS      9967    10398   .      +      0      ID=KFDOKKAG_00010;int
0;product=hypothetical protein
Chromosome   Prodigal:2.6   CDS      10385   11752   .      -      0      ID=KFDOKKAG_00011;eC
ion:Prodigal:2.6,similar to AA sequence:UniProtKB:P0ACV0;locus_tag=KFDOKKAG_00011;product=Lipid A bid
Chromosome   Prodigal:2.6   CDS      11883   13139   .      -      0      ID=KFDOKKAG_00012;int
2;product=hypothetical protein
Chromosome   Prodigal:2.6   CDS      13136   13828   .      -      0      ID=KFDOKKAG_00013;eC
on:Prodigal:2.6,similar to AA sequence:UniProtKB:Q45589;locus_tag=KFDOKKAG_00013;product=Cyclic di-AM
Chromosome   Prodigal:2.6   CDS      14205   15545   .      +      0      ID=KFDOKKAG_00014;eC
on:Prodigal:2.6,similar to AA sequence:UniProtKB:Q09049;locus_tag=KFDOKKAG_00014;product=Cytochrome b
Chromosome   Prodigal:2.6   CDS      15557   16618   .      +      0      ID=KFDOKKAG_00015;eC
ion:Prodigal:2.6,similar to AA sequence:UniProtKB:P26458;locus_tag=KFDOKKAG_00015;product=Cytochrome
Chromosome   Prodigal:2.6   CDS      16716   18020   .      -      0      ID=KFDOKKAG_00016;int

```

Seqid	source	type	start	end	score	strand	phase	attributes
Chr1	Prodigal	exon	234	1543	.	+	.	gene_id "gene1"; transcript_id "transcript1"; "prediction:.., protein motif..."
Chr1	Snap	CDS	577	1543	.	+	0	gene_id "gene1"; transcript_id "transcript1";

Bacterial Genome Annotation

Exercises


```
Summarized benchmarking in BUSCO notation for file /proj/g2019006/nobackup/lucile/data/raw_computes/Escherichia_coli_k_12.fa
BUSCO was run in mode: genome
```

```
C:98.6%[S:98.6%,D:0.0%],F:0.0%,M:1.4%,n:148
```

```
146    Complete BUSCOs (C)
146    Complete and single-copy BUSCOs (S)
0      Complete and duplicated BUSCOs (D)
0      Fragmented BUSCOs (F)
2      Missing BUSCOs (M)
148    Total BUSCO groups searched
```

```
# Summarized benchmarking in BUSCO notation for file /proj/g2019006/nobackup/lucile/data/raw_computes/Streptococcus.fa
# BUSCO was run in mode: genome
```

```
C:82.4%[S:22.3%,D:60.1%],F:4.7%,M:12.9%,n:148
```

```
122    Complete BUSCOs (C)
33     Complete and single-copy BUSCOs (S)
89     Complete and duplicated BUSCOs (D)
7      Fragmented BUSCOs (F)
19     Missing BUSCOs (M)
148    Total BUSCO groups searched
```

```
Summarized benchmarking in BUSCO notation for file /proj/g2019006/nobackup/lucile/data/raw_computes/Chlamydia_trachomatis_a_363.f
BUSCO was run in mode: genome
```

```
C:75.7%[S:75.7%,D:0.0%],F:4.7%,M:19.6%,n:148
```

```
112    Complete BUSCOs (C)
112    Complete and single-copy BUSCOs (S)
0      Complete and duplicated BUSCOs (D)
7      Fragmented BUSCOs (F)
29     Missing BUSCOs (M)
148    Total BUSCO groups searched
```



```
INFO To reproduce this run: python /opt/miniconda3/bin/busco -i /home/data/byod/Annotation/data/bacterial_annotation/prokka_Esch
erichia/PROKKA_05102019.faa -o Eschi_busco_prot -l /home/data/opt-byod/busco/lineages/bacteria_odb9/ -m proteins -c 8 -sp E_coli_K1
2
INFO Check dependencies...
INFO Check input file...
INFO Temp directory is ./tmp/
INFO Running HMMER on the proteins:
INFO 07/01/2019 10:54:28 => 0% of predictions performed (148 to be done)
INFO 07/01/2019 10:54:29 => 10% of predictions performed (18/148 candidate proteins)
INFO 07/01/2019 10:54:29 => 20% of predictions performed (32/148 candidate proteins)
INFO 07/01/2019 10:54:29 => 30% of predictions performed (49/148 candidate proteins)
INFO 07/01/2019 10:54:29 => 40% of predictions performed (61/148 candidate proteins)
INFO 07/01/2019 10:54:29 => 50% of predictions performed (79/148 candidate proteins)
INFO 07/01/2019 10:54:29 => 60% of predictions performed (91/148 candidate proteins)
INFO 07/01/2019 10:54:29 => 70% of predictions performed (106/148 candidate proteins)
INFO 07/01/2019 10:54:30 => 80% of predictions performed (120/148 candidate proteins)
INFO 07/01/2019 10:54:30 => 90% of predictions performed (135/148 candidate proteins)
INFO 07/01/2019 10:54:30 => 100% of predictions performed
INFO Results:
INFO C:100.0%[S:100.0%,D:0.0%],F:0.0%,M:0.0%,n:148
INFO 148 Complete BUSCOs (C)
INFO 148 Complete and single-copy BUSCOs (S)
INFO 0 Complete and duplicated BUSCOs (D)
INFO 0 Fragmented BUSCOs (F)
INFO 0 Missing BUSCOs (M)
INFO 148 Total BUSCO groups searched
```

```

INFO  To reproduce this run: python /opt/miniconda3/bin/busco -i /home/data/byod/Annotation/data/bacterial_annotation/prokka_Stre
ptococcus/PROKKA_05102019.faa -o strepto_busco_prot -l /home/data/opt-byod/busco/lineages/bacteria_odb9/ -m proteins -c 8 -sp E_coli
_K12
INFO  Check dependencies...
INFO  Check input file...
INFO  Temp directory is ./tmp/
INFO  Running HMMER on the proteins:
INFO  07/01/2019 10:53:29 => 0% of predictions performed (148 to be done)
INFO  07/01/2019 10:53:30 => 10% of predictions performed (17/148 candidate proteins)
INFO  07/01/2019 10:53:30 => 20% of predictions performed (33/148 candidate proteins)
INFO  07/01/2019 10:53:30 => 30% of predictions performed (46/148 candidate proteins)
INFO  07/01/2019 10:53:30 => 40% of predictions performed (61/148 candidate proteins)
INFO  07/01/2019 10:53:30 => 50% of predictions performed (78/148 candidate proteins)
INFO  07/01/2019 10:53:30 => 60% of predictions performed (91/148 candidate proteins)
INFO  07/01/2019 10:53:30 => 70% of predictions performed (106/148 candidate proteins)
INFO  07/01/2019 10:53:31 => 80% of predictions performed (120/148 candidate proteins)
INFO  07/01/2019 10:53:31 => 90% of predictions performed (136/148 candidate proteins)
INFO  07/01/2019 10:53:31 => 100% of predictions performed
INFO  Results:
INFO  C:83.7%[S:18.2%,D:65.5%],F:6.1%,M:10.2%,n:148
INFO  124 Complete BUSCOs (C)
INFO  27 Complete and single-copy BUSCOs (S)
INFO  97 Complete and duplicated BUSCOs (D)
INFO  9 Fragmented BUSCOs (F)
INFO  15 Missing BUSCOs (M)
INFO  148 Total BUSCO groups searched

```

```
INFO To reproduce this run: python /opt/miniconda3/bin/busco -i /home/data/byod/Annotation/data/bacterial_annotation/prokka_Chla
mydia/PROKKA_05102019.faa -o chlamydia_busco_prot -l /home/data/opt-byod/busco/lineages/bacteria_odb9/ -m proteins -c 8 -sp E_coli_
K12
INFO Check dependencies...
INFO Check input file...
INFO Temp directory is ./tmp/
INFO Running HMMER on the proteins:
INFO 07/01/2019 10:50:52 => 0% of predictions performed (148 to be done)
INFO 07/01/2019 10:50:52 => 10% of predictions performed (17/148 candidate proteins)
INFO 07/01/2019 10:50:52 => 20% of predictions performed (32/148 candidate proteins)
INFO 07/01/2019 10:50:52 => 30% of predictions performed (46/148 candidate proteins)
INFO 07/01/2019 10:50:52 => 40% of predictions performed (63/148 candidate proteins)
INFO 07/01/2019 10:50:53 => 50% of predictions performed (77/148 candidate proteins)
INFO 07/01/2019 10:50:53 => 60% of predictions performed (92/148 candidate proteins)
INFO 07/01/2019 10:50:53 => 70% of predictions performed (107/148 candidate proteins)
INFO 07/01/2019 10:50:53 => 80% of predictions performed (120/148 candidate proteins)
INFO 07/01/2019 10:50:53 => 90% of predictions performed (136/148 candidate proteins)
INFO 07/01/2019 10:50:53 => 100% of predictions performed
INFO Results:
INFO C:81.1%[S:81.1%,D:0.0%],F:4.1%,M:14.8%,n:148
INFO 120 Complete BUSCOs (C)
INFO 120 Complete and single-copy BUSCOs (S)
INFO 0 Complete and duplicated BUSCOs (D)
INFO 6 Fragmented BUSCOs (F)
INFO 22 Missing BUSCOs (M)
INFO 148 Total BUSCO groups searched
```