

Creation of Spectral Database and ZPTW Profiles

Eric Nussbaumer ^{*1}

¹*National Center for Atmospheric Research, Boulder, CO, USA*

September 2015

Abstract

This document outlines the creation of the spectral database as well as the profiles for pressure, temperature, and water vapor.

^{*}corresponding author: ebaumer@ucar.edu +1 (303) 497-1861

1 Introduction

The spectral database and ZPTW (altitude, pressure, temperature, and water vapor) profiles are necessary pre-processing steps to retrievals. The spectral database holds information pertaining to each of the measurements. A spectral database is unique to each site. The majority of information in the spectral database comes from the OPUS file itself; however, we append meteorological data from local weather stations. There are several steps in creating the spectral database:

1. Creating the initial spectral database
2. Formatting the house log data files
3. Formatting the external station weather data
4. Appending the initial spectral database with house and external station weather data

Note that not all sites have house or external station weather data.

Station	House Data	External Station Data
MLO	Yes	Yes (CMDL)
TAB	Yes	No
FL0	No	Yes (EOL)

The necessary python files are located in the sfit-processing-environment git repository.

2 Spectral Database

2.1 Initial Spectral Database

The initial spectral database file is created by running `ckopus` on the various raw OPUS files. A python program is created to manage the creation of the initial spectral database file (`mkSpecDB.py`). The program will create a new spectral database file or append an already existing file. Associated with `mkSpecDB.py` is an input file. The input file allows one to specify the starting and ending date to process, the station, and the various directories and files to use. In addition, one can specify additional `ckopus` flags to use in the `ckopus` call. There are logical flags which control the creating of a file which list the folders processed and whether `bnr` files are created. These files are located under the `SpectralDatabase` folder of the git repository.

Program	Description
<code>mkSpecDB.py</code>	Main program to create initial spectral database
<code>specDBInputFile.py</code>	Input file for <code>mkSpecDB.py</code> program

2.2 House Data

House data is data that is recorded by the FTS autonomous system, such as outside temperature, pressure, wind direction, etc. The format of this data has changed for each station over time as the instrument gets modified or upgraded. A python program (`station_house_reader.py`) is created to read the various formats and create a standardized file. There is one file for each year. There are no input files for the `station_house_reader.py` program. The time range, station identifier, and directories are specified directly in the program under the main function. An excel spreadsheet describes the various formats for the house log files for MLO and TAB.

Program	Description
<code>station_house_reader.py</code>	Main program to read house data files
<code>HouseReaderC.py</code>	Supporting program with formats of previous house data files
<code>HouseDataLog.xlsx</code>	Excel file with format of house log files

These programs are located in the ExternalData folder of the git repository.

2.3 External Station Data

There are currently two external station data sources used (EOL for FL0, and CMDL for MLO) only the EOL data needs to be pre-processed. The original format of this data is in netcdf files. The program `read_FL0_EOL_data.py` reads the daily netcdf files and creates a yearly text file. There are no input files for `read_FL0_EOL_data.py` program. The year of interest and directories of data are specified directly in the program under the main function. The program `pullAncillaryData.py` pulls the CMDL and EOL data from each individual ftp site.

Program	Description
<code>pullAncillaryData.py</code>	Program to automatically pull EOL and CMDL data
<code>read_FL0_EOL_data.py</code>	Main program to read EOL and CMDL data

These programs are located in the ExternalData folder of the git repository.

2.4 Append Spectral Database File

The final step is appending the initial spectral database file with the house and external station weather data. A python program was created to accomplish this (`appendSpecDB.py`). The program `appendSpecDB.py` reads in the initial spectral database file. It then searches the house and external station files for weather data at the time of observation, plus a certain number of minutes specified by the user. The mean of the data collected is calculated and a new spectral database file is created. If no data is present missing values are used. Associated with `appendSpecDB.py` is an input file. The input file allows one to specify directories and files, year to process, station, how many minutes to use for averaging, and whether to create a comma separated or pre-specified formatted new spectral database file.

Program	Description
appendSpecDB.py	Program to create the append spectral database file
appndSpecDBInputFile.py	Input file for appendSpecDB.py

Note: A warning message will often appear when running this program originating from the python numpy module. This warning is a result of numpy taking the mean of an empty array. This is handled by the main program.

2.5 Steps for Creating Spectral Database

1. Download external station data using pullAncillaryData.py
2. Create initial database file by running mkSpecDB.py
3. Format house data by running station_house_reader.py
4. Format external station data by running read_FL0_EOL_data.py
5. Create appended spectral database file by running appendSpecDB.py

3 ZPTW Profiles

The pressure, temperature, and water vapor profiles can be created from several outside sources. Temperature and pressure profiles are taken from NCEP nmc data; while currently only water profiles are taken from NCEP I and ERA-Interim re-analysis data. Both NCEP and ERA-Interim data are interpolated with WACCM data to reach 120km vertical height. The profiles are daily averages and they reside in the data directories (/data1/tab,mlo,fl0/).

The following is a table showing the various reference profiles, their sources, along with the associated file names.

Profile Type	Source	File Name
Temperature	NCEP nmc	ZPT.nmc.120
Pressure	NCEP nmc	ZPT.nmc.120
Water Vapor	WACCM	w-120.v1
Water Vapor	NCEP I	w-120.v3
Water Vapor	ERA-Interim	w-120.v4
Water Vapor	Retrieved	w-120.YYYYMMDD.HHMMSS.v99
Water Vapor	Retrieved Daily	w-120.v5

The following table shows the various sources for the data.

Data	Source
WACCM	Local (otserver:/data/Campaign/TAB,MLO,FL0/waccm/
NCEP nmc	ftp://ftp.cpc.ncep.noaa.gov/ndacc/ncp/
NCEP I re-analysis	ftp://ftp.cdc.noaa.gov/Datasets/ncp.reanalysis.dailyavgs/
ERA-Interim re-analysis	/glade/p/rda/data/ds627.0/ei.oper.an.pl/

3.1 Pulling Data

The supporting data is pulled with a program using wget. The program is pullAncillaryData.py and is located at: /data/bin/. This program has been setup in cron tab to pull data everyday. The program pullAncillaryData.py gets the following data: NCEP nmc, NCEP I re-analysis, EOL, and CMDL.

ERA-Interim data must be manually pulled through the server data-access.ucar.edu.

The following table shows the local storage of the ancillary data

Data	Local Storage
WACCM	otserver:/data/Campaign/TAB,MLO,FL0/waccm/
NCEP nmc Height	otserver:/data1/ancillary_data/NCEP_NMC/height/
NCEP nmc Temp	otserver:/data1/ancillary_data/NCEP_NMC/temp/
NCEP I Height	otserver:/data1/ancillary_data/NCEPdata/NCEP_hgt/
NCEP I Shum	otserver:/data1/ancillary_data/NCEPdata/NCEP_Shum/
NCEP I Temp	otserver:/data1/ancillary_data/NCEPdata/NCEP_Temp/
NCEP I Trpp	otserver:/data1/ancillary_data/NCEPdata/NCEP_trpp/
ERA-Interim	otserver:/data1/ancillary_data/ERAdata/
EOL	otserver:/data1/ancillary_data/fl0/eol/
CMDL Hourly	otserver:/data1/ancillary_data/mlo/cmdl/Hourly_Data/
CMDL Minute	otserver:/data1/ancillary_data/mlo/cmdl/Minute_Data/

3.2 Pressure & Temperature Profiles

Pressure and temperature profiles in the ZPT.nmc.120 files come from NCEP nmc data. The NCEP nmc data is vertically interpolated with WACCM data to reach 120km. In the event that the NCEP nmc data is not available for a particular day, the WACCM data is substituted.

The NCEP nmc data must first be formatted. This is done using the program NCEPnmcFormat.py.

After forming the NCEP nmc data one can create the altitude, pressure, and temperature profiles using the program MergPrf.py. This program also creates water profiles from WACCM data (v1).

Program	Description
NCEPnmcFormat.py	Program to format the NCEP nmc data
MergPrf.py	Main program to create ZPT and water files from NCEP data

3.3 NCEP I & ERA Interim Water Profiles

The ERA-Interim daily profiles are calculated from 6 hourly data. Both the 6 hourly and daily data for profiles are created. The ERA-Interim data is housed locally at NCAR in the CISL Research Data Archive. There is a three month lag between the current date and when the data becomes available. The data is hosted on /glade/ and can be accessed through the data-access.ucar.edu server. The data can be found at: /glade/p/r-da/data/ds627.0/ei.oper.an.pl/. The following steps should be used to pre-process the data:

1. Copy over the data from glade
2. Convert GRIB format files to NetCDF files using `cnvrtNC.py`
3. Create water profiles using `ERAWaterPrf.py`

The NCEP I re-analysis data are already daily averages. The grid resolution of NCEP I is less than ERA-interim. In addition ERA-Interim assimilates GPS occultation data. It is preferable to use ERA-Interim over NCEP I. The program to create water profiles from NCEP I data is `NCEPwaterPrf.py`.

Program	Description
<code>cnvrtNC.py</code>	Program to convert ERA-Interim GRIB files to NetCDF files
<code>ERAWaterPrf.py</code>	Program to extract daily averaged water profiles from ERA-Interim
<code>NCEPwaterPrf.py</code>	Program to create daily water profiles from NCEP I

4 Retrieved Water Profiles

For all sites (MLO,TAB, and FL0) water is retrieved when available. This water can be used as a prior for other retrievals. The program `retWaterPrf.py` creates `w-120.YYYYMMDD.HHMMSS.v99` for each retrieval. These files are stored in the data directories (/data1/tab,mlo,fl0/). A daily average of these profiles can be created using the program `retWaterPrfDaily.py`. These daily averages are also stored in the main data directories (/data1/tab,mlo,fl0/).

Program	Description
<code>retWaterPrf.py</code>	Program to create water profiles from water retrieval
<code>retWaterPrfDaily.py</code>	Program to create daily average profiles from water retrievals