

Advances in Generating Locally Relevant Climate Change Information

Climate change is affecting the amount and timing of water reaching our rivers and streams. To continue to manage systems effectively and minimize risks, water managers and planners conduct climate impact assessments, which use computer models to project future changes, their associated risks, and opportunities for adaptation. This requires careful consideration on how to make global-scale climate change information (with spatial resolutions of hundreds of kilometers) appropriate for local-scale impact assessments.


THE CHALLENGE

While climate change is occurring globally, changes are felt locally. However, global climate models do not produce information at locally relevant scales. Therefore, water managers and planners use “downscaling” methods to convert global climate data to decision-relevant spatial scales.

One of the more reliable ways to perform this downscaling is through the use of high-resolution regional climate models; however, such models are too computationally expensive to be run for long time periods. As a result, water managers often rely on simpler, statistical methods to downscale climate data. Such statistical methods however do not include atmospheric dynamics, and, therefore, may not represent local climate changes as well.

FACING THIS CHALLENGE

Scientists and engineers in RAL's Hydrometeorological Applications Program at the National Center for Atmospheric Research are collaborating with the U.S. Army Corps of Engineers, the Bureau of Reclamation, and the University of Bergen to develop a more computationally efficient form of regional climate modeling, the Intermediate Complexity Atmospheric Research model (ICAR). ICAR combines large-scale atmospheric circulation from climate models with Linear Mountain Wave theory, a mathematical solution for determining the effect of mountains on atmospheric winds, to generate wind fields and calculate the motion of air parcels within the atmosphere. Then ICAR performs detailed full-physics calculations relating three-dimensional water vapor and temperature to the formation of clouds, rain, and snow. The resulting precipitation predictions are similar to those predicted by a full-complexity regional climate model (WRF), and to the PRISM observed dataset (illustrated in the figure). The source code for ICAR, along with a growing library of documentation, is freely available at <https://www.github.com/NCAR/icar>.



Precipitation totals [mm] simulated by (a) WRF, (b) ICAR, and estimated by (c) PRISM for the winter of 2000-2001.

SCIENTIFIC ADVANCES

ICAR provides a new way of generating local climate information that captures local atmospheric dynamics *and* is computationally efficient, freeing water managers and planners from needing to choose between statistical and dynamical downscaling methods.

- *Physical processes.* ICAR provides a method of downscaling that is derived from physical principles, and incorporates some of the most advanced cloud and precipitation calculations in the scientific literature.
- *Computational efficiency.* ICAR simulations can be performed 100 to 1000 times faster than a comparable simulation with a traditional regional climate model.
- *Skill in current climate.* ICAR's representation of the precipitation that forms mountain snowpack, which is of great importance to those managing water resources in the western U.S., is excellent. ICAR explains 85-95% of the variability represented in a much more complex regional climate model, and is highly correlated with observed precipitation.
- *Flexibility.* ICAR has been designed to use a variety of model inputs and can easily be used to evaluate the influence of different physical assumptions on the downscaling process.

MOVING FORWARD

- Current work is focused on improving the skill and physical process representation of ICAR, particularly for convective precipitation and land surface feedbacks.
- ICAR is being used to generate downscaled climate datasets for the United States, including Alaska and Hawaii.
- ICAR is being evaluated in a variety of environments, including tests in the Alps, Norway, the Himalayas, and Chile.

MORE INFORMATION

Barstad, I., and S. Gronas, 2006: Dynamical structures for southwesterly airflow over southern Norway: the role of dissipation, *Tellus Series a-Dynamic Meteorology and Oceanography*, **58**(1), 2–18, doi:10.1111/j.1600-0870.2006.00152.x

Gutmann, E., I. Barstad, M.P. Clark, J. Arnold, and R. Rasmussen, 2016: The Intermediate Complexity Atmospheric Research Model, *Journal of Hydrometeorology*, **17**(3), 957–973, doi: 10.1175/JHM-D-15-0155.1

Gutmann, E., T. Pruitt, M.P. Clark, L. Brekke, J.R. Arnold, D.A. Raff, and R.M. Rasmussen, 2014: An intercomparison of statistical downscaling methods used for water resource assessments in the United States, *Water Resources Research*, **50**(9), 7167–7186, doi:10.1002/2014WR015559

Gutmann, E.D., R.M. Rasmussen, C. Liu, K. Ikeda, D.J. Gochis, M.P. Clark, J. Dudhia, and G. Thompson, 2012: A Comparison of Statistical and Dynamical Downscaling of Winter Precipitation over Complex Terrain, *Journal of Climate*, **25**(1), 262–281, doi:10.1175/2011JCLI4109.1

PROJECT TEAM AND WEBSITE

NCAR: Ethan Gutmann (PI), Martyn Clark, Trude Eidhammer, Roy Rasmussen

Uni Research (Bergen, Norway): Idar Bartstad

U.S. Army Corps of Engineers: Jeff Arnold

Bureau of Reclamation: Levi Brekke

Contact: Ethan Gutmann - gutmann@ucar.edu | Julie Vano - jvano@ucar.edu

Website: <https://ral.ucar.edu/hap/computational-hydrology>