BROWSER-BASED HANDS-ON

Try EVB with the Kemp reaction


Follow one reaction. See how EVB connects the starting and ending states. Then try changing two model settings yourself.

No installation is needed for the saved examples. Use the links on the workshop website to open each exercise.

Workshop website: nd7996.github.io/Kemp-EVB-Workshop

What changes?

The substrate is the molecule the enzyme acts on. Asp162, part of the enzyme, accepts a proton (a hydrogen nucleus) from it. As the C–H bond breaks, the small ring opens and the product forms.



2 · BEFORE AND AFTER

Two descriptions of the same atoms

REACTANT STATE

Before

The substrate holds the proton and its ring is closed. Asp162 is ready to accept it.

PRODUCT STATE

After

Asp162 holds the proton. The substrate ring is open.

EVB uses both descriptions to model the bond changes.

What do the settings mean?

TWO STATES

Reactant and product

Each has its own bond and charge description for the same atoms.

A

Energy offset

Moves the product-state energy up or down relative to the reactant.

H12

Coupling

Controls how strongly the reactant and product descriptions mix.

Neither α nor H12 is the reaction barrier. Both affect the profile, so real calculations need evidence to choose them.

Prepare, relax, then sample

PREPARE

Build R and P

Use the same atoms and atom order. Assign each state's bonds, charges, force-field terms, and EVB settings. Check mapping, total charge, proton position, solvent, and ions.

RELAX

Settle both endpoints

Minimize and equilibrate R and P separately. Check geometry and stability. This prepares the endpoints; it does not calculate the reaction barrier.

FEP / PROFILE

Sample lambda windows

Sample the path between R and P. Check window overlap and convergence, compare replicas, then analyze the profile, barrier, and reaction free energy.

This workshop includes saved Kemp data for learning the analysis, not the full live scripts for these three simulation stages.

4 · YOUR TURN

Try changing the settings

Open the small EVB example. Move one control at a time. Watch how the curve changes when you change the energy offset, then when you change the coupling.

Open the EVB example

Take about 5 minutes. The small example teaches the controls; it is not the full enzyme calculation.

5 · READ A PROFILE

What do the axes show?

ACROSS

Energy gap

Compares the reactant- and product-state energies. It is not distance or time.

UP

Free energy

Shown relative to the reactant. The rise estimates the barrier; the product level gives the reaction free energy.

A peak in a binned profile is not, by itself, a picture of the molecular transition state.

6 · COMPARE SETTINGS

One barrier does not choose one model

Try the parameter comparison. Different offset/coupling pairs can give similar peak heights while giving different product-versus-reactant energies.

Open the settings comparison

These are teaching examples made from saved data. Matching a target is not a parameter fit or a validated prediction.

END · THE POINT OF THE EXERCISE

What should you take away?

- EVB connects reactant and product bonding descriptions.
- The offset and coupling shape the profile in different ways.
- Reference data and adequate sampling are needed before interpreting a profile physically.

The short OpenMM example is for learning how to read a profile. It does not reproduce the paper’s longer Q6 calculation.

Listov et al., Nature (2025). DOI: 10.1038/s41586-025-09136-2