Atlantic cod (*Gadus morhua*)

Summary Scores

- **Vulnerability:** Moderate - High
- **Climate Exposure:** Moderate
- **Biological Sensitivity:** High - Very High

Sensitivity

Sensitivity Attributes	Range Score	Range Data Quality	Range Tally	Eastern Gulf of Maine Score	Eastern Gulf of Maine Data Quality	Eastern Gulf of Maine Tally	Western Gulf of Maine Score	Western Gulf of Maine Data Quality	Western Gulf of Maine Tally	Georges Bank Score	Georges Bank Data Quality	Georges Bank Tally	Southern New England Score	Southern New England Data Quality	Southern New England Tally
Adult Mobility	1.91	2		1.8	2		1.84	2		1.92	2		2.08	2	
Complexity in Reproductive Strategy	3.46	2.5		3.48	2.5		3.48	2.5		3.32	2.5		3.56	2.5	
Dispersal of Early Life Stages	2.12	2.22		2.16	2.2		2.16	2.2		2.04	2.3		2.12	2.2	
Egg and Larval Survival and Settlement Requirements	3.41	2.4		3.36	2.4		3.44	2.4		3.28	2.4		3.56	2.4	
Historical Variability in Temperature	3.14	2.6		3.08	2.6		3.2	2.6		3.16	2.6		3.12	2.6	
Juvenile Habitat Requirements	2.42	2.7		2.44	2.7		2.44	2.7		2.2	2.7		2.6	2.7	
Other Stressors	2.37	2.2		2.2	2.2		2.44	2.2		2.24	2.2		2.6	2.2	
Population Growth Rate	2.7	1.85		2.76	1.9		2.84	1.9		2.48	1.8		2.72	1.8	
Prey Specificity	2.22	2.2		2.24	2.2		2.2	2.2		2.2	2.2		2.24	2.2	
Relationship to Calcified Organisms (due to impacts from Ocean Acidification)	1.63	2.15		1.6	2.3		1.64	2.1		1.6	2.1		1.68	2.1	
Spawning Cycle	3.2	2.5		3.24	2.5		3	2.5		3.24	2.5		3.32	2.5	
Stock Size/Status	3.93	2.6		3.96	2.6		3.92	2.6		3.92	2.6		3.92	2.6	
Total Sensitivity | Certainty	**High (3) | 0.87**			**High (3) | 0.82**			**High (3) | 0.66**			**High (3) | 0.92**			**Very High (4) | 0.64**		

Exposure

Exposure Factors	Range Mean	Range SD	Range Tally	Eastern Gulf of Maine Mean	Eastern Gulf of Maine SD	Eastern Gulf of Maine Tally	Western Gulf of Maine Mean	Western Gulf of Maine SD	Western Gulf of Maine Tally	Georges Bank Mean	Georges Bank SD	Georges Bank Tally	Southern New England Mean	Southern New England SD	Southern New England Tally
Bottom Temperature	*2.37*	*1.10*		*2.01*	*0.99*		*2.89*	*0.94*		*1.93*	*1.02*		*3.2*	*0.88*	
Bottom Oxygen	2.8	1.01		2.79	0.74		3.46	0.77		2.21	0.95		3.07	0.96	
Bottom Salinity	*2.58*	*1.04*		*2.42*	*0.83*		*1.82*	*0.88*		*3.34*	*0.91*		*2.82*	*0.44*	
Bottom Aragonite Solubility	1.22	0.61		1.03	0.11		1.02	0.11		1.52	0.85		1	0.00	
Small zooplankton integrated biomass	1.27	0.28		1.3	0.26		1.31	0.25		1.1	0.14		1.52	0.38	
Medium zooplankton integrated biomass	1.35	0.28		1.28	0.19		1.47	0.29		1.25	0.22		1.43	0.40	
Large zooplankton integrated biomass	*1.3*	*0.36*		*1.36*	*0.43*		*1.29*	*0.31*		*1.26*	*0.33*		*1.09*	*0.15*	
Total Exposure	**Moderate (1.97)**	**0.93**		**Moderate (1.67)**	**0.83**		**Moderate (2.25)**	**0.89**		**Moderate (1.76)**	**0.94**		**Moderate (2.45)**	**0.77**	
Total Exposure - Important Variables	***Low (1.29)***	***0.63***		***Low (1.01)***	***0.10***		***Low (1.22)***	***0.54***		***Low (1.42)***	***0.77***		***Moderate (1.87)***	***0.72***	

Vulnerability

Overall Vulnerability Rank	Range Mean	Range SD	Range Tally	Eastern Gulf of Maine Mean	Eastern Gulf of Maine SD	Eastern Gulf of Maine Tally	Western Gulf of Maine Mean	Western Gulf of Maine SD	Western Gulf of Maine Tally	Georges Bank Mean	Georges Bank SD	Georges Bank Tally	Southern New England Mean	Southern New England SD	Southern New England Tally
Total Vulnerability	**Moderate (6.17)**	**3.09**		**Moderate (5.01)**	**2.50**		**Moderate (6.75)**	**2.68**		**Moderate (5.28)**	**2.82**		**High (9.8)**	**3.09**	
Total Vulnerability - Important Variables	***Moderate (4.09)***	***2.23***		***Low (3.02)***	***0.31***		***Moderate (3.65)***	***1.61***		***Moderate (4.25)***	***2.30***		***High (7.48)***	***2.88***	

Low Moderate High Very High

Figure 1: Summary Table of Sensitivity, Exposure, and Vulnerability Scores

Changes from Previous Assessment(s)

Table 1: Changes to Vulnerability

Metric	Current Assessment	Previous Assessment(s)	Change
Vulnerability	Moderate - High	Moderate	None - Increase
Exposure	Moderate	High	Decrease
Sensitivity	High – Very High	Moderate	Increase
Directional Effect	Negative	Negative	None

Stocks




Figure 2: Atlantic cod stocks on the Northeast Shelf.

Atlantic cod has four stocks on the Northeast Shelf: Eastern Gulf of Maine (EGOM), Western Gulf of Maine (WGOM), Georges Bank (GBK), and Southern New England (SNE).

Score Descriptions:

Vulnerability:




Figure 3: Atlantic cod Vulnerability

Mean vulnerability scores within the **Eastern Gulf of Maine, Western Gulf of Maine, and Georges Bank** stocks were **Moderate**, ranging from 5.01 to 6.75 (+/- 2.50 - 2.82). Mean scores in the **Southern New England** stock were **High** (9.8 +/- 3.09). This was driven by the higher Sensitivity score within the stock.

Very High Vulnerability scores were found south of Cape Cod, due to high exposure and high sensitivity within the Southern New England stocks. High Vulnerability scores were found along the coast of the Gulf of Maine and over Georges Bank due to high exposure, whereas exposure was Low within the Gulf of Maine basin.

If Exposure scores based only on factors important to the species distribution model were used, Vulnerability scores dropped to Low in the Eastern Gulf of Maine stock. The results in the remaining stocks did

not change. In this case, Vulnerability was still high south of Cape Cod, but decreased throughout the Gulf of Maine, only remaining high along Georges Bank.

Vulnerability scores in the previous Assessment were Moderate for Atlantic cod, similar to our results (Hare et al. 2016).

Climate Exposure:




Figure 4: Atlantic cod Exposure

Mean (+/- standard deviation) exposure scores within all four cod stocks were **Moderate** when all exposure factors were considered.Across the species range, exposure scores were greatest along the coastline, southeast of Cape Cod, and over Georges Bank. These patterns were driven by high exposure to changes in bottom temperature, salinity, and oxygen. High bottom temperature exposure south of Cape Cod and Long Island drove the higher scores observed in the Western Gulf of Maine and Southern New England Stocks.

In the previous assessment, cod received a High exposure score, driven by high exposure to increased sea surface temperatures and ocean acidification (Hare et al. 2016).




Figure 5: Exposure Factor Maps for Atlantic cod

Monthly Variability




Figure 6: Atlantic cod Exposure and Monthly Variability

While total exposure was relatively consistent across months for cod across their entire range, there was variability throughout the year within the different stocks. Georges Bank, Western Gulf of Maine, and Southern New England stocks had the highest variability. Exposure within the Georges Bank and Western Gulf of Maine stocks ranged from Low to High, whereas exposure ranged from Moderate to Very High in Southern New England. Exposure was generally highest in the summer months, due to high bottom temperature, oxygen and zooplankton biomass exposures.




Figure 7: Atlantic cod Exposure Factor Monthly Variability

Exposure from Significant Variables




Figure 8: Atlantic cod Exposure and Monthly Variability

Bottom temperature, bottom salinity, and large zooplankton biomass were the most important dynamic variables in the Atlantic Cod species distribution model. Cod are piscivores, feeding on other small fishes that likely prey on zooplankton (Bowman 1975). Spawning is closely linked to temperature (Lough 2004).When only important factors were used to calculate exposure, exposure in the Eastern and Western Gulf of Maine stocks, as well as the Georges Bank stock, decreased to Low. The mean exposure score only remained Moderate in the Southern New England stock.

Biological Sensitivity:




Figure 9: Atlantic cod Sensitivity

Of the four cod stocks, three (**Eastern Gulf of Maine, Western Gulf of Maine, and Georges Bank**) received **High** sensitivity scores. The **Southern New England** stock received a **Very High** score. Certainty was highest for the Georges Bank (92%) stock, followed by the Eastern Gulf of Maine (82%), Western Gulf of Maine (66%), and Southern New England (64%).

Stock size, complexity in reproductive strategy, and egg and larval survival and settlement requirements drove the scores across all four stocks. Spawning cycle and historical variability in temperature also scored above 3 for all stocks.

Cod are currently overfished in the Gulf of Maine (Northeast Fisheries Science Center (NEFSC) 2021). Spawning is coupled with copepod production (Brander and Hurley 1992) and may be impacted by warming temperatures (Behan et al. 2023; Drinkwater 2005). Warming temperatures may also reduce adult fitness, and thus reduce egg quality (Klein, Smith, and Kritzer 2016). Recruitment success is linked to retentive circulation patterns in the Gulf of Maine and on Georges Bank (Cong, Sheng, and Thompson

1996; Lough and Bolz 1989), which can be altered by large scale climate patterns and overall regional warming (Pershing et al. 2015).

Atlantic cod sensitivity has increased since the last assessment. Atlantic cod were previously scored as Moderate (Hare et al. 2016). Stock size/status and egg and larval survival and settlement requirements (formally early life history requirements) drove the scores in both assessments.

Data Quality Metrics

Life History Data Quality:

82% of the data quality scores were greater than or equal to 2 for the Georges Bank, Western Gulf of Maine, and Southern New England stocks. This value was 83% for the Eastern Gulf of Maine stock, indicating strong data quality. For all stocks, only Population Growth Rate was considered a data gap (mean attribute data quality score < 2).

Model Confidence:

Experts had moderate confidence in the Atlantic cod model (mean confidence = 2.1 +/- 0.32). Experts noted that cod are often associated with rocky bottoms, which are undersampled by traditional trawl surveys used to build the models. The ensemble model AUC was 0.84, indicating good performance.

Additional Metrics

Distribution Shifts:




Figure 10: Atlantic cod distribution shifts on the Northeast Shelf.

Note: For an animated version of this plot, please see the online version of this document.

While cod populations have declined over the past 30 years, distributions have not shifted significantly. The weighted centroid of the distribution shifted 3.8 km ESE between 1993-1997 and 2019-2023. Significant declines in latitudinal range led to significant losses in core and total habitat area. This was driven by the southern edge of the species' range moving further north.

In the next ten years, the distribution centroid is expected to shift a further 7.3 km ENE in comparison to its current location. Core habitat and total habitat areas are expected to decline, in comparison to 2019-2023 averages, but models predict a significant increase in total habitat area from 2025-2035. This is likely driven by the predicted cooling of bottom temperatures along the Northeast Shelf.

Table 2: Distribution Shifts

Time Period	Centroid Shift (km & bearing)	Core Habitat Area (change) (km2)	Habitat Total Area (change) (km2)	Latitude Range (change) (deg)	Longitude Range (change) (deg)
1993-1997	NA	65,441	159,006	3.43	4.97
2019-2023	3.8 km ESE	60,182 (-5,258)	154,872 (-4,133)	3.3 (-0.12)	4.97 (0)
2031-2035	7.3 km ENE	57,770 (-2,412)	153,955 (-917)	3.3 (0)	4.97 (0)

Directional Effect:

All four stocks of Atlantic Cod will likely be negatively impacted by future climate change (100% certainty in expert scores). This is similar to the previous Assessment (Hare et al. 2016). Continued warming will likely drive declines in recruitment and shifts in spawning phenology.

Climate Effects on Abundance and Distribution

Recent warming in the Gulf of Maine relating to a northward shift in the Gulf Stream and changes in the Atlantic Multidecadal Oscillation and Pacific Decadal Oscillation led to reduced Atlantic cod recruitment and increased mortality (Pershing et al. 2015). Rising temperatures may affect Atlantic cod spawning locations and timing (Behan et al. 2023). Drinkwater (2005) suggested Atlantic cod spawning grounds will experience a shift northward, with spawning timing occurring later in the season as a response to warming ocean temperatures. Climate change may also increase in natural mortality in adult Atlantic cod due to declines in fish condition (Behan et al. 2023). Drinkwater (2005) predicted large-scale declines in Atlantic cod in the Gulf of Maine if bottom temperatures increase above 12°C, either through changes in distribution to colder waters or natural mortality. Reduced fitness in Atlantic cod adults from climate impacts may result in lower quality eggs (Klein, Smith, and Kritzer 2016); however, there may be some capacity to adapt to slower rates of warming (Langan et al. 2020).

Life History Synopsis

Cod in the Northwest Atlantic range from Greenland to Cape Hatteras, NC, with a majority of biomass in the Gulf of Maine and on Georges Bank (Behan et al. 2023). Spawning timing varies by location - later in the year with increasing latitude - although peaks generally occur in the winter and spring (Lough 2004) in water temperatures between 5-7°C. In spring, Atlantic cod spawning matched the variation in peak abundance of *Calanus finmarchicus*. In fall, cod spawned in some, but not all, areas where *Calanus* were abundant. This is consistent with the hypothesis that cod spawning is coupled to copepod

production (Brander and Hurley 1992). Atlantic cod eggs are pelagic and hatch after 8-60 days, varying with temperature (Hardy 1978). Recruitment success increases when dispersion of larvae by currents is reduced (Cong, Sheng, and Thompson 1996). On Georges Bank, a clockwise gyre circulation pattern tends to retain and isolate eggs and larvae (Lough and Bolz 1989). Transformation from larvae to the juvenile stage occurs at sizes greater than 20 mm, where they then move to the bottom (Lough 2004). Early juveniles are dependent on gravel substrates or grass beds in nearshore nurseries to favor survival through predator avoidance and increased food availability (Lough and Bolz 1989). In the Gulf of Maine, juveniles prey on mollusks and other invertebrates, while adults consume a greater proportion of fish including herring, redfish, mackerel, and other cod (Bowman 1975). Adult cod have few predators, other than large sharks and seals (Neuenhoff et al. 2019). Recent modeling studies illustrate growing populations of grey seals, harp seals, and dogfish are a major barrier to cod population recovery (Behan et al. 2023). Adults in the Newfoundland stock migrate, leaving overwintering areas in deep oceanic waters and follow tongues of deep, relatively warm, oceanic waters ("highways") across the shelf to summer feeding areas nearshore. They then move northward along the Newfoundland coast in late summer, and eventually return to wintering areas (Lough 2004). Previous research suggested that in the middle part of their range, Atlantic cod only undertake minor seasonal movements in reaction to changing temperatures. However, more recent research suggests that there are inshore/offshore migrations that correspond to spawning and feeding activity in the Gulf of Maine (Zemeckis et al. 2017). As of 2025, Atlantic cod are divided into 4 stock units (Western & Eastern Gulf Maine, Georges Bank, and Southern New England) based on spawning locations, larval transport pathways, and genetic markers (New England Fishery Management Council (NEFMC) 2025). All stocks are managed under the New England Fishery Management Council's Northeast Multispecies Fishery Management Plan (Klein-MacPhee 2012). The stock is overfished and overfishing is occurring in the Gulf of Maine (Northeast Fisheries Science Center (NEFSC) 2021).

References

Behan, J., A. Tyrell, A. Hart, S. Large, A. Hansell, K. Lankowicz, R. Morse, I. Andruschchenko, J. Gross, and L. Kerr. 2023. "Profile of Ecosystem and Climate Influences on Atlantic Cod." NOAA. https://doi.org/10.25923/c267-zd92.

Bowman, R. E. 1975. "Food Habits of Atlantic Cod, Haddock, and Silver Hake in the Northwest Atlantic, 1969-1972." Lab. Ref. 75-1. U.S. Natl. Mar. Fish. Serv., Northeast Fish. Cent.

Brander, K., and P. C. F. Hurley. 1992. "Distribution of Early-Stage Atlantic Cod (Gadus Morhua), Haddock (Melanogrammus Aeglefinus), and Witch Flounder (Glyptocephalus Cynoglossus) Eggs on the Scotian Shelf: A Reappraisal of Evidence on the Coupling of Cod Spawning and Plankton Production." *Canadian Journal of Fisheries and Aquatic Sciences* 49 (2): 238–51.

Cong, L., J. Sheng, and K. T. Thompson. 1996. "A Retrospective Study of Particle Retention on the Outer Banks of the Scotian Shelf 1956-1993." 170. Canadian Technical Report of Hydrography; Ocean Sciences.

Drinkwater, K. F. 2005. "The Response of Atlantic Cod (Gadus Morhua) to Future Climate Change." *ICES Journal of Marine Science* 62 (7): 1327–37. https://doi.org/10.1016/j.icesjms.2005.05.015.

Hardy, J. D., Jr. 1978. "Development of Fishes of the Mid-Atlantic Bight: An Atlas of Egg, Larval and Juvenile Stages. Vol. 2 Anguillidae Through Syngnathidae." Report FWS/OBS-78/12. U.S. Fish; Wildlife Service, Biological Services Program.

Hare, Jonathan A, Wendy E Morrison, Mark W Nelson, Megan M Stachura, Eric J Teeters, Roger B Griffis, Michael A Alexander, et al. 2016. "A Vulnerability Assessment of Fish and Invertebrates to Climate Change on the Northeast US Continental Shelf." *PloS One* 11 (2): e0146756.

Klein, E. S., S. L. Smith, and J. P. Kritzer. 2016. "Effects of Climate Change on Four New England Groundfish Species." *Reviews in Fish Biology and Fisheries* 27: 317–38. https://doi.org/10.1007/s11160-016-9444-z.

Klein-MacPhee, G. 2012. "Atlantic Cod/Gadus Morhua Linnaeus 1758." In *Fishes of the Gulf of Maine*, edited by B. B. Collette and G. Klein-MacPhee, 3rd ed., 228–35. Washington D.C.: Smithsonian Institution Press.

Langan, J. A., M. C. McManus, D. R. Zemeckis, and J. S. Colli. 2020. "Abundance and Distribution of Atlantic Cod (Gadus Morhua) in a Warming Southern New England." *Fisheries Bulletin* 118 (2): 145–56. https://doi.org/10.7755/FB.118.2.4.

Lough, R. G. 2004. "Essential Fish Habitat Source Document: Atlantic Cod, Gadus Morhua, Life History and Habitat Characteristics." NOAA Technical Memorandum NMFS-NE-190.

Lough, R. G., and G. R. Bolz. 1989. "The Movement of Cod and Haddock Larvae onto the Shoals of Georges Bank." *Journal of Fish Biology* 35 (Suppl. A): 71–79.

Neuenhoff, R. D., D. P. Swain, S. P. Cox, M. K. McAllister, A. W. Trites, C. J. Walters, and M. O. Hammill. 2019. "Continued Decline of a Collapsed Population of Atlantic Cod (Gadus Morhua) Due to Predation-Driven Allee Effects." *Canadian Journal of Fisheries and Aquatic Sciences* 76 (1): 168–84. https://doi.org/10.1139/cjfas-2017-0190.

New England Fishery Management Council (NEFMC). 2025. "Northeast Multispecies Fishery Management Plan Amendment 25." https://d23h0vhsm26o6d.cloudfront.net/250305_Groundfish_Amendment-25_final_submission.pdf.

Northeast Fisheries Science Center (NEFSC). 2021. "Gulf of Maine Atlantic Cod 2021 Update Assessment Report." https://apps-st.fisheries.noaa.gov/sis/docServlet?fileAction=download&fileId=7550.

Pershing, A. J., M. A. Alexander, C. M. Hernandez, L. A. Kerr, A. Le Bris, K. E. Mills, J. A. Nye, et al. 2015. "Slow Adaptation in the Face of Rapid Warming Leads to Collapse of the Gulf of Maine Cod Fishery." *Science* 350: 809–12. https://doi.org/10.1126/science.aac9819.

Zemeckis, D. R., C. Liu, G. W. Cowles, M. J. Dean, W. S. Hoffman, D. Martins, and S. X. Cadrin. 2017. "Seasonal Movements and Connectivity of an Atlantic Cod (Gadus Morhua) Spawning Component in the Western Gulf of Maine." *ICES Journal of Marine Science* 74: 1780–96.