## Sensitivity

| Sensitivity Attributes | Range | | | Eastern Gulf of Maine | | | Western Gulf of Maine | | | Georges Bank | | | Southern New England | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | Score | LH DQ | Tally | Score | LH DQ | Tally | Score | LH DQ | Tally | Score | LH DQ | Tally | Score | LH DQ | Tally |
| Adult Mobility | 1.91 | 2 | | 1.8 | 2 | | 1.84 | 2 | | 1.92 | 2 | | 2.08 | 2 | |
| Complexity in Reproductive Strategy | 3.46 | 2.5 | | 3.48 | 2.5 | | 3.48 | 2.5 | | 3.32 | 2.5 | | 3.56 | 2.5 | |
| Dispersal of Early Life Stages | 2.12 | 2.22 | | 2.16 | 2.2 | | 2.16 | 2.2 | | 2.04 | 2.3 | | 2.12 | 2.2 | |
| Egg and Larval Survival and Settlement Requirements | 3.41 | 2.4 | | 3.36 | 2.4 | | 3.44 | 2.4 | | 3.28 | 2.4 | | 3.56 | 2.4 | |
| Historical Variability in Temperature | 3.14 | 2.6 | | 3.08 | 2.6 | | 3.2 | 2.6 | | 3.16 | 2.6 | | 3.12 | 2.6 | |
| Juvenile Habitat Requirements | 2.42 | 2.7 | | 2.44 | 2.7 | | 2.44 | 2.7 | | 2.2 | 2.7 | | 2.6 | 2.7 | |
| Other Stressors | 2.37 | 2.2 | | 2.2 | 2.2 | | 2.44 | 2.2 | | 2.24 | 2.2 | | 2.6 | 2.2 | |
| Population Growth Rate | 2.7 | 1.85 | | 2.76 | 1.9 | | 2.84 | 1.9 | | 2.48 | 1.8 | | 2.72 | 1.8 | |
| Prey Specificity | 2.22 | 2.2 | | 2.24 | 2.2 | | 2.2 | 2.2 | | 2.2 | 2.2 | | 2.24 | 2.2 | |
| Relationship to Calcified Organisms (due to impacts from Ocean Acidification) | 1.63 | 2.15 | | 1.6 | 2.3 | | 1.64 | 2.1 | | 1.6 | 2.1 | | 1.68 | 2.1 | |
| Spawning Cycle | 3.2 | 2.5 | | 3.24 | 2.5 | | 3 | 2.5 | | 3.24 | 2.5 | | 3.32 | 2.5 | |
| Stock Size/Status | 3.93 | 2.6 | | 3.96 | 2.6 | | 3.92 | 2.6 | | 3.92 | 2.6 | | 3.92 | 2.6 | |
| **Total Sensitivity \| Certainty** | **High (3) \| 0.87** | | | **High (3) \| 0.82** | | | **High (3) \| 0.66** | | | **High (3) \| 0.92** | | | **Very High (4) \| 0.64** | | |

## Exposure

| Exposure Factors | Range | | | Eastern Gulf of Maine | | | Western Gulf of Maine | | | Georges Bank | | | Southern New England | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | Mean | SD | Tally | Mean | SD | Tally | Mean | SD | Tally | Mean | SD | Tally | Mean | SD | Tally |
| Bottom Temperature | 2.37 | 1.10 | | 2.01 | 0.99 | | 2.89 | 0.94 | | 1.93 | 1.02 | | 3.2 | 0.88 | |
| Bottom Salinity | 2.58 | 1.04 | | 2.42 | 0.83 | | 1.82 | 0.88 | | 3.34 | 0.91 | | 2.82 | 0.44 | |
| Large zooplankton integrated biomass | 1.3 | 0.36 | | 1.36 | 0.43 | | 1.29 | 0.31 | | 1.26 | 0.33 | | 1.09 | 0.15 | |
| **Total Exposure** | **Low (1.29)** | **0.63** | | **Low (1.01)** | **0.10** | | **Low (1.22)** | **0.54** | | **Low (1.42)** | **0.77** | | **Moderate (1.87)** | **0.72** | |

## Vulnerability

| Vulnerability | Range | | | Eastern Gulf of Maine | | | Western Gulf of Maine | | | Georges Bank | | | Southern New England | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | Mean | SD | Tally | Mean | SD | Tally | Mean | SD | Tally | Mean | SD | Tally | Mean | SD | Tally |
| **Total Vulnerability** | **Moderate (4.09)** | **2.23** | | **Low (3.02)** | **0.31** | | **Moderate (3.65)** | **1.61** | | **Moderate (4.25)** | **2.3** | | **High (7.48)** | **2.88** | |

