


American Plaice Indicators for Fall 2026 Management Track

American Plaice


Distribution Map




Socioeconomic Indicators

Average price per pound, total landings (lbs), commercial revenue, and n vessels are all below the long term mean, declining since 2020. However, landings and revenue have rebounded slightly. Average revenue per vessel, which had been declining since the mid-2010s to a low in 2020, has risen well above the long-term mean after 2022.


Average price per pound




Total commercial landings (pounds)




Total commercial revenue



Number of commercial vessels



Average revenue per vessel (\$USD2025)




Ecosystem Indicators


Mean Latitude

From DisMAP (1974-2024): “Shifts in the distributions are measured as change in latitude and change in depth. Capturing these shifts of particular stocks can have significant impacts on management decisions, particularly in a changing environment. For this module, data is harnessed from regional surveys* and filtered to include taxa that are sampled in at least 5% of tows in at least 3/4th of all survey years”

Survey Catch in Fall



Survey Catch in Spring




Bottom temperature

Water temperatures control spawning; optimum spawning temperatures range between 3-6°C but plaice can tolerate -0.5 to 13°C. Spawning occurs from March - June, peaking in April/May. Distribution appears to be regulated by warm summer and fall temperatures.

From the 2022 ESP: “Bottom temperature anomaly is a significant factor influencing depth of occurrence, with plaice shifting deeper and a more northward distribution with increasing bottom temperature. Mean latitude (both fall and spring) has gradually shifted northward since the late 1990s and bottom temperature anomaly was found to have a significant positive influence on the center of gravity of plaice in both fall and spring distribution models. Bottom temperature exhibited a significant, positive relationship on weight-at-age of plaice at age one (fall model), however, there was a negative relationship between WAA and bottom temperature at older ages when significant.”

SEASONAL BOTTOM TEMP (MAR - AUG)


As described in the 2022 working paper, March-August bottom temperature anomaly affects plaice recruitment. Much of the 2010s were warmer than the long term mean, with seasonal highs greater than 8°C and reaching 9°C in 2020, higher than the optimal spawning temperatures. The last few years have been around or below the long term mean, which may be beneficial for plaice recruitment.



ANNUAL MEAN PRIOR TO FALL SURVEY = OCT(year-1) - SEPT


Annual bottom temp anomaly leading up to fall survey (October of year-1 to September of year 0) is correlated with the mean latitude where plaice are found (fall and spring) and fall condition. There is a long-term increasing trend in annual bottom temperatures (Oct-Sept) since 1970, with highs above the long-term mean throughout much of the 2010s and early 2020s.

Since 2023, annual mean temperatures prior to the fall survey have declined from recent highs and are near the long-term mean. An overall warming trend in bottom temperatures may be contributing to the northward shift in plaice distribution and changes in fish condition.



ANNUAL MEAN PRIOR TO SPRING SURVEY = MAR(year-1) - FEB

Annual bottom temp anomaly leading up to spring survey (March-February) is correlated with the mean latitude where plaice are found (fall and spring). Annual temperatures (Mar-Feb) were above the long-term mean in the 2010s peaking 2020. The last few years have been near or below the long-term mean. Greater than average annual temperatures prior to the spring survey in the last 15 years may contribute to a northward shift in plaice distribution, however declining temperatures in 2023-2025 may suggest a shift back to lower latitudes.




Sea Surface Temperature

Sea surface temperature is an important consideration for plaice recruitment and egg and larval survival. The thermal limit for survival and incubation of American plaice eggs is 14°C.

SEASONAL = MAR - JUNE

remove 2026? or check back to see if data is updated through end of june

March-June SST anomaly affects plaice recruitment. Spring temperatures have hovered around 7-8°C since the early 1980s, however temperatures in the early 2020s exceeded the long-term mean. Seasonal temperatures have declined since 2022, which may be beneficial for plaice recruitment.



Gulf Stream Index


Currents, changes in circulation patterns, and shelf-slope dynamics are responsible for egg and larval transport.

When the strength of the Labrador Current is weak, a higher proportion of warm water coming from the Gulf Stream enters the Gulf of Maine, increasing Gulf of Maine water temperatures. The position of the Gulf Stream has shifted northward and the Gulf of Maine has seen temperature warming rates much higher than that of the global average in recent decades.

From the working paper: “GSI was a significant driver of recruitment in the fall. GSI explained more deviance in weight-at-age trends at younger ages and tended to exhibit a positive relationship on WAA.”

ANNUAL MEAN GULF STREAM POSITION PRIOR TO FALL SURVEY = OCT(year-1) - SEPT

The annual mean of the Gulf Stream position leading up to fall survey affects recruitment. There is an overall long-term increasing trend in the position of the Gulf Stream since 1970, with the Gulf Stream being further north than the long-term mean in the 2010s, which may be contributing to warmer water temperatures in the Gulf of Maine and affecting plaice recruitment. The Gulf Stream has been near to or south of the long-term mean since 2023.




NAO Index

From working paper: “Reproductive rate (recruits per spawner) of American plaice is significantly related to temperature and the North Atlantic Oscillation, where greatest recruitment rates occur at extreme cold temperatures. NAO has a significant effect on depth of occurrence in the fall.”

NAO 2 YEAR LAG

The North Atlantic Oscillation (NAO) index is a measure of the strength and position of the westerly winds across the North Atlantic. Strong positive phases of the NAO tend to be associated with above-normal temperatures in the eastern United States. The NAO index has a 1-2 year lag effect on American plaice recruitment. The NAO index has remained near the long-term mean throughout much of the time series (since 1970), with a few years of extreme negative NAO values early 2010s and extreme positive NAO values in 2020.

NAO (6 month mean) was significantly related to mean latitude in the spring.



AMO Index

From working paper:


- AMO was the most prevalent significant variable between both the condition index and WAA analyses, higher AMO = better condition and weight at age. Recruits per spawner increased with increasing AMO values up to a point, but then declined.
- AMO explained the most deviance in both fall and spring recruitment models.
- AMO was the most consistent significant variable across the age classes with a tendency for more variance explained in deviance of weight-at-age in older ages and a negative relationship with increasing AMO.
- AMO correlated with mean latitude in fall bottom trawl surveys

AMO has been greater than +1 SD of the long term mean since 2020, indicating possible better condition and recruitment for plaice.

AMO (short, unsmoothed) data from <https://psl.noaa.gov/data/correlation/amon.us.data>. Last updated January 2023.


AMO 6 months prior to fall survey (April-September)

6 months AMO prior to fall survey (Apr-Sept) affects recruitment and fall condition. There is a long-term increasing trend in the AMO index (Apr-Sept), with recent peaks in the early 2020s. The AMO index has been above the long-term mean since 2020, which may be contributing to better plaice condition and recruitment, or AMO may have crossed the tipping point for plaice and could be negatively impacting recruitment.



AMO 6 months prior to spring survey (September - February)


6 months AMO prior to spring survey (Sept-Feb) affects recruitment. There is a long-term increasing trend in the AMO index (Apr-Sept), with recent peaks in the 2023 and 2024. The AMO index has been above the long-term mean since 2020, which may be contributing to higher recruitment, or AMO may have crossed the tipping point for plaice and could be negatively impacting recruitment.



Condition

Condition of American plaice exhibited lows in the 2000s to 2010, indicating thinner, lower-quality fish, especially in the Gulf of Maine. Condition values have increased overall since 2010, with most fish on Georges Bank in above average or good condition. Condition of American plaice in the Gulf of Maine and Scotian Shelf vary, but are experiencing fewer years of poor or below-average condition compared to lows in the 2000s.

From ESP: “SSB had the most significant influence on mean condition, followed by annual bottom temperature anomaly, and a 6-month mean AMO variable. Plaice condition increased with increasing SSB and bottom temperature, but exhibited a curvilinear response to AMO.”



Key Takeaways

- The fishery is consolidating into fewer vessels, with fewer landings and lower revenue, but higher revenue per vessel.
- Recent cooling in bottom temperatures may be beneficial for plaice recruitment, but it is unclear how long this cooling will last.
- Warmer temperatures in the 2010s and early 2020s may have contributed to northward shifts in distribution and lower condition.
- A positive trend in AMO may result in lower weight-at-age, a northward shift in distribution in spring, and lower recruits per spawner.