



Started on	Tuesday, 14 October 2025, 7:27 PM
State	Finished
Completed on	Tuesday, 14 October 2025, 7:37 PM
Time taken	9 mins 24 secs
Marks	1.00/1.00
Grade	4.00 out of 4.00 (100%)

Given an array A of sorted integers and another non negative integer k, find if there exists 2 indices i and j such that $A[j] - A[i] = k$, $i \neq j$.

Input Format:

First Line n - Number of elements in an array

Next n Lines - N elements in the array

k - Non - Negative Integer

Output Format:

1 - If pair exists

0 - If no pair exists

Explanation for the given Sample Testcase:

YES as $5 - 1 = 4$

So Return 1.

For example:

Input	Result
3 1 3 5 4	1

Answer: (penalty regime: 0 %)

```

1  #include <stdio.h>
2
3  int main() {
4      int n, i, j, k;
5      scanf("%d", &n);
6      int A[n];
7      for(i = 0; i < n; i++) {
8          scanf("%d", &A[i]);
9      }
10     scanf("%d", &k);
11
12     for(i = 0; i < n; i++) {
13         for(j = 0; j < n; j++) {
14             if(i != j && A[j] - A[i] == k) {
15                 printf("1\n");
16                 return 0;
17             }
18         }
19     }
20     printf("0\n");
21     return 0;
22 }
23

```

	Input	Expected	Got	
✓	3 1 3 5 4	1	1	✓

	Input	Expected	Got	
✓	10 1 4 6 8 12 14 15 20 21 25 1	1	1	✓
✓	10 1 2 3 5 11 14 16 24 28 29 0	0	0	✓
✓	10 0 2 3 7 13 14 15 20 24 25 10	1	1	✓

Passed all tests! ✓

Correct

Marks for this submission: 1.00/1.00.

[Back to Course](#)