

MULTIVARIATE STATISTICS

Stats and R workshop 5th – 6th March 2018

Susan Jarvis

What is multivariate statistics?

- Univariate statistics – **single response**
- Multivariate statistics – **multiple responses**

What is multivariate statistics?

- Univariate statistics – **single response**
- Multivariate statistics – **multiple responses**



What does multivariate statistics do?

- Summarise multivariate data into a smaller number of variables that represent the main sources of variation in the data (ordination)

Site	Sp1	Sp2	Sp3	Sp4	Sp5
1	0	4	3	5	1
2	0	0	0	3	0
3	1	1	2	2	0
4	2	2	0	2	1
5	6	0	4	0	8
6	0	0	6	0	2




Site	Axis 1	Axis 2
1	-1.04	0.14
2	-1.12	-0.70
3	-0.65	0.13
4	-0.51	-1.48
5	2.87	-0.91
6	0.46	2.82

What does multivariate statistics do?

- Summarise multivariate data into a smaller number of variables that represent the main sources of variation in the data (ordination)


Site	Sp1	Sp2	Sp3	Sp4	Sp5
1	0	4	3	5	1
2	0	0	0	3	0
3	1	1	2	2	0
4	2	2	0	2	1
5	6	0	4	0	8
6	0	0	6	0	2



What does multivariate statistics do?

- Find groups of samples that share similar properties based on multiple variables (e.g. cluster analysis)

Site	Sp1	Sp2	Sp3	Sp4	Sp5
1	0	4	3	5	1
2	0	0	0	3	0
3	1	1	2	2	0
4	2	2	0	2	1
5	6	0	4	0	8
6	0	0	6	0	2



```
dis  
hclust(*,"average")
```

What does multivariate statistics do?

- Test for differences between groups of samples based on multiple variables (e.g. MANOVA)


Site	Sp1	Sp2	Sp3	Sp4	Sp5	Group
1	0	4	3	5	1	A
2	0	0	0	3	0	A
3	1	1	2	2	0	A
4	2	2	0	2	1	A
5	6	0	4	0	8	B
6	0	0	6	0	2	B

	Df	SS	MS	F	R2	P
Group	1	0.65	0.65	4.61	0.53	0.06
Residuals	4	0.56	0.14		0.46	
Total	5	1.21				

What does multivariate statistics do?

- Test whether patterns in variation are explained by independent variables (constrained ordination)


Site	Sp1	Sp2	Sp3	Sp4	Sp5	V1	V2
1	0	4	3	5	1	4.5	1.2
2	0	0	0	3	0	2.4	1.1
3	1	1	2	2	0	5.3	2.1
4	2	2	0	2	1	2.4	1.4
5	6	0	4	0	8	2.5	1.3
6	0	0	6	0	2	5.5	0.8



What does multivariate statistics do?

- Test whether patterns in variation are explained by independent variables (constrained ordination)

Site	Sp1	Sp2	Sp3	Sp4	Sp5	V1	V2
1	0	4	3	5	1	4.5	1.2
2	0	0	0	3	0	2.4	1.1
3	1	1	2	2	0	5.3	2.1
4	2	2	0	2	1	2.4	1.4
5	6	0	4	0	8	2.5	1.3
6	0	0	6	0	2	5.5	0.8



Practical

- Use R package 'vegan'
- Includes:
 - Ordination
 - Constrained ordination
 - MANOVA with dissimilarities
 - Clustering methods