an open access 🔓 journal

SCIENCE



Citation: Betzel, R. F., Fukushima, M., He, Ye, Zuo, Xi-Nian, Sporns, O. (2016) Dynamic fluctuations coincide with periods of high and low modularity in resting-state functional brain networks Network Neuroscience, 1

DOI:

http://dx.doi.org/10.1162/NETN-00001

Supporting Information: http://dx.doi.org/10.7910/DVN/PQ6ILM

Received: 20 October 2016 Accepted: 7 November 2016 Published: 26 January 2016

Competing Interests: The authors have declared that no competing interests exist.

Corresponding Author: George Chacko netelabs@nete.com

Handling Editor: Xi-Nian Zuo

Copyright: © 2019

Massachusetts Institute of Technology Published under a Creative Commons Attribution 4.0



RESEARCH ARTICLE

Co-citations in context: disciplinary heterogeneity is relevant

James Bradley¹, Sitaram Devarakonda², Avon Davey², Dmitriy Korobskiy², Siyu Liu², Tandy Warnow³ and George Chacko²

¹Raymond A. Mason School of Business, College of William and Mary, Williamsburg, VA, USA
²Netelabs, NET Esolutions Corporation, McLean, VA 22102, USA
³Department of Computer Science, University of Illinois at Urbana-Champaign, Champaign, IL 61820, USA

Keywords: (bibliometrics, co-citation analysis, random graphs)

ABSTRACT

Citation analysis of the scientific literature has been used to study and define disciplinary boundaries, to trace the dissemination of knowledge, and to estimate impact. Co-citation, the frequency with which pairs of publications are cited, provides insight into how documents relate to each other and across fields. Co-citation analysis has been used to characterize combinations of prior work as conventional or innovative and to derive features of highly cited publications. Given the organization of science into disciplines, a key question is the sensitivity of such analyses to frame of reference. Our study examines this question using semantically-themed citation networks. We observe that trends reported to be true across the scientific literature do not hold for focused citation networks, and we conclude that co-citation analysis requires a contextual perspective.

INTRODUCTION

Citation and network analysis of scientific literature reveals information on semantic relationships between publications, collaboration between scientists, and the practice of citation itself (de Solla Price, 1965; Garfield, 1955; Newman, 2001; Patience, Patience, Blais, & Bertrand, 2017; Shi, Leskovec, & McFarland, 2010). Co-citation, the frequency with which two documents are cited together in other documents provides additional insights, including the identification of semantically related documents, fields, ideas, and specializations in science (Boyack & Klavans, 2010; Marshakova-Shaikevich, 1973; Small, 1973; ?).

(Uzzi, Mukherjee, Stringer, & Jones, 2013) used a novel approach for co-citation analysis of 17.9 million articles and their cited references from the Web of Science (WoS) to characterize a subset of highly cited articles with respect to both novel and conventional combinations of prior research. Observed co-citation frequencies, mapped to journals, were computed across the dataset by Monte Carlo simulation under a random graph model and normalized (shifted and scaled) to expected values. These normalized journal pair frequencies were termed *z-scores* (Materials and Methods). Thus, every article was associated with multiple *z-scores* corresponding with the journal pairs represented in its citations. For each article, positional statistics of these *z-scores* were calculated to describe conventionality: high conventionality (HC) if the median *z-score* for an article was greater than the median of median *z-scores* of all articles and low conventionality for the converse (LC). Similarly,

an article was deemed to have high novelty (HN) if the tenth percentile of its z-scores was negative and low novelty (LN) for the converse. Accordingly, each article was labeled with respect to conventionality and novelty, e.g, HCHN (denoting that the article exhibits high conventionality and high novelty), with all four combinations being possible. (Uzzi et al., 2013) observed that HCHN articles were twice as likely to be highly cited, suggesting that novel combinations of ideas flavoring a body of conventional thought were a recipe for impact across the scientific literature.

Key to Uzzi et al., however, is their random graph model and its underlying assumptions. A careful examination of their Monte Carlo simulation reveals that random substitutions of references used to generate expected journal pair frequencies are equiprobable, which means that a reference in an article can be substituted by any other reference randomly chosen from those published in the same year, irrespective of subject matter, disciplinary focus and citation count. For example, the model permits substitution of a reference in quantum physics with equal probability by a reference in quantum biology (i.e., the same field) or in some entirely different field, such as classical literature, evolutionary biology, or anthropology. Such substitutions poorly model documented citation behavior (Garfield, 1979; Klavans & Boyack, 2017; Moed, 2010; Wallace, Lariviere, & Gingras, 2012). In addition, under this random model, a reference cited over 100 times in a given year is selected with the same probability as a reference cited only once, which is inconsistent with the actual frequency of citation and also with the power law or lognormal citation distributions observed in the literature (Perline, 2005; Stringer, Sales-Pardo, & Amaral, 2010). Lastly, as we will show here, while the citation switching algorithm used in Uzzi et al. preserves the number of publications and the number of references in each publication, it does not preserve the proportion of cited references from within the disciplinary network. Accordingly, the expected value calculations generated by the Uzzi et al. (2013) simulations and used in characterizing journal pairs in terms of conventionality and novelty can be reasonably questioned on grounds of model mis-specification.

A follow-up study by (Boyack & Klavans, 2014) explored the impact of discipline and journal effects on the definition of conventionality and novelty. While their study had some methodological differences from (Uzzi et al., 2013) in the use of Scopus data and a χ^2 calculation for the expected number of journal pair citations, Boyack and Klavans find the same basic trend that HCHN is more probably in highly cited papers. However, they note that "only 64.4% of 243 WoS subject categories" in the Uzzi *et al.* study met the criterion of having the highest probability of hit papers in the HCHN category. Further, they observed that journals vary widely in terms of size and influence and that 20 journals accounted for nearly 15% of co-citations in their measurements. Lastly, they noted that three multidisciplinary journals accounted for 9.4% of all atypical combinations, suggesting strong effects from both disciplines and journals that were not reported by Uzzi et al.

Despite different methods used to generate expected values, both (Uzzi et al., 2013) and (Boyack & Klavans, 2014) measured observed co-citation frequencies across the scientific literature without disciplinary constraints and subsequently used normalized frequencies to examine disciplinary subsets. In contrast, we chose to first construct semantically related sets of documents (disciplinary networks) and to measure observed and expected frequencies within these networks in order to preserve the proportion of citations from a citing document's subject area under random sampling.

We hypothesized that assessing articles' novelty, conventionality, and impact on datasets focused in disciplinary areas would reduce model mis-specification and better simulate citation practice, in turn, leading to different and more relevant conclusions. Accordingly, we used keyword searches of the scientific literature to create three citation networks themed around academic disciplines. Within these disciplinary frameworks, we calculated expected journal pair frequencies using a random graph model that accounted for existing citation frequencies and an efficient Monte Carlo simulation algorithm that permitted a substantially greater number of simulations than do previous studies. We found our model did, in fact, preserve the citation patterns found in the literature whereas the approach of Uzzi et al. does not. The varying citation patterns when sampling from a disciplinaryfocused dataset versus a broader dataset due, was a cause of journal pair z-scores changing signs in 28.6% of instances: the effect of any one of these journal pairs on an article's novelty or conventiality is contradictory between broad and, more narrow disciplinary datasets. We contend that the interpretation due to a disciplinary dataset, which preserved citation patterns, is more appropriate and an improvement on current methods. Furthermore, our analysis of these semantically-themed citation networks challenges the universality of articles in the HCHN category having the highest probability of being hits as found in Uzzi et al. (2013). For example, while HC remains highly correlated with hit articles in immunology and metabolism research, HN is not. In addition, while HN is highly correlated with hit articles in applied physics, HC is not. Furthermore, we found that the categories demonstrating the highest percentage of hits is not robust with respect to varying parameters that define hit articles and the threshold for highly novel citation patterns.

MATERIALS AND METHODS

Bibliographic data We have previously developed ERNIE, an open source knowledge platform into which we parse the Web of Science (WoS) Core Collection (Keserci, Davey, Pico, Korobskiy, & Chacko, 2018). WoS data stored in ERNIE spans the period 1900-2019 and consists of over 72 million publications. For this study, we generated an analytical dataset from years 1985 to 2005 from ERNIE. The total number of publications in this dataset was just over 25 million publications (25,134,073). For each of these years, we further restricted analysis to those of type Article. Since WoS data also contains incomplete references or references that point at other indexes, we also considered only those references for which there were complete records (Table 1). For example, WoS data for year 2005 contained 1,753,174 publications, which after restricting to type Article and considering only those references described above resulted in 916,573 publications, 6,095,594 unique references (set of references), and 17,167,347 total references (multiset of references). Given consistent trends in the data, we analyzed the two boundary years (1985 and 2005) and the mid-point (1995) performing 1,000 simulations for each dataset.

Disciplinary datasets We constructed three disciplinary datasets based on keyword searches. (i) immunology (ii) metabolism (iii) applied physics. For the first two, rooted in biomedical research, we searched Pubmed for the term 'immunology' or 'metabolism' in the years 1985, 1995, and 2005. Pubmed IDs (pmids) returned were matched to WoS IDs (wos_ids) and used to retrieve relevant articles. For the applied physics dataset, we directly searched subject labels in WoS for 'applied physics'.

Normalization of observed and expected values Building upon prior work Uzzi et al. (2013), all $\binom{n}{2}$ reference pairs were generated for each publication, where n is the number of cited references in the publication. These reference pairs were then mapped to the journals they were published in using ISSN numbers to create journal pairs. Where multiple ISSN numbers ex-

ist for a journal, the most frequently used one in the WoS was assigned to the journal. In addition, publications containing fewer than two references were discarded. Journal pair frequencies were summed across the dataset to create observed frequencies (F_{obs}). In contrast to the preceding study (Uzzi et al., 2013), we generated 1,000 rather than 10 null models for each dataset by randomly shuffling references while preserving the number of publications, the number of references in each publication, and the frequency with which these references were cited within the year of interest. In contrast to the preceding study (Uzzi et al., 2013), we generated 1,000 rather than 10 null models for each dataset. Expected values (F_{exp}) for journal pairs were generated by averaging the result of 1,000 simulations. z-scores were calculated for each journal-pair using the formula $(F_{obs} - F_{exp})/\sigma$ where σ is the standard deviation of the frequencies generated by simulation. As a result of these calculations, each publication becomes associated with a set of z-scores corresponding to the journal pairs derived from pairwise combinations of its cited references and positional statistics (quantiles) of z-scores were calculated for each publication. Publications were subsequently labeled according to conventionality and novelty: (i) HC if the median z-score exceeded the median of median z-scores for all publications and LC if the median z-score was equal to or less than the median of median z-scores for all publications, and (ii) HN if the tenth percentile of z-scores for a publication was less than zero, and LN if the tenth percentile of z-scores for a publication was greater than zero. (Should we comment here that while we are using the term z-score, reusing it from Uzzi, we do not imply that these are related to a normal distribution?)

We investigated multiple definitions of hit articles, as did Uzzi et al., with hits defined as the 1%, 2%, 5%, and 10% top-cited articles. Also, mention different thresholds, 1st and 10th, for novelty threshold.

RESULTS

Chacko

In our study, we also use a Monte Carlo approach to simulate under a random graph model. A principal consideration, however, was to restrict model misspecification arising from disciplinarily irrelevant references. We addressed this consideration by restricting random substitutions to only those references in the disciplinary network being studied. We estimated model misspecification by measuring the Kullback-Leibler (K-L) Divergence (?) between observed and simulated frequencies for the set of journals common to both a disciplinary network and the WoS superset (Table 3). The results indicate that simulations under our model consistently have a lower K-L divergence compared to simulations that draw from the WoS superset and its attendant substitutions that are ectopic with respect to field and discipline (Introduction).

Bradley

Figure 2 shows that the z-scores for the same journal pair can be positive (negative) when computed with respect to one data set but be negative (positive) for another data set. The journal-pair z-scores in Figure 2 have consistent signs for both Immunology and WoS data sets in 71.4% of the instances and different signs for 28.6% of the journal pairs. A negative journal pair z-score in one context increases the likelihood that articles citing it will be deemed to be novel while articles citing it in another context, where the journal pair z-score is positive, are less likely to be classified as novel and possibly more likely to be classified as conventional. Figure 2 reflects that the WoS data set has approximately 44,000 fewer

Table 1. Summary of WoS Analytical Dataset. UP: unique publications, UR: unique references, TR: total references. The number of publications, unique references, total references and the ratio of total references to unique references increases monotonically with each year indicating that both the number of documents and citation activity increase over time. Data for reference years is flanked by horizontal lines and shown in boldface. Only publications of type Article and references with complete WoS records are included in these counts.

| Year | UP | UR | TR | TR/UR |
|------|--------|---------|----------|-------|
| 1985 | 418495 | 2281297 | 5615496 | 2.46 |
| 1986 | 402309 | 2316451 | 5708796 | 2.46 |
| 1987 | 412936 | 2427347 | 5998513 | 2.47 |
| 1988 | 426001 | 2545647 | 6354917 | 2.50 |
| 1989 | 443144 | 2673092 | 6749319 | 2.52 |
| 1990 | 458768 | 2827517 | 7209413 | 2.55 |
| 1991 | 477712 | 2977784 | 7729776 | 2.60 |
| 1992 | 492181 | 3134109 | 8188940 | 2.61 |
| 1993 | 504488 | 3278102 | 8676583 | 2.65 |
| 1994 | 523660 | 3458072 | 9255748 | 2.68 |
| 1995 | 559685 | 3692575 | 9897946 | 2.68 |
| 1996 | 663110 | 4144581 | 11641286 | 2.81 |
| 1997 | 677077 | 4340733 | 12135104 | 2.80 |
| 1998 | 693531 | 4573584 | 12728629 | 2.78 |
| 1999 | 709827 | 4784024 | 13280828 | 2.78 |
| 2000 | 721926 | 5008842 | 13810746 | 2.76 |
| 2001 | 727816 | 5203078 | 14261189 | 2.74 |
| 2002 | 747287 | 5464045 | 15001390 | 2.75 |
| 2003 | 786284 | 5773756 | 16024652 | 2.78 |
| 2004 | 826834 | 6095594 | 17167347 | 2.82 |
| 2005 | 916573 | 6629595 | 19066249 | 2.88 |
| | | | | |

Table 2. Disciplinary Datasets. PubMed and WoS were searched for articles using search terms, 'immunology', 'metabolism', and 'applied physics'. Counts of retrieved publications are shown for each of the three years analyzed.

| Year | Immunology | Metabolism | Applied Physics |
|------|------------|------------|-----------------|
| 1985 | 21,606 | 78,998 | 10,298 |
| 1995 | 29,320 | 121,247 | 21,012 |
| 2005 | 37,296 | 200,052 | 35,600 |

negative z-scores than does the immunology data set, which contributes to its significantly lower percentage of high-novelty articles as shown in Figure 2.

Figure 3, Panels (a) and (b), compares hit rates for the four categories among the Immunology, Applied Physics, and WoS datasets for 1995: the hit rate is defined as the number of hit articles in each category divided by the number of articles in the category. We evaluated the statistical significance of the categorical hit rates using multiple methods, some of which we describe here. Our first test was based on the null hypotheses that hits were distributed randomly among the four categories with uniform probability in proportion to the number of articles in each category. Using a Chi-Square Goodness of Fit test, rejecting the null hypothesis in favor of the alternate hypothesis supports a non-uniform dispersion of hits: that is, some of the four categories are individually associated with higher than




Figure 1. Fraction of publications with high conventionality (HC) and high novelty (HN) signatures relative to citation count.




Figure 2. Journal pair z-scores vary with reference dataset. Scatter plot points (319,005) indicate journal pair z-scores for the 1995 Immunology dataset along the x-axis and the 1995 Web of Science dataset on the y-axis. Black indicates journal pairs whose z-scores have the same sign when computed for both reference datasets while red points indicate the 28.6% of journal pairs whose z-scores change sign across datasets. Regions with deeper hues indicate higher point densities and the histograms show the marginal distributions for each dataset separately.




Figure 3. Effect of Context on Journal-Pair z-scores and Categorical Hit Rates: Immunology, Applied Physics, and WoS for 1995. Panels (a) and (b) show hit rates for the LNLC, LNHC, HNLC, and HNHC categories for the immunology, applied physics, and WoS datasets when hit articles are defined as the top 1% and top 10% of articles, respectively. Novelty in both panels is defined at the 10th percentile of articles' z-score distributions. The number of data points in the immunology, applied physics, and WoS data sets are 21,917, 18,305, and 476,288, respectively. The results for the WoS data set mirrored previous results from Uzzi et al. Uzzi et al. (2013) where the highest hit rate was for the HNHC category. The highest hit rates for the immunology dataset, in contrast, are in the LNHC category. The LNHC category often had the highest hit rate for the immunology dataset for various parameter settings with the HCHN category having the highest hit rate in other cases: metabolism results are similar. The applied physics dataset shows further contrast as the hit rate for it was highest in the HNLC category. The HNLC category demonstrated the highest hit rate often for applied physics: otherwise the highest hit rate was for HNHC articles.

Table 3. Measuring Model Misspecification. For the set of journal pairs in common between a disciplinary network and the full WoS dataset, Kullback-Leibler (K-L) divergences between empirical and simulated journal pair frequencies were computed for the years 1985, 1995, and 2005 for the three disciplinary datasets (applied_physics, immunology, and metabolism) using either the disciplinary network as background or the WoS superset (all_wos) to generate the null model (Background). K-L divergence was calculated using the R seewave package with a base (logarithm) of 2. The ratio between the K-L divergence for disciplinary networks versus the full WoS ranges from 1.96 to 2.77 and is greater than 2.0 for eight out of nine cases, strongly suggesting that simulations that constrain substitutions to the given disciplinary network better model the observed data.

| | Disciplinary Network | Year | Background | K-L Divergence | Ratio |
|----|----------------------|------|--------------|----------------|-------|
| 1 | appl_physics | 1985 | appl_physics | 1.21 | |
| 2 | | 1985 | all_wos | 2.37 | 1.96 |
| 3 | | 1995 | appl_physics | 0.86 | |
| 4 | | 1995 | all_wos | 2.37 | 2.77 |
| 5 | | 2005 | appl_physics | 0.95 | |
| 6 | | 2005 | all_wos | 2.35 | 2.47 |
| 7 | immunology | 1985 | immunology | 0.75 | |
| 8 | | 1985 | all_wos | 1.68 | 2.24 |
| 9 | | 1995 | immunology | 0.78 | |
| 10 | | 1995 | all_wos | 1.70 | 2.19 |
| 11 | | 2005 | immunology | 0.73 | |
| 12 | | 2005 | all_wos | 1.92 | 2.63 |
| 13 | metabolism | 1985 | metabolism | 1.11 | |
| 14 | | 1985 | all_wos | 2.24 | 2.02 |
| 15 | | 1995 | metabolism | 1.07 | |
| 16 | | 1995 | all_wos | 2.33 | 2.17 |
| 17 | | 2005 | metabolism | 1.19 | |
| 18 | | 2005 | all_wos | 2.60 | 2.18 |

expected, or lower than expected hit rates. The null hypothesis was rejected at a p < 0.001 in all cases in Figure 3, with the exception of the immunology and applied datasets where hit articles are designated as the top 1% of articles: valid tests were not possible in those instances due to too few expected hits. The null hypothesis was rejected with p < 0.001 for all valid tests for all parameter settings, all datasets, and all years: hypotheses tests were valid in 73 of 96 instances. We conclude that it is likely that the distribution of hits among categories is not uniform but, rather, hit rates vary among the categories in all datasets. (Should we insert a table with all statistical results?)

We computed hit rates for the WoS dataset, which mirrored Uzzi et al.'s results whereby the largest hit rates were for the HNHC category, despite our methodological improvement of sampling citations in proportion to their frequency. We found contrary results in the 1995 immunology dataset where articles in the LNHC category often had the greatest hit rates, as reflected in Table 4 and Figure 3. Across all year's data, all datasets, and all parameter settings, the highest hit rates in the immunology datasets were sometimes in the LNHC category and sometimes in the HNHC category. The metabolism hit rates reflected this same pattern. The greatest hit rates for the applied physics data were often in the HNLC category as reflected in Table 4 and Figure 3, and otherwise in the HNHC category. We conclude that Uzzi et al.'s finding of high hit rates in the HNHC category does not hold generally for disciplinary-based datasets and that novel citation patterns are not always

Table 4. Hit Rates by Category. The last four columns indicate the proportion of publications that are hits for each respective category.

| | Hits as % | Novelty | | | | |
|----------|-------------|------------|-------|-------|-------|-------|
| Data Set | of Articles | Percentile | LNLC | LNHC | HNLC | HNHC |
| Imm95 | 1% | 10% | 0.000 | 0.019 | 0.005 | 0.014 |
| Imm95 | 10% | 10% | 0.017 | 0.128 | 0.076 | 0.129 |
| Metab95 | 1% | 10% | 0.001 | 0.017 | 0.006 | 0.014 |
| Metab95 | 10% | 10% | 0.019 | 0.130 | 0.074 | 0.133 |
| AP95 | 1% | 10% | 0.002 | 0.007 | 0.012 | 0.010 |
| AP95 | 10% | 10% | 0.047 | 0.079 | 0.123 | 0.109 |
| WoS95 | 1% | 10% | 0.004 | 0.013 | 0.009 | 0.017 |
| WoS95 | 10% | 10% | 0.056 | 0.115 | 0.104 | 0.156 |

indicative of impactful research, as was the case with immunology. Furthermore, the categories displaying the greatest hit rate vary with parameter settings and with the year. The lack of stable results across parameter settings suggests that parameters must be selected judiciously.

We also tested the explanatory power of each framework dimension by classifying articles as Low Novelty (LN) or High Novelty (HN) and, separately, as having Low Conventionality (LC) or High Conventionality (HC). We tested the null hypothesis that hits are distributed between LN and HN (LC and HC) in proportion to the total number of articles assigned to those categories. That null hypothesis was rejected for the WoS data along both dimensions. Consistent with previous analysis, hit articles were overrepresented in the HC category in every instance of WoS data at a p < 0.001 and hit articles were overrepresented in the HN category at a p < 0.001 in all but two cases. The p-values, in those cases, were 0.002 and 0.007. Hits in the immunology and metabolism data were overrepresented in the HC category with the same statistical significance as for WoS. The relationship of novelty with hits in the immunology and metabolism data differed dramatically from the WoS, however, with statistically significant findings of hit articles being sometimes overrepresented in the LN category, and sometimes being underrepresented. Of the 12 tests for applied physics, the statistical significance supporting a positive relationship between hit articles and HN were all p < 0.10, and 10 of 12 were p < 0.05. These tests also indicated strong support for the relationship between LN and hit articles in applied physics in a limited number of tests, with p < 0.10 in 5 of 12 instances and p < 0.05 in 3 of 12 instances. These results suggest that (1) both conventionality and novelty are strongly related to hits in the WoS, (2) the conventionality dimension is strongly related with hits in immunology and metabolism and novelty is not, (3) novelty is more strongly related with hits in applied physics than is conventionality. More generally, we find that the dimensions most strongly related with hit articles vary across disciplines and between disciplinary and broad data sets.

DISCUSSION

We conclude that Uzzi et al.'s finding of high hit rates in the HNHC category does not hold generally for all datasets and that novel citation patterns are not always indicative of impactful research.

The z-scores that change sign across datasets is either contradictory or an acceptable variation due to the difference in reference sets. We contend that it is the former because the inappropriate substitution of citations with those from disciplines that are implausible and because the observed citation frequencies are ignored. That fewer z-scores are negative in the WoS dataset relative to the immunology dataset may be directly due to the uniform sampling of references whereby many resulting journal pairs are never observed in the literature and so that the expected frequencies of those observed pairs have lower expected values, thus biasing their z-scores upward. In addition, treating citation as a set versus a multiset means that a random model will sample popular citations downward, thus further increasing their z-scores.

Contrary to Uzzi et al. we found the category displaying the highest hit rate to be sensitive to the experimental parameter settings, which included the percentage of articles deemed to be hits and the percentile of articles' z-score distributions that delineated between articles of low novelty and high novelty. Thus, with this lack of robustness, studies are faced with the necessity of defining which parameters are the best parameters to classify articles.

ACKNOWLEDGMENTS

We are grateful to Kevin Boyack and Dick Klavans for constructively critical discussions. We thank the authors of Uzzi et al. (2013) for sharing their Python simulation code. Research and development reported in this publication was partially supported by Federal funds from the National Institute on Drug Abuse, National Institutes of Health, US Department of Health and Human Services, under Contract Nos. HHSN271201700053C (N43DA-17-1216) and HHSN271201800040C (N44DA-18-1216). The content of this publication is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. TW receives funding from the Grainger Foundation. All the code used in this study is freely available from a Github site. Access to the bibliographic data analyzed in this study requires a license from Clarivate Analytics, which had no role in funding, experimental design, review of results, and conclusions presented.

AUTHOR CONTRIBUTIONS

This study was designed by GC, JB, SD, and TW. Simulations and analysis were performed by AD, GC, JB, and SD. Infrastructure and workflows used to generate data used in this study were developed by AD, DK, SL, SD, and GC. All authors reviewed and commented on the manuscript, which was written by GC, JB, and TW.

REFERENCES

- Boyack, K., & Klavans, R. (2010).Co-citation analysis, bibliographic coupling, and direct citation: Which citation approach represents the research front most accu-Journal of the American Society for Information Science and Technology, 61(12), 2389-2404. Retrieved 2019-01-05, from https://onlinelibrary.wiley.com/doi/ abs/10.1002/asi.21419 doi: 10.1002/asi.21419
- Boyack, K., & Klavans, R. (2014). Atypical combinations are confounded by disciplinary effects. In International conference on science and technology indicators (pp. 49-58). Leiden, Netherlands: CWTS-Leiden University.
- de Solla Price, D. J. (1965).Networks of Scientific Science, 149(3683), 510–515. Papers. Retrieved 2019-02-03, from http://www.sciencemag.org/cgi/doi/10 .1126/science.149.3683.510 doi: 10.1126/science.149.3683.510
- Garfield, E. Citation Indexes for Science: A New Dimension in Documentation through Association of Science, 122(3159), 108-111. Retrieved 2019-05-16, from https://science.sciencemag.org/content/ 122/3159/108 doi: 10.1126/science.122.3159.108
- Garfield, E. (1979). Citation Indexing-Its Theory and Application in Science, Technology, and Humanities (1st ed.). The address: John Wiley and Sons, ISI Press. (An optional note)
- Keserci, S., Davey, A., Pico, A. R., Korobskiy, D., & Chacko, G. (2018). ERNIE: A data platform for research assessment. doi: 10.1101/371955
- Klavans, R., & Boyack, K. W. (2017). Research portfolio analysis and topic prominence. Journal of Informetrics, 11(4), 1158–1174. Retrieved 2019-05-16, from http://www.sciencedirect .com/science/article/pii/S1751157717302110 doi: 10.1016/j.joi.2017.10.002
- Marshakova-Shaikevich, I. (1973). System of document connections based on references. Nauch-Techn.Inform, Ser.2, 6(4), 3-8. Retrieved 2019-02-03, from http://doi.wiley.com/ 10.1002/asi.4630240406 doi: 10.1002/asi.4630240406

- Moed, H. F. (2010). Measuring contextual citation impact of scientific journals. Journal of informetrics, 4(3), 265-277.
- Newman, M. E. J. (2001). The structure of scientific collaboration networks. Proceedings of the National Academy of Sciences, 98(2), 404-409. Retrieved 2019-03-22, from https://www.pnas .org/content/98/2/404 doi: 10.1073/pnas.98.2.404
- Patience, G. S., Patience, C. A., Blais, B., & Bertrand, F. (2017). Citation analysis of scientific categories. *Heliyon*, 3(5), e00300.
- Perline, R. (2005). Strong, Weak and False Inverse Power Laws. Statistical Science, 20(1), 68-88. Retrieved 2019-05-22, from https://www.jstor.org/stable/20061161
- Shi, X., Leskovec, J., & McFarland, D. A. (2010). Citing for high impact. In Proceedings of the 10th annual joint conference on digital libraries (pp. 49-58). New York, NY, USA: ACM. Retrieved from http://doi.acm.org/10.1145/1816123 .1816131 doi: 10.1145/1816123.1816131
- Small, H. (1973). Co-citation in the scientific literature: A new measure of the relationship between two documents. Journal of the American Society for Information Science, 24(4), 265-269. Retrieved 2019-02-03, from http://doi.wiley.com/ 10.1002/asi.4630240406 doi: 10.1002/asi.4630240406
- Stringer, M. J., Sales-Pardo, M., & Amaral, L. A. N. (2010). Statistical validation of a global model for the distribution of the ultimate number of citations accrued by papers published in a scientific journal. Journal of the American Society for Information Science and Technology, 61(7), 1377-1385. Retrieved bioRxiv. (https://www.biorxiv.org/content/early/2018/07/19/371955)2019-05-22, from https://www.ncbi.nlm.nih.gov/pmc/ articles/PMC3158611/ doi: 10.1002/asi.21335
 - Uzzi, B., Mukherjee, S., Stringer, M., & Jones, B. (2013). Atypical combinations and scientific impact. Science (New York, N.Y.), 342(6157), 468-472. doi: 10.1126/science.1240474
 - Wallace, M. L., Lariviere, V., & Gingras, Y. (2012). A Small World of Citations? The Influence of Collaboration Networks on Citation Practices. PLOS One, 7, e33339. Retrieved 2019-05-16, from https://journals.plos.org/plosone/ article?id=10.1371/journal.pone.0033339 10.1371/journal.pone.0033339