

Accelerated Broiler Production & Optimization

Course Description

Broiler production requires a high-velocity management style focused on maximizing feed conversion efficiency within a tight 6-week window. This module provides a complete medication, vaccination, and support protocol designed to achieve target slaughter weights safely and efficiently.

Lesson 4.1: High-Velocity Broiler Program (Days 1-42)

DAY	MEDICATION & VACCINATION SCHEDULE	OPERATIONAL ADMINISTRATION
DAY 1	Glucose & Multivitamin Solution	Oral (First 12 hours post-placement)
DAYS 2 - 5	Prophylactic Broad-Spectrum Antibiotics	Oral (Prevent early chick mortality)
DAY 6	Multivitamin Support Solution	Oral (Stabilize immune recovery)
DAY 7	Quality Unmedicated Drinking Water	Oral (Systemic physiological flushing)
DAY 8	1st Gumboro Vaccine (IBD)	Oral (Targeted bursal defense)
DAY 9	Multivitamin Post-Vaccination Sweep	Oral (Sustained weight gain catalyst)
DAYS 10 - 13	Anticoccidial Preventive Block	Oral (Mid-brooding intestinal shield)
DAYS 14 - 15	Multivitamin Solution Block	Oral (Metabolic optimization)
DAY 16	1st LaSota Vaccine (Newcastle Disease)	Oral (Primary systemic viral barrier)
DAYS 17 - 19	Multivitamin Maintenance Line	Oral (Skeletal expansion support)
DAYS 20 - 21	Quality Unmedicated Drinking Water	Oral (Renal clearing window)
DAY 22	2nd Gumboro Vaccine (Booster)	Oral (Final IBD lockdown phase)
DAY 23	Multivitamin Stress Minimizer Course	Oral (Post-vaccinal recovery)
DAYS 24 - 26	2nd Antibiotic Prophylactic Block	Oral (Respiratory protection sweep)
DAY 27	Multivitamin Growth Formulation	Oral (Muscle mass optimization)
DAY 28	Quality Unmedicated Drinking Water	Oral (Systemic clearing)
DAY 29	2nd LaSota Vaccine (Booster)	Oral (Final Newcastle protection)
DAYS 30 - 33	High-Density Multivitamin Solution	Oral (Accelerated meat-weight finish)
DAYS 34 - 42	Pure Drinking Water / Multivitamin Only	Oral (Strict Chemical Withdrawal Phase)

Lesson 4.2: Critical Principles of Chemical Withdrawal

The final phase (Days 34–42) is the most critical period for meat safety and quality. **All chemical treatments, therapeutic antibiotics, and synthetic coccidiostats must be completely withdrawn from the water and feed.** Only plain, clean water or low-dose multivitamins should be administered.

Why Withdrawal Matters:

- Prevents chemical and antibiotic residues from remaining in the consumer meat product.
- Helps prevent the development of drug-resistant bacterial strains.
- Ensures full compliance with international food safety standards for commercial distribution.

Lesson 4.3: Maximizing Feed Conversion Efficiency (FCR)

To optimize feed efficiency within this short 42-day cycle:

- Provide fresh feed at cooler times of the day (early morning and evening) to prevent heat stress and maintain steady consumption.
- Ensure water lines are clean and cool; a reduction in water intake directly reduces feed consumption and slows growth rates.